


Hypoglycemia (Low Blood Glucose)

Causes: Too little food or skipping a meal; too much insulin or diabetes pills; more active than usual.

Some Symptoms:

Onset: Often sudden.



What Can You



CHECK your blood glucose, right away. If you can't check, treat anyway.



TREAT by eating 3 to 4 glucose tablets or 3 to 5 hard candies you can chew quickly (such as peppermints), or by drinking 4-ounces of fruit juice, or 1/2 can of regular soda pop.



CHECK your blood glucose again after 15 minutes. If it is still low, treat again. If symptoms don't stop, call your healthcare provider.

For more information, call the Novo Nordisk Tip Line at 1-800-260-3730 or visit us online at ChangingDiabetes-us.com.


Novo Nordisk Inc. grants permission to reproduce this piece for non-profit educational purposes only on condition that the piece is maintained in its original format and that the copyright notice is displayed. Novo Nordisk Inc. reserves the right to revoke this permission at any time. Concept developed by Rhonda Rogers, RN, BSN, CDE





Hyperglycemia (High Blood Glucose)

Causes: Too much food, too little insulin or diabetes pills, illness, or stress. **Onset:** Often starts slowly.



HIGH BLOOD GLUCOSE MAY LEAD TO A MEDICAL EMERGENCY IF NOT TREATED.

What Can You Do?



CHECK BLOOD GLUCOSE

If your blood glucose levels are higher than your goal for three days and you don't know why,

CALL YOUR HEALTHCARE PROVIDER



For more information, call the Novo Nordisk Tip Line at 1-800-260-3730 or visit us online at ChangingDiabetes-us.com.

Novo Nordisk Inc. grants permission to reproduce this piece for non-profit educational purposes only on condition that the piece is maintained in its original format and that the copyright notice is displayed. Novo Nordisk Inc. reserves the right to revoke this permission at any time. Concept developed by Rhonda Rogers, RN, BSN, CDE

