Welcome to
Diabetes in the 21st Century

Beverly Dyck Thomassian, RN, MPH, BC-ADM, CDE
President, Diabetes Education Services
www.DiabetesEd.net

Diabetes in the 21st Century:
A Clinical and Educational Update

1. Describe impact of diabetes
2. Discuss prevention, management strategies
3. Discuss different types of diabetes
4. Describe insulin therapy
5. Review glucose patterns and determine how to adjust therapy to improve glucose.
6. Discuss medical nutrition therapy
7. Gain understanding of Type 2 Meds.
8. Demonstrate successful teaching strategies

Foundations of Care

▸ Education
▸ Nutrition
▸ Monitoring
▸ Physical Activity
▸ Psychosocial Care
▸ Medications
▸ Reducing Risk
▸ Getting to Best Possible Health

Diabetes in America 2016

- 29 million or > 9.3%
- 27% don't know they have it
- 37% of US adults have pre diabetes (86 mil)

Diabetes

1994 2000 2009

No Data | <4.5% | 4.5-5.9% | 6.0-7.4% | 7.5-8.9% | ≥9.0%

CDC's Division of Diabetes Translation. National Diabetes Surveillance System available at http://www.cdc.gov/diabetes/statistics

Diabetes Education SERVICES

CDC Announces

35% of Americans will have Diabetes by 2050

Boyle, Thompson, Barker, Williamson
2010, Oct 22:8(1)29
www.pophealthmetrics.com

Diabetes Education SERVICES

Type 2 in Kids

Treatment Options for type 2 Diabetes in Adolescents & Youth

- 7 fold increase 1990
- 1 in 6 overwt kids (age 12- 19) have prediabetes.
- ~2,500 to 3,700 new cases in U.S. annually.
- Highest risk: very obese, minority, female, low socioeconomic status, limited education
- In age range 12-19, less than 1% have Type 2 – NHANES
- Environmental changes to urgently needed

Diabetes Education SERVICES

Global Epidemic

- Every 10 seconds
  - 1 person dies with diabetes
  - 2 people develop diabetes
- Every year
  - 3 million deaths
  - 6 million new cases
- World Diabetes Day is November 14
- March is ADA Sound the Alert Day "find people w/ undetected diabetes"

Nov 14 – World Diabetes Day

World Diabetes Day » International Diabetes Foundation (IDF)
Celebrated on November 14 around the world to increase the global awareness of diabetes.

- 600 million people will have diabetes by 2035.
- According to IDF, 70% of type 2 cases can be prevented or delayed by adopting healthier lifestyles

Key Messages:

1. **Act to change your life today**
2. **Act to change the world tomorrow** – access to affordable healthy food is essential to reduce the global burden of diabetes.

World Diabetes Day
November 14

The right education for all

Diabetes: protect our future

The right environment for all

Diabetes: protect our future

Age-adjusted Diabetes Prevalence
20 yrs or older, by race/ethnicity— U.S. 2014

Age-adjusted* percentage of people aged 20 years or older with diagnosed diabetes, by race/ethnicity, United States, 2010–2012

Race/ethnicity	Percentage
Non-Hispanic whites	7.6
Asian Americans	9.0
Hispanics	12.8
Non-Hispanic blacks	13.2
American Indians/Alaska Natives	15.9

*Based on the 2000 U.S. standard population.
Source: 2010–2012 National Health Interview Survey and 2012 Indian Health Service's National Patient Information Reporting System.

- Among Hispanic adults, the age-adjusted rate of diagnosed diabetes was 8.5% for Central and South Americans, 9.3% for Cubans, 13.9% for Mexican Americans, and 14.8% for Puerto Ricans.
- Among Asian American adults, the age-adjusted rate of diagnosed diabetes was 4.4% for Chinese, 11.3% for Filipinos, 13.0% for Asian Indians, and 8.8% for other Asians.
- Among American Indian and Alaska Native adults, the age-adjusted rate of diagnosed diabetes varied by region from 6.0% among Alaska Natives to 24.1% among American Indians in southern Arizona.

Why Should Zip Code Determine Life Expectancy?

STOCKTON
95202
Life Expectancy
73

IRVINE
92606
Life Expectancy
88

California Endowment – look up your zip code at www.measureofamerica.org

---

Free Live Webinars and Live Seminars at DiabetesEd.net

- Free Webinars
  - Preparing to take CDE
  - New Frontiers
  - New Medications
  - BC-ADM

Role of the Pancreas
Endocrine Functions

Beta Cells - Insulin

Anabolic hormone - helps store glucose as glycogen in muscle, liver

- secreted in response to elevated glucose
- halts breakdown of glycogen in liver
- increases protein synthesis, fat storage
- powerful hypoglycemic

Beta Cells - Amylin

- secreted in 1:1 ratio with insulin
- Causes satiety
- Lowers post-prandial glucagon response
- Slows gastric emptying
- Type 1 make none
- Type 2 make less than normal amounts

Diabetes Education
SERVICES

Role of the Pancreas
Endocrine Functions

INSULIN

Normal Range

GLUCAGON

Blood Glucose Concentration

Alpha cells - Glucagon

Opposes action of insulin at the liver

- stimulated in response to low glucose levels
- stimulates liver to convert glycogen to glucose
- inhibits liver from glucose uptake
- causes hyperglycemia

Diabetes Education
SERVICES

Hormones Effect on Glucose

Hormone	Effect
Glucagon (pancreas)	⬆
Stress hormones (kidney)	⬆
Epinephrine (kidney)	⬆
Insulin (pancreas)	⬇
Amylin (pancreas)	⬇
Gut hormones - incretins (GLP-1) released by L cells of intestinal mucosa, beta cell has receptors)	⬇

Diabetes Education
SERVICES

GLP-1 Effects in Humans
Understanding the Natural Role of Incretins

GLP-1 secreted upon the ingestion of food

Promotes satiety and reduces appetite

↑ Beta-cell response

Alpha cells:
↓ Postprandial glucagon secretion

Liver:
↓ Glucagon reduces hepatic glucose output

Beta cells:
Enhances glucose-dependent insulin secretion

Stomach:
Helps regulate gastric emptying

GLP-1 degraded by DPP-4 w/in minutes

Adapted from Flint A, et al. *J Clin Invest.* 1998;101:515-520
Adapted from Larsson H, et al. *Acta Physiol Scand.* 1997;160:413-422
Adapted from Nauck MA, et al. *Diabetologia.* 1996;39:1546-1553
Adapted from Drucker DJ. *Diabetes.* 1998;47:159-169

Incretin Mimetics
Byetta, Bydureon, Trulicity, Tanzeum

▸ **Action (synthetic gut hormone)**
  ▸ Insulin release in response to meal
  ▸ Slows gastric emptying
  ▸ Causes Satiety – promotes wt loss
  ▸ Preserves Beta Cells
▸ **Details:**
  ▸ Daily and long acting version - 1x week injection
  ▸ **Efficacy:** Decreases A1c by 0.5 – 1.6%, wt by 3lbs +
▸ **Benefits/Issues** – wt loss, no hyp. Expensive, N/V
  □ Pancreatitis Warning – report signs immediately

Bariatric Surgery

▸ Consider on diabetes pts w/ BMI >35, esp with comorbidities
▸ Remission (BG normalized)
  ▸ rates range from 40 – 95%
  ▸ Better results with newer diabetes (more beta cell mass)
  ▸ Due to increase incretins (gut hormones)
▸ Still researching long term benefits, cost effectiveness and risk

Natural History of Diabetes

Normal	**Prediabetes**	**Diabetes**
FBG <100	FBG 100-125	FBG 126 +
Random <140	Random 140 - 199	Random 200 +
A1c <5.7%	A1c ~ 5.7- 6.4%	A1c 6.5% or +
	50% working pancreas	20% working pancreas

Development of type 2 diabetes happens over years or decades

Signs of Diabetes

- Polyuria
- Polydipsia
- Polyphasia
- Weight loss
- Fatigue
- Skin and other infections
- Blurry vision

- Glycosuria, H_2O losses
- Dehydration
- Fuel Depletion
- Loss of body tissue, H_2O
- Poor energy utilization
- Hyperglycemia increases incidence of infection
- Osmotic changes

Diabetes Education SERVICES

Diabetes Classifications

- Type 1
- Type 2
- Gestational
- Secondary

Diabetes Education SERVICES

Case Study

1. Pt profile: 5'8", 192 lb male
Diabetes 12 years, on insulin 3 yrs
What type of DM and how do you know?

2. 5'6", 108 lb female
On insulin 3u Regular before meals, 10u NPH at bedtime
What type of DM and how do you know?

Type 1 Diabetes – Genetics and Risk Factors

- 1- 400 to 1-1000 = Risk of type 1 in gen pop
- 1-20 to 1-50 in offspring of diabetes parents
- Combo of genes and disease susceptibility
- Risk Factors:
  - Autoimmunity tends to run in families
  - Higher rates in non breastfed infants
  - Viral triggers: congenital rubella, coxsackie virus B, cytomegalovirus, adenovirus and mumps.

Incidence of Type 1 in Youth

- **General Pop 0.3%**
- **Sibling 4%**
- **Mother 2-3%**
- **Father 6-8%**
- Rate doubling every 20 yrs
- Many trials underway to detect and prevent (Trial Net)

Type 1 – 10% of all Diabetes
Genetics and Risk Factors

➢ Auto-immune pancreatic beta cells destruction
➢ Most commonly expressed at age 10-14
➢ Insulin sensitive (require 0.5 - 1.0 units/kg/day)

 Combo of genes and environment:
   Autoimmunity tends to run in families
   Higher rates in non breastfed infants
   Viral triggers: congenital rubella, coxsackie virus
      B, cytomegalovirus, adenovirus and mumps.

Autoantibodies Assoc w/ Type 1

Panel of autoantibodies –
   ▸ GAD65 - Glutamic acid decarboxylase –
   ▸ ICA - Islet Cell Cytoplasmic Autoantibodies
   ▸ IAA - Insulin Autoantibodies

Medalist Study – Harvard
Joslin Diabetes Center

▸ After 50 years with diabetes
   ▸ Many still produced some insulin
   ▸ Many had no eye disease

Type 1 Diabetes Associated with other immune conditions

- Celiac disease (gluten intolerance)
- Thyroid disease
- Addison's Disease
- Rheumatoid arthritis
- Other

Type 1 Summary

- Autoimmune pancreatic destruction
- Need insulin replacement therapy
- Often first present in DKA
- At risk for other autoimmune diseases
- Eval coping strategies

Type 1 in Hospital

- 43 yr old admitted to evaluate angina.
- Morning blood sugar is 92.
- Based on Regular insulin sliding scale, no insulin required.
- Breakfast tray shows up and patient says, I need my insulin shot before I eat.

What do you say?

BMI Categories

17	19	22	24	26	29	33	37	40
BMI	<20	20 - 25		26 - 29		>30		
Underweight		Normal weight		Slightly Overweight		Overweight		Extremely Overweight

Diabetes Education
SERVICES

Visceral Fat and Subcutaneous Fat

- subcutaneous fat
- abdominal muscle layer
- visceral fat
- intestines

Diabetes Education
SERVICES

Natural Progression of Type 2 Diabetes

Plasma Glucose

126 mg/dL

Postprandial glucose

Fasting glucose

Relative β-Cell Function

Insulin resistance

Insulin secretion

-20   -10   0   10   20   30
Years of Diabetes

Prior to diagnosis   After diagnosis

Adapted from Bergenstal et al. 2000; International Diabetes Center.

Diabetes Education
SERVICES

Cardio Metabolic Risk -
5 Hypers -

- Hyperinsulinemia (resistance)
- Hyperglycemia
- Hyperlipidemia
- Hypertension
- Hyper"waistline"emia (35" women, 40" men)

Manifestations of Insulin Resistance

---

2. Classification and DM Diagnosis

- Pre Diabetes & Type 2- Screening Guidelines
- Start screening at age 45 or for anyone who is overweight (BMI ≥ 25, Asians BMI ≥ 23) with one or > additional **risk factor**:

  - First-degree relative w/ diabetes
  - Member of a high-risk ethnic population
  - Habitual physical inactivity
  - PreDiabetes
  - History of heart disease

Diabetes 2 - Who is at Risk?
(ADA Clinical Practice Guidelines)

Risk factors cont'd

- HTN - BP > 140/90
- HDL < 35 or triglycerides > 250
- baby >9 lb or history of Gestational Diabetes Mellitus
- Polycystic ovary syndrome (PCOS)
- Other conditions assoc w/ insulin resistance:
  - Severe obesity, acanthosis nigricans (AN)

Acanthosis Nigricans (AN)

‣ Signals high insulin levels in bloodstream
‣ Patches of darkened skin over parts of body that bend or rub against each other
   ‣ Neck, underarm, waistline, groin, knuckles, elbows, toes
   ‣ Skin tags on neck and darkened areas around eyes, nose and cheeks.
‣ No cure, lesions regress with treatment of insulin resistance

_____
_____
_____
_____
_____
_____
_____

Diabetes Detectives Needed

‣ On average – takes 6.5 years to diagnose diabetes
‣ 1/4 of all people with diabetes don't know they have it

_____
_____
_____
_____
_____
_____
_____

Ominous Octet

Decreased satiation neuro-transmission

Decreased amylin, β-cell secretion 80% loss at dx

Decreased Gut hormones

Increased lipolysis

Increase glucose production

Decreased glucose uptake

_____
_____
_____
_____
_____
_____
_____

SGLT2 Inhibitors- "Glucoretics"

Action: "Glucoretic" decreases renal reabsorption in the proximal tubule of the kidneys (reset renal threshold and increase glycosuria)

Decreases Glucose Reabsorption

SGLT2 Inhibitors		
• Decrease glucose reabsorption in kidneys • "glucoretic."	Canagliflozin (Invokana)	100–300 mg once daily
	Dapagliflozin (Farxiga)	5–10 mg once daily
	Empagliflozin (Jardiance)	10–25 mg once daily

- Benefits: Lowers A1c 0.7 – 1.5%, lowers wt 1-3 lbs, no hypo
- Issues: Can initially lower GFR, monitor kidney function and lytes. Watch for hypotension/ GU infections. Expensive

Diabetes Education SERVICES

EMPA-REG OUTCOME®: Summary

- Empagliflozin, as used in this trial, for 3 years in 1,000 patients with type 2 diabetes at high CV risk:

- Empagliflozin reduced hospitalisation for heart failure by 35%
  ¤ 14 fewer hospitalisations for heart failure (42 vs 28)

- Empagliflozin reduced CV death by 38%
  ¤ 25 lives saved (82 vs 57 deaths)
    - 22 fewer CV deaths (59 vs 37)

- Empagliflozin improved survival by reducing all-cause mortality by 32%
  ¤ 53 additional genital infections (22 vs 75)

Diabetes Education SERVICES

Comparison of
Type 1,Type 2, LADA

	Type 1	Type 2
Obesity	x	xxx
Insulin dependence	xxx	30%
Respond to oral agents	0	xxx
Ketosis	xxx	x
Antibodies present	xxx	0
Typical Age of onset	teens	adult
Insulin Resistance	0	xxx

Diabetes Education SERVICES

Comparison of Type 1 and Type 2

Feature	Type 1	Type 2
▸ Obesity	x	xxx
▸ Insulin dependence	xxx	30%
▸ Respond to oral agents	x	xxx
▸ Antibodies present	xxx	0
▸ Typical age of onset	puberty	40-65
▸ Insulin Resistance	x	xxx

Gestational DM ~ 7% of all Pregnancies

- ▸ GDM prevalence increased by
  - ▸ ~10–100% during the past 20 yrs
- ▸ Native Americans, Asians, Hispanics, African-American women at highest risk
- ▸ Immediately after pregnancy, 5% to 10% of GDM diagnosed with type 2 diabetes
- ▸ Within 5 years, 50% chance of developing DM in next 5 years.

Diabetes in pregnant mothers associated with …

- ▸ Offspring
  - ▸ Fetal Complications
  - ▸ Obesity and diabetes later in life
- ▸ Mother
  - ▸ More complicated pregnancy and delivery
  - ▸ Diabetes later in life

- ▸ Intrauterine environment is important

Postnatal Health: Maternal Behavior

- Encourage breastfeeding for one year
    - (25% of women achieving this goal)
- Screening 6-12 weeks post partum using non-pregnant OGTT criteria (50%)
- Repeat at 3 yr intervals or signs of DM
- Encourage weight control and exercise
- Make sure connected with health care
- Preconception counseling

Start Metformin therapy

- For women with PreDiabetes and History of GDM

Metformin – 1st agent of choice

- **Action:** decrease hepatic glucose (glycogen)
- Metformin (Glucophage)
    - Starting dose: 500 BID, max 2500mg daily
    - Metformin XR - extended release – less GI upset
- **Efficacy:**
    - Decrease fasting plasma glucose 60-70 mg/dl
    - Reduce A1C 1.0-2.0%
- Benefits / Issues
    - Cheap, no weight gain; some lose weight, lowers LDL, no hypo
    - ADA Stds 2016 suggests GFR may be a more appropriate measure. If GFR <45, max dose is 1000mg a day. If GFR <30, stop metformin.

Other Causes of Hyperglycemia

- Steroids
- Agent Orange
- Tube feedings / TPN
- Transplant medications
- Cystic Fibrosis

Regardless of cause, requires treatment
- Insulin always works
- Sign of pancreatic malfunction

Diabetes Education SERVICES

Diabetes is also associated with

- Fatty liver disease
- Obstructive sleep apnea
- Cancer; pancreas, liver, breast
- Alzheimer's
- Depression

Diabetes Education SERVICES

DiaBingo

B Frequent skin and yeast infections
B A BMI of _____ or greater is considered overweight
B To reduce complications, control **A**1c, **B**lood pressure, **C**holesterol
B PreDiabetes – fasting glucose level of ____ to _____
B Erectile dysfunction indicates greater risk for _____
B Diabetes – fasting glucose level____ or greater
B Type 1 diabetes is best described as an _____ disease
B People with diabetes are _____ times more likely to die of heart dx
B Elevated triglycerides, < HDL, smaller dense LDL
B Each percentage point of A1c = _____ mg/dl glucose
B At dx of type 2, about __% of the beta cell function is lost
B Diabetes – random glucose ____ or greater

Diabetes Education SERVICES

Life Study – Mrs. Jones

Mrs. Jones is 62 years old, overweight and complaining of feeling tired and urinating several times a night. She is admitted with a urinary tract Infection. Her WBC is 12.3, glucose 237. She is hypertensive with a history of gestational diabetes. No ketones in urine.

‣ What are her risk factors, signs of diabetes
‣ What type of diabetes does she have?
‣ Does she have insulin resistance?

Strategies – One Step at a Time, Focus on Survival Skills

Look for
"teaching moment"
opportunities

Diabetes Education
SERVICES

What Do You Say?
Mrs. Jones asks you

‣ What is type 2 diabetes?

‣ Will this go away?

‣ Will I get complications?

‣ Will I need to take diabetes medication for the rest of my life?

‣ How come I got diabetes?

‣ Do I have to check my blood sugars?

Diabetes Education
SERVICES

No one is Unmotivated

…. to lead and long and healthy life

▸ **These are the 3 usual Critical Barriers**
  ▸ Perceived worthlessness
  ▸ Too many personal obstacles
  ▸ Absence of support and resources

Bill Polonsky, PhD, CDE

Overcoming barriers

▸ Confront the key misbelief. Ask the question, does dm cause complications?
▸ Offer pts evidence based hope message –
▸ Frequent contact
▸ Paired glucose testing

▸ Ask pt, "Tell me 1 thing that is driving you crazy about your diabetes"
▸ Discuss medication beliefs
▸ To improve outcomes, see pts more often

Bill Polonsky, PhD, CDE

How will blood glucose testing help me?

▸ See if your treatment plan is working
▸ Make decisions regarding food and/or med adjustment when exercising
▸ Find out how that pizza affected your BG
▸ Avoid unwanted weight gain
▸ Enhanced athletic performance
▸ Find patterns
▸ Manage illness

How Often Should I Check?

▸ Be realistic!!
▸ Type 2 on orals – Medicare covers 100 strips for 3 months
▸ Based on individual - Consider:
  ▸ Types and timing of meds
  ▸ Goals
  ▸ Ability (physical and emotional)
  ▸ Finances / Insurance

Diabetes Education SERVICES

"The highest form of wisdom is kindness."
The Talmud

How many times has a person arrived disheartened?

This moment of discouragement and despair provides us an opportunity.

By modeling kindness and understanding, we can encourage them to be a kinder self-coach from this day forward.

Diabetes Education SERVICES


Give the gift of Non-Judgment

out beyond ideas
of wrongdoing
and rightdoing,
there is a field.

i'll meet you there
-rumi

Diabetes Education SERVICES

Complications - Why?

- Degree of hyperglycemia "glucose toxicity"
- Duration of hyperglycemia
- Genes
- Multiple risk factors: smoking, vascular disease, dyslipidemia, hypertension, other

Diabetes Education SERVICES

Diabetes Complications

- Heart disease leading cause of death.
- CAD death rates are about 2 -4x's as high as adults without diabetes (it's not getting better)
- Risk of stroke is 2 - 4 times higher
- 60% - 65% of people with DM have HTN.
- DM accounts for 40% of new cases of ESRD
- 60 - 70% have mild - severe forms of neuropathy
- Diabetes is the leading cause of blindness
- Accounts for 50% of lower limb amputations

Diabetes Education SERVICES

Control Matters

- **Prevention**
- **Trials**
- **Practice Recommendations**

Diabetes Education SERVICES

Financial Advisor

‣ Mid 30s, friendly, he smiles to greet you and you notice his gums are inflamed. You'd guess a BMI of 26 or so, with most of the extra weight in the waist area.

‣ If you could give him some health related suggestions, what would they be?

Can Type 2 be Prevented in Older Adults?

• Physical activity (30 mins a day)
• Dietary score (higher fiber intake, low saturated fat and *trans*-fat, lower mean glycemic index)
• Not Smoking
• Alcohol use (up to 2 drinks a day);
• BMI <25 and waist circumference

Dariush Mozaffarian, MD,
Arch Intern Med. 2009;169(8):798–807.

Overall, 9 of 10 new cases of diabetes attributable to these 5 lifestyle factors.

89% risk reduction when all at goal.

35% rel risk reduction for each additional

Can we stop pre diabetes from progressing?

3, 234 people w/ Pre-Diabetes randomized:

‣ Placebo
‣ Diet/Exercise or
‣ Metformin

over a three year period

NATIONAL Diabetes PREVENTION PROGRAM

Diabetes Prevention Program (DPP) 2001

Diabetes Prevention Program

- Standard Group - 29% developed DM
- Lifestyle Results - 14% developed DM
  - 58% (71% for 60yrs +) Risk reduction
    - 30 mins daily activity
    - 5-7% of body wt loss
- Metformin 850 BID - 22% developed DM
  - 31% risk reduction (less effective with elderly and thinner pt's)

Weight loss and Prevention

- For every 2.2 pounds of weight loss, risk of type 2 diabetes was reduced by 13%.

Use Technology to Prevent Diabetes

- Recent studies support content delivery through virtual small groups, internet social networks, cell phones and mobile devices.
  - Validated studies that these approaches can:
    - Support wt loss
    - Reduce A1c (prediabetes)
  - The CDC Diabetes Prevention Program is incorporating these tools into their program content

Diabetes Education SERVICES

Over 1200 Diabetes Apps -

AADE Diabetes Goal Tracker

Diabetes Education SERVICES

Goals of Care

Diabetes Education SERVICES

ABCs of Diabetes –

- A1c less than 7% (avg 3 month BG)
  - Pre-meal BG 80-130
  - Post meal BG <180
- Blood Pressure < 140/90
- Cholesterol
  - DM and 40 yrs, start statin
  - HDL >40
  - Triglyceride < 150
- E xercise, Education
- H ealthy Eating

Glucose and BP Control Matter

- 1% decrease in A_1c reduces microvascular complications by 35%
- 1% decrease in A_1c reduces diabetes related deaths by 25%

- B/P control (144/82) reduced risk of:
  - Heart failure (56%)
  - Stroke (44%)
  - Death from diabetes (32%)

Lancet 352: 837-865, 1998

## 6. Glycemic Targets

- **Adult non pregnant A1c goals**
  - **A1c < 7%** - a reasonable goal for adults.
  - **A1c < 6.5%** - may be appropriate for those without significant risk of hypoglycemia or other adverse effects of treatment.
  - **A1c < 8%** - may be appropriate for patients with history of hypoglycemia, limited life expectancy, or those with longstanding diabetes and vascular complications.

Goals for Older Adults (Std 10)

Table 10.1—Framework for considering treatment goals for glycemia, blood pressure, and dyslipidemia in older adults with diabetes

Patient characteristics/ health status	Rationale	Reasonable A1C goal‡	Fasting or preprandial glucose	Bedtime glucose	Blood pressure	Lipids
Healthy (few coexisting chronic illnesses, intact cognitive and functional status)	Longer remaining life expectancy	<7.5% (58 mmol/mol)	90–130 mg/dL (5.0–7.2 mmol/L)	90–150 mg/dL (5.0–8.3 mmol/L)	<140/90 mmHg	Statin unless contraindicated or not tolerated
Complex/intermediate (multiple coexisting chronic illnesses* or 2+ instrumental ADL impairments or mild-to-moderate cognitive impairment)	Intermediate remaining life expectancy, high treatment burden, hypoglycemia vulnerability, fall risk	<8.0% (64 mmol/mol)	90–150 mg/dL (5.0–8.3 mmol/L)	100–180 mg/dL (5.6–10.0 mmol/L)	<140/90 mmHg	Statin unless contraindicated or not tolerated
Very complex/poor health (LTC or end-stage chronic illnesses** or moderate-to-severe cognitive impairment or 2+ ADL dependencies)	Limited remaining life expectancy makes benefit uncertain	<8.5%† (69 mmol/mol)	100–180 mg/dL (5.6–10.0 mmol/L)	110–200 mg/dL (6.1–11.1 mmol/L)	<150/90 mmHg	Consider likelihood of benefit with statin (secondary prevention more so than primary)

Diabetes Education Services

A1c and Estimated Avg Glucose (eAG) 2008

A1c (%)	eAG
5	97
6	126
7	154
8	183
9	212
10	240
11	269
12	298

Order teaching tool kit free at diabetes.org

eAG = 28.7 x A1c-46.7 ~ 29 pts per 1%

Translating the A1c Assay Into Estimated Average Glucose Values – ADAG Study

Diabetes Care: 31, #8, August 2008

Diabetes Education Services

"Legacy Effect"

‣ For participants of DCCT and UKPDS
  ‣ long lasting benefit of early intensive BG control prevents
    ‣ microvascular complications
    ‣ Macrovascular complications (15-55% decrease)
  ‣ Even though their BG levels increased over time
  ‣ Message – Catch early and Treat aggressively

Diabetes Education Services

Updated Recommendations

TABLE 7. Recommendations for Statin and Combination Treatment in People With Diabetes

Age	Risk factors	Recommended statin intensity*
<40 years	None	None
	ASCVD risk factor(s)**	Moderate or high
	ASCVD	High
40–75 years	None	Moderate
	ASCVD risk factors	High
	ASCVD	High
	ACS and LDL cholesterol >50 mg/dL (1.3 mmol/L) in patients who cannot tolerate high-dose statins	Moderate plus ezetimibe
>75 years	None	Moderate
	ASCVD risk factors	Moderate or high
	ASCVD	High
	ACS and LDL cholesterol >50 mg/dL (1.3 mmol/L) in patients who cannot tolerate high-dose statins	Moderate plus ezetimibe

*In addition to lifestyle therapy.
**ASCVD risk factors include LDL cholesterol ≥100 mg/dL (2.6 mmol/L), high blood pressure, smoking, overweight and obesity, and family history of premature ASCVD. ACS, acute coronary syndrome.

▸ Statin – lowers cholesterol production in liver
▸ Ezetimibe (Zetia) – blocks absorption of cholesterol in intestine

---

Vaccinations- Immunizations

▸ Flu vaccine
  ▸ every year starting 6 months
▸ Pneumococcal starting at 2 years.
  ▸ One time Revaccination for those over 64 and had first vaccine >5 years prior
▸ Hepatitis B Vaccine (ADA Stds 2013, pg s28)
  ▸ For diabetes pts age 19 – 59 (not previously vaccinated)
  ▸ Double risk of Hep B due to lancing devices/ glucose meter exposure

---

Education

▸ People with diabetes and pre diabetes should receive DSME
  ▸ Monitor for effective self-management and quality of life
  ▸ Address psychosocial issues and emotional well being
  ▸ Results in cost savings and improved outcomes, should be reimbursed by third party payers.

Exercise Recommendations

- **Activity update –Don't sit more than 90 minutes**
- Evidence supports that everyone, including with diabetes should be encouraged to reduce sedentary time, by not sitting for more than 90 minutes at a time.
- It is recommended that people with pre diabetes and diabetes engage in 150 minutes of activity a week and at **least 2 weekly sessions of resistance exercise.**

Good Exercise Info / Quotes

- 20 % of people walk 30 mins a day
- Exercise decrease A1c 0.7%
- No change in body wt, but 48% loss in visceral fat
  - ADA PostGrad 2010

- "If you don't have time for exercise, you better make time for disease."

"I don't have time to exercise, I MAKE time."

Mike Huckabee

Vaccinations- Immunizations

- Flu vaccine
  - every year starting 6 months
- Pneumococcal starting at 2 years.
  - One time Revaccination for those over 64 and had first vaccine >5 years prior
- Hepatitis B Vaccine
  - For diabetes pts age 19 – 59 (not previously vaccinated)
  - Double risk of Hep B due to lancing devices/ glucose meter exposure

Pneumonia Vaccination Update

- Pneumonia polysaccharide PPSV23 vaccine to all patients starting at age 2
- **Adults ≥ 65 years of age**, if not previously vaccinated, should receive pneumococcal conjugate vaccine 13 (PCV13), followed by PPSV23 6-12 months after initial vaccination.
- **Adults ≥ 65 years of age**, if previously vaccinated with PPSV23 should receive a follow-up ≥ 12 months with PCV13.

DiaBingo- G

G ADA goal for A1c is less than _____%

G People with DM need to see their provider at least every month

G Blood pressure goal is less than

G People with DM should see eye doctor (ophthalmologist) at least

G The goal for triglyceride level is less than

G Goal for my HDL cholesterol is more than

G The goal for blood sugars 1-2 hours after a meal is less than:

G People with DM should get this shot every year

G People with DM need to get urine tested yearly for _____

G Periodontal disease indicates increased risk for heart disease

G The goal for blood sugar levels before meals is:

G The activity goal is to do ____ minutes on most days

Diabetes Care Guidelines- ADA

Test / Exam	Frequency
A1c	At least twice a year
B/P	Each diabetes visit
Cholesterol (LDL, HDL, Tri)	Yearly (less if normal)
Weight	each diabetes visit
Microalbumin/GFR/Creat	Yearly
Eye exam	Yearly
Dental Care	At least twice a year
Comprehensive Foot Exam	Yearly (more if high risk)
Physical Activity Plan	As needed to meet goals
Preconception counseling	As needed

Mr. Jones - What are Your Recommendations?

Patient Profile

64 yr old with type 2 for 11 yrs. Hx of CVD.

Labs:

- A1c 9.3%
- HDL 37 mg/dl
- Triglyceride 260mg/dl
- Proteinuria - neg
- B/P 152/94

Self-Care Skills

- Walks dog around block 3 x's a week
- Bowls every Friday
- 3 beers daily
- *What meds?*
- *What referrals?*
- *My foot hurts*

Glucose Management and Hospitalized Patients

- In hospitalized patients with critical illness, hyperglycemia is a signal that warrants our attention.

Hospitals and Hyperglycemia – What's the Big Deal?

- Hyperglycemia is associated with increased morbidity and mortality in hospital settings.
  - Acute Myocardial Infarction
  - Stroke
  - Cardiac Surgery
  - Infection
  - Longer lengths of stay

WHAT SHOULD WE AIM FOR?

Critically Ill pts
- BG > 180- Start insulin
- BG goal 140-180

Non Critically Ill patients BG Goals
- Premeal <140
- Post meal <180

•Insulin therapy preferred treatment

Consensus: Inpt Hyperglycemia, Endocr Pract. 2009;15 (No.4)

Management of Hyperglycemia and Diabetes

▸ Stop oral agents (ie) metformin & sulfonylurea on admission
▸ "The sole use of Sliding Scale insulin is discouraged" – ADA 2014
▸ For discharge, oral meds can be resumed

Start Basal/bolus therapy
▸ NPH and Regular insulin
▸ Long-acting and rapid-acting insulin
▸ Premixed insulin

Now What?

▸ Nurse had an emergency and pt already ate lunch?

▸ Nurse administered insulin and pt only ate a few bites of turkey and drank non sugar tea?

▸ You just gave 3 units of Regular and patient needs to go to OR NOW!

Foot Care

Lift the sheets and look at the Feets!

Diabetes Education

Foot Wounds

| Blisters | Ulcers | Bone infection |
| Calluses | | |

Diabetes Education

No Bathroom Surgery

Diabetes Education

5.07 monofilament = 10gms linear pressure

Instrument in use

10-guage nylon monofilament

Free Monofilaments
http://www.hrsa.gov/leap/

Mr. Jones - What are Your Recommendations?

Patient Profile

64 yr old with type 2 for 11 yrs. Hx of CVD.

Current Status:

‣ A1c 9.3%
‣ On Metformin 500mg BID
‣ Partial foot amputation
‣ Lives alone
‣ What resources, teaching?

Three Most Important Foot Care Tips

‣ Inspect and apply lotion to your feet every night before you go to bed.

‣ Do NOT go barefoot, even in your house. Always wear shoes!

‣ Every time you see your doctor, take off your shoes and show your feet.

Bottom Line

- 30-40% of hospitalized patients have diabetes
  - 10% aren't officially diagnosed
- Cardiovascular disease is the leading cause of hospitalization for people with diabetes
- Look for patients with hyperglycemia and cardiometabolic risk factors: smokers, HTN, central obesity, abnormal lipids, Acanthosis.
- Provide education and promote self-advocacy

"Getting diabetes saved my life."
~ Sherri Sheperd

Sherri Shepard decided to embrace diabetes and use it as a motivator to improve her health.

Insulin Therapy
From Ants to Analogs:

Insulin – the Ultimate Hormone Replacement Therapy

Objectives:
- Discuss the actions of different insulins
- Describe using pattern management as an insulin adjustment tool.

Human Proinsulin Molecule

Diabetes Education Services

Psychological Insulin Resistance (PIR)

- 50% of providers in study threatened pts "with the needle".
- Less than 50% of providers realized insulins' positive effect on type 2 dm
- Most pts don't believe that insulin would "better help them manage their diabetes".
- Solutions: Find the root of PIR and address

Diabetes Attitudes, Wishes, Needs Study - Rubin

Diabetes Education Services

Needle Size often a Barrier
Size *Does* Matter

(Actual Size)
BD Original 12.7mm (½") BD Short 8 mm (5/16") BD Mini 5 mm (3/16") BD Nano 4 mm (5/32")
BD Nano 4mm and BD Mini 5mm only available in pen needles

- Use more short needles – 4 mm
- Effective for pts with BMI of 24- 49
- Keeps it subq
- If pt thin, inject at angle
- To avoid leakage, count to 10 before withdrawing needle
- ½ the patients who could benefit from insulin are not using it due to needle phobias

Diabetes Education Services

Physiologic Insulin Secretion: 24-Hour Profile

Insulin (µU/mL)

50
25
0

Bolus Insulin

Basal Insulin

⇧ Breakfast ⇧ Lunch ⇧ Dinner

Glucose (mg/dL)

150
100
50
0

Mealtime Glucose

Basal Glucose

7 8 9 10 11 12 1 2 3 4 5 6 7 8 9
A.M.   P.M.

Time of Day

Diabetes Education Services

Insulin Action Teams

▸ Bolus: lowers after meal glucose levels
  ▸ Rapid Acting
    ▸ Aspart, Lispro, Glulisine, Afrezza
  ▸ Short Acting
    ▸ Regular
▸ Basal: controls glucose between meals, hs
  ▸ Intermediate
    ▸ NPH
  ▸ Long Acting
    ▸ Detemir (Levemir)
    ▸ Glargine (Lantus, Basaglar)
    ▸ Degludec (Tresiba)

Diabetes Education Services

Case Study

▸ 70 yr old, weighs 100kg
▸ History of CABG, tobacco
▸ A1c – 11.3%, BG 400-500 for past weeks
▸ Insulin – 100+ units Lantus at hs (solostar)
▸ Oral Meds: Metformin, Invokana
▸ What is a better insulin dosing strategy?
▸ Pt can't afford insulin pen – what other option
▸ Diabetes Meds on a Budget - 2014 - provides practical and affordable strategies to manage hyperglycemia

Diabetes Education Services

Cost Per Vial in Northern CA

Per vial cost	Walmart	Walgreens	Costco
Regular Insulin	$25*	$92	$99
NPH	$25*	$92	$99
70/30	$25*	$92	$101
Humalog	$200	$220	$178
Novolog	$197	$217	$178
Apidra	$180	$246	$178
Levemir	$300	$300	$300
Lantus	$226	$221	$206

Despite the valiant efforts of the research group,
the insulin suppository still had one major problem.

®1999 Kings Publishing

Bolus Insulins
(½ of total daily dose ÷ meals)

Name	Onset	Peak Action
‣ Lispro (Humalog)	15-30 min	1-1.5 hrs
‣ Aspart (NovoLog)		
‣ Glulisine (Apidra)		
‣ Afrezza (Inhaled)		
‣ Regular	30 mins	2-4 hrs

Bolus Insulin Summary

‣ Regular, Novolog, Humalog, Apidra,
‣ Starts working fast (15-30 mins)
‣ Gets out fast (3-6 hours)
‣ Post meal BG reflects effectiveness
‣ Should comprise about ½ total daily dose
‣ Covers food or hyperglycemia.
‣ 1 unit
  ‣ Covers ≈ 10 -15 gms of carb
  ‣ Lowers BG ≈ 30 – 50 points

Bolus Insulin Timing

‣ How is the effectiveness of bolus insulin determined?
  ‣ 2 hour post meal (if you can get it)
  ‣ Before next meal blood glucose

‣ Glucose goals (ADA) – may be modified by provider/pt
  ‣ 1-2 hours post meal <180
  ‣ Before next meal – 70 - 130

Pattern Management –AKA

How to think like a pancreas

Pattern Management

- Safety 1st!! - Evaluate 3 day patterns
- **Hypo:** eval 1st and fix:
  - If possible, decrease medication dose
  - Timing of meals, exercise, medications
- **Hyperglycemia:** evaluate 2nd
  - Identify patterns
  - Before increase insulin, make sure not missing something (carbs, exercise, omission)

Type 2 – BMI 32. New diagnosis, No meds. What Patterns? Recommendations? Meds?

	Break	Lunch	Dinner	HS
Day 1	164			181
Day 2		124	106	195
Day 3	149		102	242
Day 4	151	81		211

Bolus – Insulin Sliding Scale
Starts at 150, 2 units for every 50 mg/dl >150

	Break	Lunch	Dinner	HS
Day 1	94 no insulin	212 4 uR	148 no insulin	254 6 uR
Day 2	243 4uR	254 6 uR	201 4uR	199 no insulin
Day 3	189 2uR	243 4uR	162 2uR	244 4uR
Day 4	66 No insulin	287 6uR	144 none	272 6uR

Basal Insulins
(½ of total daily dose)

Intermediate Acting	Peak Action	Duration
▸ NPH	4-12 hrs	12-24

Long Acting	Peak Action	Duration
▸ Detemir (Levemir)	No Peak	20 hrs
▸ Glargine (Lantus)		24 hrs
▸ Glargine (Basaglar)		24 hrs
▸ Degludec (Tresiba)		42 hrs

Fasting BG reflects efficacy of basal

Degludec and Ryzodeg

▸ Degludec (Tresiba)
  ▸ An ultra long acting insulin - lasts up to 42 hours
  ▸ Takes 3-4 days to reach steady state
  ▸ Available in u-100 and u-200 pens
  ▸ Seems to cause less hypo
  ▸ Adjust dose every 3-4 days
  ▸ Wait at least 8 hours between doses
  ▸ Good at room temp for 8 wks

▸ Ryzodeg 70/30
  ▸ mixture of insulin degludec and aspart

Insulin Degludec | A new-Generation-ultra-long acting basal insulim

---

Basal Insulin Summary

▸ NPH, Levemir, Lantus, Degludec
▸ Covers in between meals, through night
▸ Starts working slow (4 hours)
▸ Stays in long (12-42hours)
  ▸ NPH 12 hrs
  ▸ Levemir, Lantus 20-24 hrs
  ▸ Degludec – 42 hours
▸ Fasting blood glucose reflects effectiveness

Basal + Metformin
Type 2, 80kg – A1c 8.7%

	Break	Lunch	Dinner	HS
Mo 1	170s			298 10uNPH
Mo 2	160s			233 20uNPH
Mo 4	140s	283	265	206 40uNPH

Diabetes Education SERVICES

Figure 3. Approach to starting & adjusting insulin in T2DM

Basal Insulin (usually with metformin +/- other non-insulin agent)
- **Start:** 10U/day or 0.1-0.2 U/kg/day
- **Adjust:** 10-15% or 2-4 U once-twice weekly to reach FBG target.
- **For hypo:** Determine & address cause; ↓dose by 4 units or 10-20%.

If not controlled after FBG target is reached (or if dose > 0.5 U/kg/day) treat PPG excursions with meal-time insulin. (Consider initial GLP-1-RA trial.)

Add 1 rapid insulin* injections before largest meal
- **Start:** 4U, 0.1 U/kg or 10% basal dose. If A1c<8%, consider ↓basal by same amount.
- **Adjust:** ↑dose by 1-2 U or 10-15% once-twice weekly until SMBG target reached.
- **For hypo:** Determine and address cause; ↓corresponding dose by 2-4 U or 10-20%.

Change to premixed insulin* twice daily
- **Start:** Divide current basal dose into 2/3 AM, 1/3 PM or 1/2 AM, 1/2 PM.
- **Adjust:** ↑dose by 1-2 U or 10-15% once-twice weekly until SMBG target reached.
- **For hypo:** Determine and address cause; ↓corresponding dose by 2-4 U or 10-20%.

Add ≥2 rapid insulin* injections before meals ('basal-bolus')
- **Start:** 4U, 0.1 U/kg or 10% basal dose/meal. If A1c<8%, consider ↓basal by same amount.
- **Adjust:** ↑dose by 1-2 U or 10-15% once-twice weekly to achieve SMBG target.
- **For hypo:** Determine and address cause; ↓corresponding dose by 2-4 U or 10-20%.

When is it Too much basal insulin?

Basal Insulin (usually with metformin +/- other non-insulin agent)
- **Start:** 10U/day or 0.1-0.2 U/kg/day
- **Adjust:** 10-15% or 2-4 U once-twice weekly to reach FBG target.
- **For hypo:** Determine & address cause; ↓dose by 4 units or 10-20%.

If not controlled after FBG target is reached (or if dose > 0.5 U/kg/day) treat PPG excursions with meal-time insulin. (Consider initial GLP-1-RA trial.)

ulin* injections gest meal

premixed i

Diabetes Education SERVICES

Next Steps

- At max basal dose
  - 80 x 0.5 = 40 units
- Add Sulfonylurea?
- Consider Adding a SGLT-2 Inhibitor
- Consider a GLP-1 Agonist
- Start bolus insulin at largest meal
- Or switch to 70/30 Insulin

Combo Sub-Q Insulin

Insulin Type	Onset	Peak
Humalog Mix 75/25: 75% NPL, 25% lispro 50/50: 50% NPL, 50% lispro	0.25 - 0.5 hr	0.5-6.5 hrs
NovoLog Mix 70/30: 70% NPA, 30% aspart	0.25 - 0.5 hr	1 – 4 hrs
NPH + Reg Combo 70/30: 70%N /30%R 50/50: 50%N /50%R	0.5 – 1.0 hr	2 - 16 hrs

Considerations:
▫ Pre-mixed, difficult to fine tune therapy

Next Steps – Switch from 40 units basal to 70/30 Insulin

- Switch to 70/30 Insulin
- Take current dose and give 2/3 in am and 1/3 in pm.
  - 2/3 of basal in am
    - 40 units x 0.6 = 24 units 70/30
  - 1/3 of basal in *pm
    - 40 units x 0.4 = 16 units 70/30
  - *pm = before dinner

24u 70/30 am, 16 u 70/30 pm
Patterns? Changes needed?

	Break	Lunch	Dinner	HS
Day 1	102	63	92	181
Day 2	112	67	106	195
Day 3	98	56	112	201
Day 4	99	71	132	211

Diabetes Education
SERVICES

_____

_____

_____

_____

_____

_____

_____

Type 2 – Glyburide 20mg AM, 10u Lantus pm

	Break	Lunch	Dinner	HS
Day 1	164	94	66	162
Day 2	169		59	195
Day 3		84	81	242
Day 4	159		43	211

Diabetes Education
SERVICES

_____

_____

_____

_____

_____

_____

_____

What Medications Cause Hypoglycemia?

▸Insulin
▸Sulfonylureas
▸Meglitinides
▸ Or any combo medication
  that includes these

Diabetes Education
SERVICES

_____

_____

_____

_____

_____

_____

_____

Sulfonylureas - Squirts

- Action: Increase endogenous insulin secretion throughout day
- Efficacy:
  - Decrease FPG 60-70 mg/dl
  - Reduce A1C by 1.0-2.0%
- Side Effects:
  - Weight gain, hypoglycemia
- Benefits:
  - Cheap, effective

Hypoglycemia = "Limiting Factor"

- Defined as glucose of 70mg/dl or below
- 50% of episodes occur during the night
- Higher mortality rate with severe hypoglycemia secondary to sulfonylureas
  - Especially (glyburide) Micronase°, Diabeta°
- Blood glucose levels don't describe severity, response is individual

---

Hypoglycemic Symptoms

- Autonomic
  - Anxiety
  - Palpitations
  - Sweating
  - Tingling
  - Trembling
  - Hypoglycemic Unawareness
- Neuroglycopenia
  - Irritability
  - Drowsiness
  - Dizziness
  - Blurred Vision
  - Difficulty with speech
  - Confusion
  - Feeling faint

Treatment of Hypoglycemia

▸ If blood glucose **70**mg/dl or below:
  - 10-15 gms of carb to raise BG 30 - 45mg/dl

- Retest in 15 minutes, if still low, treat again, even without symptoms
- Follow with usual meal or snack
- If BG less than 40, allow recovery time

15 - 20 Gms Carb Sources

- 3 - 4 Glucose Tablets
- 8 - 10 Lifesavers candy
- 8 - 10 Hard candies
- 2 Tablespoons Raisins
- 4 - 6 oz's Nondiet soda
- 4 - 6 oz's Fruit Juice
- 8 oz Milk (non fat)

Basal Bolus – What Adjustments?
Pt weighs 80kg

	Break	Lunch	Dinner	HS
Day 1	69 7R	79 5R	245 8R	190 22u NPH
Day 2	81 7R	87 5R	170 8R	133 22u NPH
Day 3	73 7R	94 5R	194 8R	110 22u NPH
Day 4	62 7R	83 5R	211 8R	127 22u NPH

Intensive Diabetes Therapy
Insulin Dosing Strategy

50/50 Rule
▸ 0.5-1.0 units/kg day

▸ Basal = 50% of total
  ● Glargine QD
  ● NPH or Detemir BID

✱ Bolus = 50% of total
  ● usually divided into 3 meals

Example
▸ Wt 50kg x 0.5 = 25 units of insulin/day

▸ Basal dose: 13 units
  ● Glargine 13 units QD
  ● NPH/Detemir 6u BID

▸ Bolus dose: 12 units
  ▸ 4 units NovoLog, Apidra Humalog, Regular each meal

Intensive Diabetes Therapy
Insulin Dosing Strategy

50/50 Rule
▸ 0.5-1.0 units/kg day

▸ Basal = 50% of total
  ● Glargine QD
  ● NPH or Detemir BID
✱ Bolus = 50% of total
  ● usually divided into 3 meals

Example – You Try
▸ Wt 60 kg x 0.5 = ____ units of insulin/day

▸ Basal dose: ____ units
  ● Glargine ____ QD
  ● NPH/Detemir ___ BID

▸ Bolus dose: ____ units
  ____ units NovoLog, Apidra Humalog, Reg each meal

Basal Bolus – <u>Using 50/50 Rule</u> - Pt weighs 80kg

	Break	Lunch	Dinner	HS
Day 1	84 6H	89 7H	145 7H	190 20 u Det
Day 2	81 6H	97 7H	107 7H	133 20u Det
Day 3	79 6H	104 7H	124 7H	110 20u Det
Day 4	69 6H	103 7H	208 7H	193 20u Det

Diabetes Education SERVICES

Insulin Teaching Keys

- Bolus insulin with meals
- Basal 1-2xs daily
- Abdomen preferred injection site
- Stay 1" away from previous site
- Don't re-use ultra fine syringes
- Keep unopened insulin in refrigerator

- Toss opened insulin vial after 28 days
- Proper disposal
- Review patients ability to withdraw and inject.
- Side effects include hypoglycemia/wt gain
- Insulin pens –
  - Prime needle to assure accurate insulin dose given
  - Hold needle in for 5 seconds after injection
  - Roll 70/30 pens

Diabetes Education SERVICES

Sharps Disposal: Product and Info

- Look in the Government section white pages for a household hazardous waste listing for your city or county.
- Call 1-800-CLEANUP (1-800-253-2687)
- Search for collection centers on the California Integrated Waste Management Board (CIWMB) Web site: http://www.ciwmb.ca.gov/HHW/HealthCare/Collection/
-

Diabetes Education SERVICES

Diabetes Vacations

"The greatest glory in living lies not in never falling, but in rising every time we fall."
~Nelson Mandela

DiaBingo - I

I Injected hormone that is an analog of amylin

I Glargine, Detemir, NPH are types of

I Breakdown of glycogen into glucose

I Anabolic hormone

I Insulin is released when glucose levels are low

I Once opened, insulin vials are good for one _____

I Elevated post-prandial glucose indicate need for pre-meal

I Epinephrine increases insulin resistance

I Creation of glucose from amino acids and lactate

I Decreasing renal function for people on insulin can cause

I Bolus insulins

I A hormone that increases blood glucose levels

U.S. Weight - 68% overweight or obese

▸ 34% BMI 25-29
▸ 34% BMI 30 +
▸ 1/3 of all overwt people don't get diabetes
▸ We burn 100 cals less a day at work
▸ Overall, food costs ~ 10-15% of income
▸ Calorie Intake is on the rise

Average Daily Per Capita Calories Adjusted for Waste. Source: Economic Research Service of the United States Department - sility (Per Capita) Data System.

Average American Consumes 25 teaspoons of sugar a day (400 cals)

‣ Warning label on sodas proposed

‣ One soda has 12 teaspoons sugar

‣ On avg, 1 person consumes 40 gallons of soda each year

‣ ADA guidelines "limit sodas and beverages with sugar, High Fructose Corn Syrup, (HFCS)

Bacterial Cells Outnumber Human Cells 10 to 1

• 10 trillion human cells
• Host 100 trillion bacterial and fungal cells

The Human Microbiome Project

Poll Question 1

‣ How much does your gut bacteria weigh?

A. 24 ounces

B. 3 pounds

C. Less than 1 pound

D. 1.5 pounds

E . Not sure

3 lbs of Microbes in our Gut

- This community of bacteria can be thought of as an extra 'organ' "microbiome".
- We have evolved together with our microbiome over millions of years.
- Ratios of these communities has changed over the past 30 years
- Mirrors global spikes in obesity, diabetes, allergic and inflammatory diseases
- What are we doing to change these bacteria?

Diabetes Education

Gut Microbiome

- Part of endocrine axis
- Stabilized by 3 years of age
- Influenced by:
  - Birth method
  - Breast fed
  - Early Antibiotic use
  - Environment
  - Travel
- Help us
  - utilize energy
  - fight off invaders

Microbes maketh man

Human Intestine Friends

- The majority belong 2 major phyla:
- Firmicutes
  - includes *Clostridium*, *Enterococcus*, *Lactobacillus* and *Ruminococcus*
- Bacteroidetes
  - includes *Bacteroides* and *Prevotella*

in proportions determined in part by birth, breastfeeding, diet

Weight and Gut Bacteria
New and Early Research

- Leaner people appear to have more bacterial diversity and a higher proportion of bacteroidetes
  - Gut bacteria less efficient at converting food to calories
- Obese people appear to have higher levels of firmicutes
  - Gut bacteria very efficient at calorie extraction
- Bacteria tend to run in families

Getting to Better Gut Bacterial Health

Eat more PREbiotics
- Foods with indigestible fibers that nourish the good bacteria:
  - High fiber foods like, whole grains, fruits, veggies, nuts
  - High in prebiotic fibers include: Jerusalem artichokes, onions, kale, Brussels sprouts, bananas, dandelion greens & more

PRObiotics
- These foods contain healthy bacteria like *Bifidobacterium* and *lactobacillus.*
  - Yogurt, Kefir – look for "live or active cultures"
 - Fermented foods like: Sauerkraut, Kimchi, Miso soup, kombucha

---

Follow Your Gut – Dr. Rob Knight

THE GARDEN OF YOU

Check out Dr. Knight's:
- TED Talk
- Website – AmericanFoodProject.org
- Articles in Nature and all over

Medical Nutrition Therapy – ADA

▸ Focus on the Individual
▸ Maintain pleasure of eating
▸ Provide positive messages about food
▸ Limit food choices only when backed by science
▸ Provide practical tools
▸ Refer to a RD and Diabetes Education – Lowers A1c by 1-2%

Approach Depends on Patient

• New Type 2
  • Portion Control
  • Plate Method
  • Record Keeping
  • Education
• On Insulin?
  • Carb counting
  • Post prandial checks

What are next steps?

▸ 72 yr old, thin, lives alone, A1c 7.3%. History of MI, stroke. DM for 12 yrs, "diet controlled". Limited income. Creat 1.4.

DPP-4 Inhibitors – "Incretin Enhancers"
Januvia (sitagliptin) – Tradjenta (linagliptin)
Onglyza (saxagliptin) Nesina (alogliptin)

▸ **Action:**
  ▸ Increase insulin release w/ meals
  ▸ Suppress glucagon
▸ **Dosing**: Januvia – 100mg a day
          Onglyza – up to 5mg a day
          Tradjenta – 5mg a day
          Nesina – up to 25 mg a day
▸ **Efficacy:** Decreases A1c by 0.6 -0.8%
▸ **Benefits/ Issues**: weight neutral, no hypo, few
  side effects. Expensive

Losing 2-8kg Early in diagnosis Type 2 Helpful
ADA 2014

▸ Weight Loss –
  ▸ *The optimal macronutrient intake to lose weight not known*
  ▸ *The literature does not support one particular nutrition therapy to reduce weight, but rather a spectrum of eating patterns that result in reduced energy intake.*

  ▸ To lose one pound – avoid 3,500 cals
    ▸ Decrease intake 250-500 cals daily + exercise

Successful weight loss strategies include

▸ Weekly self-weighing
▸ Eat breakfast
▸ Reduce fast food intake.
▸ Decrease portion size
▸ Increase physical activity
▸ Use meal replacements
▸ Eat healthy foods

Diabetes Prevention Program
Focus on fat = wt loss success

To help you lose weight and improve your health, stay as close as possible to your fat and calorie goals.
Find your starting weight below. Your fat and calorie goals are in the same row. Circle your fat and calorie goals.

Weight (lb)	Fat Goal (grams)	Calorie Goal
120-174	33	1,200
175-219	42	1,500
220-249	50	1,800
>250	55	2,000

http://www.cdc.gov/diabetes/prevention/recognition/curriculum.htm

Diabetes Education SERVICES

How nutrients affect blood sugar

Diabetes Education SERVICES

Carbohydrate Needs for Most Adults

	Grams	Servings
Each Meal	45-60 gm	3 - 4
Snacks	15-30 gm	1- 2

Carbs affect Post Meal Blood Glucose

Diabetes Education SERVICES

Choose Healthy Carbs

o Carbs have fiber, vitamins, minerals and phytonutrients

o 25 gms of fiber a day

o Power Carbs include:

  o Beans

  o Veggies

  o Fruits

  o Whole grain foods

Another plate example

---

10 Superfoods

‣ Beans
‣ Dark Green Leafy Vegs
‣ Citrus Fruit
‣ Sweet Potatoes
‣ Berries

‣ Tomatoes
‣ Fish High in Omega-3 Fatty Acids
‣ Whole Grains
‣ Nuts
‣ Fat-Free Milk and Yogurt

Carb Counting - Starch

Each Food has:
80 Calories
15 grams carb

1/2 cup cooked beans

1 small ear of corn or 1/2 cup corn

1/3 cup cooked pasta

3/4 cup cold cereal

1 slice bread

1/3 cup cooked rice

1 small potato

1/2 English muffin

1 small tortilla

5-6 small crackers

Diabetes Education SERVICES

Carb counting- fruit

Each Food has:
60 Calories
15 grams carb

1 small fresh fruit

½ cup fruit juice

½ banana

½ cup unsweetened apple sauce

1 slice bread

17 small grapes

1 cup melon

¼ cup dried fruit

2 tbsp raisins

1 1/4 cup strawberries

Diabetes Education SERVICES

Carb Counting - Milk

Each Food has:
90-150 calories
12-15 grams carb

8 oz buttermilk

1 packet diet hot cocoa

1 slice bread

6 oz plain yogurt

8 oz milk

8 oz soy milk

6 oz light fruit yogurt

Diabetes Education SERVICES

Carb Counting - Sweets

Each Food has:
Calories vary
15 grams carb

2 inch square cake or brownie, unfrosted

½ cup diet pudding

½ cup regular jello

2 tbsp light syrup

1 slice bread

2 small cookies

1 tbsp syrup, jam, jelly, table sugar, honey

½ cup ice cream or frozen yogurt

½ cup sherbet

¼ cup sorbet

Diabetes Education SERVICES

Move toward the Tomato

Diabetes Education SERVICES

USDA Plate Method
www.myplate.gov

ChooseMyPlate.gov

Balancing Calories
- Enjoy your food, but eat less.
- Avoid oversized portions.

Foods to Increase
- Make half your plate fruits and vegetables.
- Make at least half your grains whole grains.
- Switch to fat-free or low-fat (1%) milk.

Foods to Reduce
- Compare sodium in foods like soup, bread, and frozen meals — and choose the foods with lower numbers.
- Drink water instead of sugary drinks.

Diabetes Education SERVICES

Mindful Eating

Mindful Eating
www.eatingmindfully.com

Aware
Tasting vs. mindless munching

Savor
Notice the texture, aroma, and flavor. (Is it crunchy, sweet, salty smooth, spicy?)

Observe
Notice your body, rumbling stomach, low energy, stressed out, satisfied, full, empty.

Nonjudgment
Speak mindfully and compassionately. Notice when "shoulds," rigid rules or guilt pop into your mind.

In-the-Moment
Be fully present. Turn off the T.V. Sit down. When you eat, just eat.

Susan Albers PsyD 2012# Eat, Drink & Be Mindful

Diabetes Education SERVICES

Using Alcohol Safely

- Women- 1 or fewer alcoholic drinks a day
- Men 2 or fewer alcoholic drinks a day
  - 1 alcoholic drink equals
    - 12 oz beer, 5 oz glass of wine, or 1.5 oz distilled spirits (vodka, gin etc)
- If drink, limit amount and drink w/ food.
- Ask HCP if safe for you to drink. Tell them your usual quantity and frequency.
- Can cause hypo and worsen neuropathy

Diabetes Education SERVICES

Ms. Gonzales' Daily Meal plan

Break	Lunch	Dinner	Night
5 corn tortillas, 1/2 c. beans, salsa, peppers, egg beaters	Sandwich, low fat potato chips, 1c. juice, 2-4 lowfat cookies	Lg bowl low salt soup, 1c. rice, BBQ meat, salad & cooked vegs 1 glass wine	1 bowl of cereal
Avg BG 120's	**Avg BG 200's**	**Avg BG 200's**	**Avg BG 180's**

Diabetes Education SERVICES

Thank You

- Questions?
- Email
  bev@diabetesed.net
- Web
  www.diabetesed.net

Diabetes Education
SERVICES

Diabetes Bingo - "DiaBingo"
Shout out Right Answer

DiaBingo - N

N DPP demonstrated that exercise and diet reduced risk of DM by ___%
N An _____ a day can help prevent heart attack and stroke
N Rebound hyperglycemia
N Scare tactics are effective at motivating patients to change behavior
N Losing ___ % of body weight, can improve blood glucose, BP, lipids
N Drugs that can cause hyperglycemia
N 2/3 cups of rice equals _____ serving carbohydrate
N A1c of 7% equals glucose of
N One % drop in A1c reduces risk of complications by ___ %
N 1 gm of fat equal _____ kilo/calories

N Metabolic syndrome = hyperglycemia, hyperlipidemia, hypertension

N Average American consumes 25 teaspoons of sugar a day.