

Combination Oral Medications PocketCard™



Medications	Avail Doses in mg	Medications	Avail Doses in mg
ACTOplus Met* (pioglitazone/metformin)	15 / 500 15 / 850	Jentadueto (linagliptin/metformin)	2.5 / 500 2.5 / 850 or 2.5 / 1000
ACTOplus Met XR (pioglitazone/metformin)	15 / 1000 30 / 100	Kazano (alogliptin/metformin)	12.5 / 500 12.5 / 1000
Avandamet (rosiglitazone/metformin)	2 / 100	Kombligize XR (onglyza/metformin XR)	2.5 / 1000 5 / 500 or 5 / 1000
Duetact* (pioglitazone/glimepiride)	30 / 2 30 / 4	Metaglip* (glipizide/metformin)	2.5 / 250 2.5 / 500 or 5 / 500
Glucovance* (glyburide/metformin)	1.25 / 250 2.5 / 500 5 / 500	Oseni (aloglitpin/pioglitazone)	12.5/15 or 25 / 15 12.5 / 30 or 25 / 30 12.5 / 45 or 25 / 45
Glyxambi (empagliflozin and linagliptin)	10/5 25/5	Prandimet (repaglinide/metformin)	1 / 500 2 / 500
Invokamet (canagliflozin/metformin)	50 / 500 or 50 / 1000 150 / 500 or 150 / 1000	Synjardy (empagliflozin/metformin)	5 / 500 or 12.5 / 500 5 / 1000 or 12.5 / 1000
Janumet (sitagliptin/metformin)	50 / 500 50 / 1000	Synardy XR (empagliflozin/metformin XR)	5/1000 or 10/1000 12.5/1000 or 25/1000
Janumet XR (sitagliptin/metformin)	50 / 500 50 / 1000 or 100 / 1000	Xigduo XR (dapagliflozin/metformin)	5 / 500 or 10 / 500 5 / 1000 or 10 / 1000

**Available in generic. Observe precautions of each component drug. The information listed here are not guidelines. Please consult prescribing information for details.*

Other Oral Diabetes Meds

Download **FREE CDE® Coach App** for latest PocketCard versions and priority notifications | DiabetesEd.Net



Class/Main Action	Name(s)	Daily Dose Range	Considerations
Thiazolidinediones “TZDs” Increases insulin sensitivity	pioglitazone (Actos) rosiglitazone (Avandia)	15 – 45 mg daily 4 – 8 mg daily	Black Box Warning: TZDs may cause or worsen CHF. Monitor for edema and weight gain. Increased peripheral fracture risk. Actos may increase risk of bladder cancer. Lowers A1c 0.5% – 1.0%
Glucosidase Inhibitors Delays carb absorption	acarbose (Precose) miglitol (Glyset)	25 – 100 mg w/meals; 300 mg max daily dose	Start low dose, increase at 4-8 wk intervals to decrease GI effects. Caution with liver or kidney problems. In case of hypo, treat w/ glucose tabs. Lowers A1c 0.5– 1.0%.
Meglitinides Stimulates rapid insulin burst	repaglinide (Prandin)	0.5 – 4 mg w/meals (metabolized in liver)	Take before meals. Side effects may include hypoglycemia and weight gain.
	nateglinide (Starlix)	60 – 120 mg w/meals (eliminated via kidney)	Lowers A1c 1.0% – 2.0%.
Dopamine Receptor Agonists Resets circadian rhythm	bromocriptine mesylate— Quick Release “QR” (Cycloset)	1.6 to 4.8 mg a day (each tab 0.8 mg)	Take within 2 hrs of waking. Side effects: nausea, headache, fatigue, hypotension, syncope, somnolence. Lowers A1c 0.6% – 0.9%.
Bile Acid Sequestrants Decreases cholesterol / BG levels.	Colesevelam HCL (Welchol)	Up to six (6) 625 mg pills (3 tabs am, 3 tabs pm) 3.75gm packet in 4-8 ounces of fluid	Do not use if history of bowel obstruction, triglycerides >500, or pancreatitis. Can decrease absorption of certain meds, soluble vitamins. Lowers LDL by 15-30%. Side effects GI in nature. Lowers A1c 0.5%

Note: Content is for educational purposes only: please consult prescribing information for details.

REV 10/2016 © 2016