Exploring Diabetes – a Curiosity Based Approach

Beverly Thomassian, RN, MPH, BC-ADM, CDE
President, Diabetes Education Services

Schedule of Topics

- Where we are at with diabetes
- Patient Centered Approach
- Head to Toe



Spiritual Care

CDC Announces

35% of Americans will have Diabetes by 2050

Boyle, Thompson, Barker, Williamson
2010, Oct 22:8(1)29
www.pophealthmetrics.com

Diabetes Education SERVICES

Diabetes in America 2016

- 29 million or > 9.3%
- 27% don't know they have it
- 37% of US adults have pre diabetes (86 mil)



Diabetes Education SERVICES

Why Should Zip Code Determine Life Expectancy?

California Endowment – look up your zip code at www.measureofamerica.org

Diabetes Education SERVICES

Table 2: Estimated Diagnosed Diabetes Prevalence (%) 2012-2013, NM BRFSS

Has a doctor, nurse, or other health professional EVER told you that you have diabetes?

	Percent of Adults			Percent of Adults
New Mexico	9.6		**Health Region**	
Female	9.2		Northwest	12.1
Male	10.1		Northeast	7.3
			Metro	8.8
Age Group			Southeast	11.0
18-39 years	2.0		Southwest	10.9
40-59 years	12.1			
60 years & older	20.1			
Household Income			**Race/Ethnicity**	
< $15,000	15.2		American Indian/Alaska Native	18.1
$15-24,999	11.8		Asian/Native Hawaiian /Other Pacific Isle*	8.4
$25-34,999	11.1		Black/African American	9.0
$35-49,999	9.4		Hispanic	12.6
> $50,000	5.8		White	6.1

Data Source: New Mexico Department of Health, Behavioral Risk Factor Surveillance System (BRFSS), 2012 & 2013. Two years of data are used to increase the reliability of some of the rates. All rates except age group rates are age-adjusted to the 2000 US standard.

Diabetes Education SERVICES



Diabetes Education SERVICES

Type 2 in Kids

TODAY Treatment Options for type 2 Diabetes in Adolescents & Youth

- 7 fold increase 1990
- 1 in 6 overwt kids (age 12- 19) have prediabetes.
- ~2,500 to 3,700 new cases in U.S. annually.
- Highest risk: very obese, minority, female, low socioeconomic status, limited education
- In age range 12-19, less than 1% have Type 2 – NHANES
- Environmental changes to urgently needed

Diabetes Education SERVICES

Global Epidemic

- Every 10 seconds
  - 1 person dies with diabetes
  - 2 people develop diabetes
- Every year
  - 3 million deaths
  - 6 million new cases
- World Diabetes Day is November 14
- March is ADA Sound the Alert Day "find people w/ undetected diabetes"

Diabetes Education SERVICES

World Diabetes Day

November 14

Diabetes Education SERVICES

"The highest form of wisdom is kindness."
The Talmud

Diabetes Education Services
Published by Beverly Thomassian [?] · July 7 ·

Kindness matters!
Learning to be less harsh or judgmental and more compassionate to oneself may help people with diabetes manage their disease and stave off depression, a recent study suggests.

Self-compassion may help people with diabetes achieve better glucose control and less depression

By Reyna Gobel(Reuters Health) – Learning to be less harsh or judgmental and more...

REUTERS.COM | BY REYNA GOBEL

Diabetes Education SERVICES

Path to Type 2 Diabetes

Diabetes Education SERVICES

BMI Categories



Diabetes Education SERVICES

Patti Labelle

"divabetic"

"I have diabetes, it doesn't have me"

Diabetes Education SERVICES



2. Classification and DM Diagnosis

- Pre Diabetes & Type 2- Screening Guidelines
- Start screening at age 45 or for anyone who is overweight (BMI ≥ 25, Asians BMI ≥ 23) with one or > additional **risk factor**:
  - First-degree relative w/ diabetes
  - Member of a high-risk ethnic population
  - Habitual physical inactivity
  - PreDiabetes
  - History of heart disease

Diabetes Education SERVICES

Diabetes 2 - Who is at Risk?

(ADA Clinical Practice Guidelines)

Risk factors cont'd

- HTN - BP > 140/90
- HDL < 35 or triglycerides > 250
- baby >9 lb or history of Gestational Diabetes Mellitus
- Polycystic ovary syndrome (PCOS)
- Other conditions assoc w/ insulin resistance:
  - Severe obesity, acanthosis nigricans (AN)

Diabetes Education SERVICES

Acanthosis Nigricans (AN)

- Signals high insulin levels in bloodstream
- Patches of darkened skin over parts of body that bend or rub against each other
  - Neck, underarm, waistline, groin, knuckles, elbows, toes
  - Skin tags on neck and darkened areas around eyes, nose and cheeks.
- No cure, lesions regress with treatment of insulin resistance

Diabetes Education SERVICES

Acanthosis Nigricans

Diabetes Education SERVICES

Diabetes Detectives Needed

- On average – takes 6.5 years to diagnose diabetes
- 1/4 of all people with diabetes don't know they have it

Diabetes Education SERVICES

Natural Progression of Type 2 Diabetes



Adapted from Bergenstal et al. 2000; International Diabetes Center.

Diabetes Education SERVICES

Natural History of Diabetes



Multiple, Complex Pathophysiological Abnormalities in T2DM

Adapted from: Inzucchi SE, Sherwin RS in: *Cecil Medicine* 2011



Diabetes Education SERVICES

DiaBingo

B Frequent skin and yeast infections
B A BMI of ____ or greater is considered overweight
B To reduce complications, control **A**1c, **B**lood pressure, **C**holesterol
B PreDiabetes – fasting glucose level of ___ to ____
B Erectile dysfunction indicates greater risk for ____
B Diabetes – fasting glucose level____ or greater
B Type 1 diabetes is best described as an _____ disease
B People with diabetes are _____ times more likely to die of heart dx
B Elevated triglycerides, < HDL, smaller dense LDL
B Each percentage point of A1c = _____ mg/dl glucose
B At dx of type 2, about __% of the beta cell function is lost
B Diabetes – random glucose ____ or greater

Diabetes Education SERVICES

Life Study – Mrs. Jones

Mrs. Jones is 62 years old, overweight and complaining of feeling tired and urinating several times a night. She is admitted with a urinary tract Infection. Her WBC is 12.3, glucose 237. She is hypertensive with a history of gestational diabetes. No ketones in urine.

- What are her risk factors, signs of diabetes
- What type of diabetes does she have?
- Does she have insulin resistance?

What Do You Say? Mrs. Jones asks you

- What is type 2 diabetes?
- Will this go away?
- Will I get complications?
- Will I need to take diabetes medication for the rest of my life?
- How come I got diabetes?
- Do I have to check my blood sugars?

Diabetes Education SERVICES

Strategies – One Baby Step at a Time

Pause

Diabetes Education SERVICES

No one is Unmotivated

…. to lead and long and healthy life

- **These are the 3 usual Critical Barriers**
  - Perceived worthlessness
  - Too many personal obstacles
  - Absence of support and resources

Bill Polonsky, PhD, CDE

Diabetes Education SERVICES

Overcoming barriers

- Confront the key misbelief. Ask the question, does dm cause complications?
- Offer pts evidence based hope message –
- Frequent contact
- Paired glucose testing
- Ask pt, "Tell me 1 thing that is driving you crazy about your diabetes"
- Discuss medication beliefs
- To improve outcomes, see pts more often

Bill Polonsky, PhD, CDE

Diabetes Education SERVICES

How Often Should I Check?

- Be realistic!!
- Type 2 on orals – Medicare covers 100 strips for 3 months
- Based on individual - Consider:
  - Types and timing of meds
  - Goals
  - Ability (physical and emotional)
  - Finances / Insurance

Diabetes Education SERVICES

Give the gift of Non-Judgment



Diabetes Education SERVICES

Complications - Why?

- Degree of hyperglycemia "glucose toxicity"
- Duration of hyperglycemia
- Genes
- Multiple risk factors: smoking, vascular disease, dyslipidemia, hypertension, other

Diabetes Education SERVICES

Control Matters

- **Prevention**
- **Trials**
- **Practice Recommendations**

Diabetes Education SERVICES

Financial Advisor

- Mid 30s, friendly, he smiles to greet you and you notice his gums are inflamed. You'd guess a BMI of 26 or so, with most of the extra weight in the waist area.
- If you could give him some health related suggestions, what would they be?

Diabetes Education SERVICES

Can Type 2 be Prevented in Older Adults?

- Physical activity (30 mins a day)
- Dietary score (higher fiber intake, low saturated fat and *trans*-fat, lower mean glycemic index)
- Not Smoking
- Alcohol use (up to 2 drinks a day);
- BMI <25 and waist circumference

Dariush Mozaffarian, MD,
Arch Intern Med. 2009;169(8):798-807.

Overall, 9 of 10 new cases of diabetes attributable to these 5 lifestyle factors.

89% risk reduction when all at goal.

35% rel risk reduction for each additional

Diabetes Education SERVICES

Exercise – Getting Active

- "Physical activity is the closest thing we have to a wonder drug,"
- "Being active is one of the most important things people of all ages, sizes, and shapes can do to improve their health."
- *Dr. Thomas Frieden, Director of CDC*

Diabetes Education SERVICES

Exercise Recommendations

- **Activity update –Don't sit more than 90 minutes**
- Evidence supports that everyone, including with diabetes should be encouraged to reduce sedentary time, by not sitting for more than 90 minutes at a time.
- It is recommended that people with pre diabetes and diabetes engage in 150 minutes of activity a week and at **least 2 weekly sessions of resistance exercise.**

Diabetes Education SERVICES

Good Exercise Info / Quotes

- 20 % of people walk 30 mins a day
- Exercise decrease A1c 0.7%
- No change in body wt, but 48% loss in visceral fat
  - ADA PostGrad 2010

- "If you don't have time for exercise, you better make time for disease."

"I don't have time to exercise, I MAKE time."

Mike Huckabee

Can we stop pre diabetes from progressing?

3, 234 people w/ Pre-Diabetes randomized:

- Placebo
- Diet/Exercise or
- Metformin

over a three year period

Diabetes Prevention Program (DPP) 2001

NATIONAL Diabetes PREVENTION PROGRAM

Diabetes Education SERVICES

Diabetes Prevention Program

- Standard Group - 29% developed DM
- Lifestyle Results - 14% developed DM
  - 58% (71% for 60yrs +) Risk reduction
    - 30 mins daily activity
    - 5-7% of body wt loss
- Metformin 850 BID - 22% developed DM
  - 31% risk reduction (less effective with elderly and thinner pt's)

Diabetes Education SERVICES

Weight loss and Prevention

- For every 2.2 pounds of weight loss, risk of type 2 diabetes was reduced by 13%.

NATIONAL Diabetes PREVENTION PROGRAM

Diabetes Education SERVICES



ABCs of Diabetes –

- A1c less than 7% (avg 3 month BG)
  - Pre-meal BG 80-130
  - Post meal BG <180
- Blood Pressure < 140/90
- Cholesterol
  - DM and 40 yrs, start statin
  - HDL >40
  - Triglyceride < 150
- E xercise, Education
- H ealthy Eating

Diabetes Education SERVICES

Glycemic Targets - ADA

- **Adult non pregnant A1c goals**
  - **A1c < 7%** - a reasonable goal for adults.
  - **A1c < 6.5%** - may be appropriate for those without significant risk of hypoglycemia or other adverse effects of treatment.
  - **A1c < 8%** - may be appropriate for patients with history of hypoglycemia, limited life expectancy, or those with longstanding diabetes and vascular complications.



Diabetes Education SERVICES

A1c and Estimated Avg Glucose (eAG) 2008

A1c (%)	eAG
5	97
6	126
7	154
8	183
9	212
10	240
11	269
12	298

Order teaching tool kit free at diabetes.org

eAG = 28.7 x A1c-46.7 ~ 29 pts per 1%

Translating the A1c Assay Into Estimated Average Glucose Values – ADAG Study
Diabetes Care: 31, #8, August 2008

Diabetes Education SERVICES

Older Adults - Considerations

- Reduced life expectancy
- Higher CVD burden
- Reduced GFR
- At risk for adverse events from polypharmacy
- More likely to be compromised from hypoglycemia

- ✓**Less ambitious targets**
- ✓**A1c <7.5–8.0%**
- ✓**Focus on drug safety**

Diabetes Care 2012;35:1364–1379
Diabetologia 2012;55:1577–1596

Diabetes Education SERVICES

Diabetes and the Cairn



Diabetes Education SERVICES

DiaBingo- G

G ADA goal for A1c is less than ____%

G People with DM need to see their provider at least every month

G Blood pressure goal is less than

G People with DM should see eye doctor (ophthalmologist) at least

G The goal for triglyceride level is less than

G Goal for my HDL cholesterol is more than

G The goal for blood sugars 1-2 hours after a meal is less than:

G People with DM should get this shot every year

G People with DM need to get urine tested yearly for _____

G Periodontal disease indicates increased risk for heart disease

G The goal for blood sugar levels before meals is:

G The activity goal is to do ___ minutes on most days

Diabetes Education SERVICES

Antihyperglycemic Therapy – 1st Step

- Lifestyle Changes
  - Weight control
  - Healthy eating
  - Activity

ADA-EASD Position Statement: Management of Hyperglycemia in T2DM

Diabetes Care **2012;35:1364–1379**
Diabetologia **2012;55:1577–1596**

Diabetes Education SERVICES

Patient Centered Approach

"...providing care that is respectful of and responsive to individual patient preferences, needs, and values - ensuring that patient values guide all clinical decisions."

- **Gauge patient's preferred level of involvement.**
- **Explore, where possible, therapeutic choices.**
- **Utilize decision aids.**
- **<u>Shared</u> decision making – final decisions re: lifestyle choices ultimately lie with the patient.**

ADA-EASD Position Statement: Management of Hyperglycemia in T2DM

Diabetes Care **2012;35:1364–1379**
Diabetologia **2012;55:1577–1596**

Diabetes Education SERVICES



Figure 6.1—Depicted are patient and disease factors used to determine optimal A1C targets. Characteristics and predicaments toward the left justify more stringent efforts to lower A1C; those toward the right suggest less stringent efforts. Adapted with permission from Inzucchi et al. (45).

ADA Standards of Care 2015

Diabetes Education SERVICES

Treating Hyperglycemia with Meds

- For all of the following case studies, we assume we are providing ongoing education on lifestyle – including referral to the Diabetes Ed Team and other resources.
- In describing what meds match the patient best, we act as patient advocates and consultants to providers.

Diabetes Education SERVICES

Life Study

- 61 year old overweight woman with type 2 diabetes 3 months. Has been trying to control diabetes with diet and exercise. GFR in 90s. Worried about weight gain.
- Most recent A1c 6.9%
  - ADA
  - AACE
  - Cash pay or
  - Good insurance

Diabetes Education SERVICES

ADA Step Wise Approach to Hyperglycemia 2015



- Start with lifestyle coaching
- When lifestyle alone is not achieving A1c goal – Metformin should be added at, or soon after diagnosis (unless contraindicated).
- Metformin has a long standing evidence base for efficacy and safety, is cheap and may reduce CV risk.

Diabetes Education SERVICES

Other Considerations

- Cost
- Hypoglycemia
- Age
- Weight
- Comorbidities
  - Kidney disease
  - Heart disease – CHF, CAD
  - Liver dysfunction

ADA-EASD Position Statement: Management of Hyperglycemia in T2DM

Diabetes Care **2012;35:1364–1379**
Diabetologia **2012;55:1577–1596**

Diabetes Education SERVICES

Oral Diabetes Medications

Diabetes Education SERVICES — 15 years

Class/Main Action	Name(s)	Daily Dose Range	Considerations
Biguanides • Decrease hepatic glucose output • First line med at diagnosis of type 2	metformin (Glucophage)	500 – 2500 mg (usually BID w/meal)	Side effects: nausea, bloating, diarrhea. Use XR to minimize. Lactic acidosis precaution: avoid in pts with creat >1.4 women, 1.5 men, during illness or surgery. Benefits: decreased cholesterol, no wt gain or hypoglycemia. Lowers A1c 1.0% – 2.0%.
	Extended Release-XR (Glucophage XR) (Glumetza) (Fortamet)	(1x daily w/dinner) 500 – 2000 mg 500 – 2000 mg 500 – 2500 mg	
Sulfonylureas • Stimulates sustained insulin release	glyburide: (Micronase, Diabeta) (Glynase)	1.25 – 20 mg 0.75 – 12 mg	Can take once or twice daily before meals. Side effects include hypoglycemia and weight gain. Eliminated via kidney. **Caution:** Glyburide most likely to cause hypoglycemia. Lowers A1c 1.0% – 2.0%.
	glipizide: (Glucotrol) (Glucotrol XL)	2.5 – 40 mg 2.5 – 20 mg	
	glimepiride (Amaryl)	1.0 – 8 mg	
DPP – 4 Inhibitors • "Incretin Enhancers" • Prolongs action of gut hormones • Increases insulin secretion • Delays gastric emptying	sitagliptin (Januvia)	100 mg daily (eliminated via kidney*)	*If creatinine elevated, see pkg insert for dosing info. No wt gain or hypoglycemia. Side effects include nasopharyngitis, headache and upper-respiratory tract infection. Report signs of pancreatitis (abdominal pain, nausea, vomiting). Lowers A1c 0.6% – 0.8%.
	saxagliptin (Onglyza)	Up to 5 mg daily (eliminated via kidney*, feces)	
	linagliptin (Tradjenta)	5 mg daily (eliminated via feces)	
	alogliptin (Nesina)	25 mg once daily (eliminated via kidney)	

More medications on back. Note: These meds are for people with Type 2 diabetes and should not be used during pregnancy. Content is for educational purposes only: please consult prescribing information for details.

REV 01/2015 ©2015

Diabetes Education SERVICES

A Diabetes PocketCard® from Diabetes Education Services

Class/Main Action	Name(s)	Daily Dose Range	Considerations
SGLT2 Inhibitors • Decrease glucose reabsorption in kidneys • "Glucoretic"	Canagliflozin (Invokana) Dapagliflozin (Farxiga) Empagliflozin (Jardiance)	100 – 300 mg 1x daily 5 – 10 mg 1x daily 10 – 25 mg 1x daily	For all, monitor B/P, K+ and renal function. If GFR<45, stop Invokana. If GFR<60, stop Farxiga. Do not start pts w/ GFR<45 on Jardiance. Side effects: hypotension, UTIs, increased urination, genital infections. Avoid Farxiga in pts w/ bladder cancer. Lowers A1c 0.7% – 1.5%, lowers wt 1 – 3 lbs.
Thiazolidinediones "TZDs" • Increase insulin sensitivity	pioglitazone (Actos) rosiglitazone (Avandia)	15 – 45 mg daily 4 – 8 mg daily	**Black Box Warning:** TZDs may cause or worsen CHF. Monitor for edema and weight gain. Increased peripheral fracture risk. **Actos may increase risk of bladder cancer.** Lowers A1c 0.5% – 1.0%
Glucosidase Inhibitors • Delay carb absorption	acarbose (Precose) miglitol (Glyset)	25 – 100 mg w/meals; 300 mg max daily dose	Start low dose, increase at 4-8 wk intervals to decrease GI effects. Caution with liver or kidney problems. In case of hypo, treat w/ glucose tabs. Lowers A1c 0.5– 1.0%.
Dopamine Receptor Agonists • Resets circadian rhythm	bromocriptine mesylate—Quick Release "QR" (Cycloset)	1.6 to 4.8 mg a day (each tab 0.8 mg)	Take within 2 hrs of waking. Side effects: nausea, headache, fatigue, hypotension, syncope, somnolence. Lowers A1c 0.6% – 0.9%.
Meglitinides • Stimulates rapid insulin burst	repaglinide (Prandin) nateglinide (Starlix)	0.5 – 4 mg w/meals (metabolized in liver) 60 – 120 mg w/meals (eliminated via kidney)	Take before meals. Side effects may include hypoglycemia and weight gain. Lowers A1c 1.0% – 2.0%.

Diabetes Education Services | Advancing Your Career in Diabetes Education | Visit DiabetesEd.net

Diabetes Education Services
(530) 893-8635
Beverly Dyck Thomassian
RN, MPH, BC-ADM, CDE

Diabetes Education SERVICES

Injectables That Lower Glucose

Class/Main Action	Name	Dose Range	Considerations
GLP-1 Agonist "Incretin Mimetic" • Increases insulin release with food • Slows gastric emptying • Promotes satiety • Suppresses glucagon Lowers A1c 0.5 – 1.6% Wt loss of ~ 3lbs	exenatide (Byetta)	5 or 10 mcg BID (renally excreted)	**Side effects for all:** Nausea, vomiting, weight loss, injection site reaction. Report signs of acute pancreatitis (severe abdominal pain, vomiting), stop med. **Black box:** Thyroid C-cell tumor warning for liraglutide, exenatide XR, albiglutide, and dulaglutide (avoid if family history of medullary thyroid cancer, notify MD of hoarseness, throat lump).
	exenatide XR (Bydureon)	2mg 1x a week (renally excreted)	
	liraglutide (Victoza)	0.6 - 1.8 mg daily	
	albiglutide (Tanzeum)	30 and 50 mg 1x a week pen injector	
	dulaglutide (Trulicity)	0.75 and 1.5 mg 1x a week pen injector	
Amylin Mimetic • Slows gastric emptying • Supresses glucagon • Promotes satiety Lowers A1c 0.5 – 1%	pramlintide (Symlin)	Type 1: 15 - 60 mcg; Type 2: 60 - 120 mcg immediately before major meals	For Type 1 or 2 on insulin. **Black box warning:** severe hypoglycemic risk 3 hrs post injection. Prevent hypoglycemia, decrease insulin dose when starting pramlintide. Side effects: nausea, weight loss.

The information listed here are general guidelines only; please consult prescribing information for details.

Diabetes Education SERVICES

CDE® Coach App – Download Success



Diabetes Education SERVICES



ADA Standards of Care 2015



When goal is to avoid weight gain

- These meds are weight neutral
  - Metformin
  - DPP-IV Inhibitors: Januvia, Onglyza, Tradjenta, Nesina
  - Acarbose
- These meds associated with wt loss
  - GLP-1 agonists (Byetta, Bydureon, Victoza, Tanzeum, Trulicity)
  - SGLT-2 Inhibitors (Canagliflozin, Dapagliflozin, Empagliflozin)
  - Symlin (Pramlintide)

When goal is to minimize cost

- Go generic.
- Oral Meds -Metformin and Sulfonylureas
  - Walmart offers 3 mo supply of following meds for ~ $10
    - Metformin and Metformin XR
    - Glipizide, Glyburide, Glimepiride
- Insulins – Oldies but Goodies
  - NPH, Regular, 70/30 mix
  - $25 a vial at Walmart – ReliOn
  - Vials and needles cheaper

Diabetes Education SERVICES

Life Study

- 61 year old overweight woman with type 2 diabetes 3 months. Has been trying to control diabetes with diet and exercise. GFR in 90s. Worried about weight gain.
- Most recent A1c 6.9%
  - ADA
  - AACE
  - Cash pay
  - Good insurance
- Solutions?

Diabetes Education SERVICES

Life Study

- 61 year old overweight woman with type 2 diabetes 3 months. Has been trying to control diabetes with diet and exercise. GFR in 90s. Worried about weight gain.
- Most recent A1c 6.9%
  - ADA
  - AACE
  - Cash pay
  - Good insurance
- Solutions?
  - Consider no meds, monitor or start Metformin (ADA - Cash)
  - Start Metformin 500 mg 1-2 x a day (AACE)
  - If can't tolerate metformin, good insurance, consider?

Diabetes Education SERVICES

Life Study

- 54 year old smoker, has CVD disease, creatinine 1.2, BMI 27. Not checking BG, even though he has glucose meter. On Metformin 500mg BID for past 4 months. Had bad experience with hypoglycemia on glyburide.
- Most recent A1c 7.9%
  - ADA
  - AACE
  - Cash pay
  - Good insurance

Diabetes Education SERVICES

When goal is to avoid Hypoglycemia

- Avoid sulfonylureas
- Careful insulin dosing
- May need to up adjust glucose goals
- Monitor kidney function
- Reinforce for patients on insulin to "TIE"
  - Test
  - Inject
  - Eat

Diabetes Education SERVICES

Life Study

- 54 year old smoker, has CVD disease, creatinine 1.2, BMI 27. Not checking BG, even though he has glucose meter. On Metformin 500mg BID for past 4 months. Had bad experience with hypoglycemia on glyburide.
- Most recent A1c 7.9%
- Solution:
  - Change to Metformin XR and double dose
    - Good insurance? Add GLP-1 or SGLT-2

If cash pay, increase Metformin, if needed add insulin or glipizide

Diabetes Education SERVICES

Diabetes Vacations



Diabetes Education SERVICES

Other Meds Needed?

- 54 year old smoker, has CVD disease, creatinine 1.2, BMI 27. Not checking BG, even though he has glucose meter. On Metformin 500mg BID for past 4 months..
- Most recent A1c 7.9%
- Other Meds to protect against CVD?
  - In addition to LifeStyle Coaching

Diabetes Education SERVICES

8. Coronary Heart Disease

- In pts with known CVD, use:
  - Aspirin
  - Statin
  - B/P Med
    - Consider ACE Inhibitor to reduce risk of CV event
    - In pts with prior MI, Beta Blockers should be continued at least 2 years after the event
  - Don't use Actos or Avandia in pts with CHF
  - In pts with stable CHF, Metformin can be used in renal function normal and stable

Diabetes Education SERVICES

Life Study

- 71 year old woman with type 2 diabetes for past year. BMI 24. Has been trying to control diabetes by limiting carbs and exercise. Creat 1.6. Good social support.
- Most recent A1c 8.6%
  - She has great insurance or
  - She is cash pay or
  - She hates needles

Diabetes Education SERVICES

Life Study

- 71 year old woman type 2 diabetes. BMI 24. Has been trying to control diabetes by limiting carbs and exercise. Creat 1.6. GFR low 30s. Good social support.
- Most recent A1c 8.6%
- Solutions
  - Great insurance – GLP1, DPP-IV Inhibitor, Analog insulin
  - She is cash pay – Sulfonylurea or NPH or 70/30
  - She hates needles – Sulfonylurea, DPP-IV Inhibitor - if doesn't work, see if she will reconsider insulin, GLP1

Diabetes Education SERVICES

Pattern Management

- Safety 1st!! - Evaluate 3 day patterns
- **Hypo:** eval 1st and fix:
  - If possible, decrease medication dose
  - Timing of meals, exercise, medications
- **Hyperglycemia:** evaluate 2nd
  - Identify patterns
  - Before increase insulin, make sure not missing something (carbs, exercise, omission)

Diabetes

80 /20 Rule – Perfect Not Required



Diabetes Education SERVICES

DiaBingo - N

N DPP demonstrated that exercise and diet reduced risk of DM by ___%
N An _____a day can help prevent heart attack and stroke
N Rebound hyperglycemia
N Scare tactics are effective at motivating patients to change behavior
N Losing ___ % of body weight, can improve blood glucose, BP, lipids
N Drugs that can cause hyperglycemia
N 2/3 cups of rice equals _____ serving carbohydrate
N A1c of 7% equals glucose of
N One % drop in A1c reduces risk of complications by ___ %
N 1 gm of fat equal _____kilo/calories

N Metabolic syndrome = hyperglycemia, hyperlipidemia, hypertension

N Average American consumes 25 teaspoons of sugar a day.

Diabetes Education SERVICES

Case Study

- 68 yr old, avid walker
- BMI 24, Weighs 90kg
- A1c – 8.6%, BG 270s during day for past wks
- On glyburide 10mg BID
- Doesn't have a glucose meter
- Tells you, "my foot hurts"

Diabetes Education SERVICES

Diabetes and Lower Extremity Ulcers

- Up to 15% of DM patients have ulcers in their lifetime
- Mortality with foot ulcers is twice usual

Lower Extremity Amputations Dropping over past 10yrs

- 60% of amputations in 7% of pop
- Higher in men, elderly, minorities, Chronic Kidney Disease (CKD)
- Lower extremity complications represent 20% of hospitalizations for elderly
- Amputations cost $40,000
- Amputation associated w/ earlier death compared to revascularization
- 10 yr survival after LEA

Risk factors for Foot Ulcers/ Amputation



- Previous amputation
- Past foot ulcer history
- Peripheral neuropathy
- Foot deformity
- Peripheral vascular disease
- Visual impairment
- Diabetic nephropathy (especially patients on dialysis)
- Poor glycemic control
- Cigarette smoking
- ADA Task Force - 2008

Pitting Edema

Peripheral Arterial Disease Assessment

Diabetes Education SERVICES

No Bathroom Surgery



You Can Make A Difference

- Assess
  - Nail condition, nail care, in between the toes
  - Who trims your nails
  - Have you ever cut your self?
  - Shoes – type and how often
  - Socks
  - Skin/skin care and vascular health
  - Ability to inspect
  - Loss of protective sensation

Diabetes Education SERVICES

Three Most Important Foot Care Tips

- Inspect and apply lotion to your feet every night before you go to bed.
- Do NOT go barefoot, even in your house. Always wear shoes!
- Every time you see your doctor, take off your shoes and show your feet. Report any foot problems right away!

Diabetes Education SERVICES

"Getting diabetes saved my life." ~ Sherri Sheperd

Sherri Shepard decided to embrace diabetes and use it as a motivator to improve her health.

Diabetes Education SERVICES

Patient Perspective Matters!

- **"I am no where."**

- **"I am now here!"**

PERHAPS FIGURING OUT
WHAT MATTERS MOST
MATTERS MOST

Diabetes Education SERVICES

Thank You

Diabetes Education SERVICES

www.DiabetesEd.net

Diabetes Education SERVICES