



Welcome to Diabetes in the 21st Century

Beverly Dyck Thomassian, RN, MPH, BC-ADM, CDE
President, Diabetes Education Services
www.DiabetesEd.net



Karen Cole, BSN, RN, CWOCN,
CDE



- Certified Diabetes Educator
- Claxton-Hepburn Medical Center Diabetes Education Program
- Serving St. Lawrence County

Shannon VanHouse, RN, CWOCN



Certified Diabetes Educator
Candidate

Claxton-Hepburn Medical Center
Diabetes Education Program
Ogdensburg, NY

315-713-5116
svanhouse@chmed.org

Serving all of St. Lawrence
County

Healthy Lifestyles
1575 Washington Street
Watertown, NY 13601
Tel: 315-785-4687
Fax: 315-785-7310



The Healthy Lifestyles Program Includes:

1. Medical Nutrition Therapy (MNT) for a variety of medical conditions, including but not limited to:
 - ❖ Diabetes
 - ❖ Renal Disease
 - ❖ Weight Management
 - ❖ Gastrointestinal Disorders
2. Diabetes Self Management Education (DSME) classes are organized by our Registered Dietitian and Registered Nurse for patients with a diagnosis of Diabetes. Patients with Diabetes should be referred for both services (MNT & DSME) at diagnosis, annually, when new complications or transitions in care arise.

Teresa Intorcja, RDN, CDE
Julie Stone, RN, Diabetes Educator





Nutrition Services Outpatient Care
Medical Nutrition Therapy and counseling is offered for medical conditions, including but not limited to:

- Diabetes
- Renal Disease
- Weight Management
- Gastrointestinal Disorders
- Nutritional Support (oral, enteral, parenteral)

Diabetes Self-Management Education Program, accredited by the American Association of Diabetes Educators, is available for clients at any stage of their diagnosis of diabetes. Class schedule and time varies.

Our Providers:



Carly R. Draper RD, CDN, CDE Susan Kim Ray MS, RD, CNSC, CDE Rebecca Redmore MSEd, RD, CDN

Contact Information:
1001 West Street
Carthage, NY 13619
P: (315) 493-1000 Ext 5589
F: (315) 493-0038
Office Hours: 8am to 4pm M- F

Diabetes in the 21st Century:
A Clinical and Educational Update

1. Describe impact of diabetes
2. Discuss prevention, management strategies
3. Discuss different types of diabetes
4. Describe insulin therapy
5. Gain understanding of Type 2 Meds.
6. Review glucose patterns and determine how to adjust therapy to improve glucose.
7. Discuss gut bacteria and healthy eating
8. Demonstrate successful teaching strategies



CDC Announces



35% of
Americans will
have Diabetes
by 2050

Boyle, Thompson, Barker, Williamson
2010, Oct 22:8(1)29
www.pophealthmetrics.com



Diabetes Education
SERVICES

Diabetes in America 2018


- ▶ 30.3 million or > 9.4%
- ▶ 24% don't know they have it
- ▶ 34 % of US adults have pre diabetes (84 mil)
- ▶ Increasing rates 3 key factors
 - ▶ Aging of U.S. Population
 - ▶ Increasing size of higher-risk minority populations
 - ▶ Declining mortality among those with diabetes



Diabetes Education
SERVICES

Age-Adjusted Prevalence of Obesity and Diagnosed Diabetes Among US Adults

2013



Diabetes Education
SERVICES

Global Epidemic


- ▶ Every 10 seconds
 - ▶ 1 person dies with diabetes
 - ▶ 2 people develop diabetes
- ▶ Every year
 - ▶ 3 million deaths
 - ▶ 6 million new cases
- ▶ World Diabetes Day is November 14
- ▶ March is ADA Sound the Alert Day “find people w/ undetected diabetes”



Diabetes Education SERVICES

World Diabetes Day

November 14




Diabetes Education SERVICES

Age-adjusted Diabetes Prevalence

20 yrs or older, by race/ethnicity— U.S. 2014

Age-adjusted* percentage of people aged 20 years or older with diagnosed diabetes, by race/ethnicity, United States, 2010–2012



*Based on the 2000 U.S. standard population.
Source: 2010–2012 National Health Interview Survey and 2012 Indian Health Service's National Patient Information Reporting System.

- Among Hispanic adults, the age-adjusted rate of diagnosed diabetes was 8.5% for Central and South Americans, 9.3% for Cubans, 13.9% for Mexican Americans, and 14.8% for Puerto Ricans.
- Among Asian American adults, the age-adjusted rate of diagnosed diabetes was 4.4% for Chinese, 11.3% for Filipinos, 13.0% for Asian Indians, and 8.8% for other Asians.
- Among American Indian and Alaska Native adults, the age-adjusted rate of diagnosed diabetes varied by region from 6.0% among Alaska Natives to 24.1% among American Indians in southern Arizona.



Diabetes Education SERVICES

Why Should Zip Code Determine Life Expectancy?



California Endowment – look up your zip code at www.measureofamerica.org




Diabetes Education
SERVICES

Now, let's get to the Nitty Gritty



Diabetes Education
SERVICES




Hormones Effect on Glucose

Hormone	Effect
▶ Glucagon (pancreas)	↑
▶ Stress hormones (kidney)	↑
▶ Epinephrine (kidney)	↑
▶ Insulin (pancreas)	↓
▶ Amylin (pancreas)	↓
▶ Gut hormones - incretins (GLP-1) released by L cells of intestinal mucosa, beta cell has receptors	↓



Diabetes Education SERVICES

GLP-1 Effects in Humans Understanding the Natural Role of Incretins



Adapted from Flint A, et al. *J Clin Invest*. 1998;101:515-520
Adapted from Larsson W, et al. *Acta Physiol Scand*. 1997;160:413-422
Adapted from Nauck MA, et al. *Diabetologia*. 1996;39:1546-1553
Adapted from Drucker DJ. *Diabetes*. 1998;47:159-169

GLP-1 degraded by DPP-4 w/in minutes



Diabetes Education SERVICES

Injectables that Lower Glucose

Class/Main Action	Name	Dose Range	Considerations
GLP-1 Receptor Agonist (GLP-1 RA) "Incretin Mimetic" <ul style="list-style-type: none"> Increases insulin release with food Slows gastric emptying Promotes satiety Suppresses glucagon 	exenatide (Byetta)	5 and 10 mcg BID	Side effects for all: Nausea, vomiting, weight loss, injection site reaction. Report signs of acute pancreatitis (severe abdominal pain, vomiting), stop med. Renally excreted. Black box warning: Thyroid C-cell tumor warning for exenatide XR, liraglutide, dulaglutide, and semaglutide (avoid if family history of medullary thyroid tumor). * Victoza significantly reduces risk of CV death, heart attack, and stroke. Lowers A1c 0.5 - 1.6% Weight loss of 1.6 to 6.0kg†
	exenatide XR (Bydureon)	2mg 1x a week	
	Pen injector - Bydureon BCise		
	liraglutide (Victoza)*	0.6 and 1.8 mg daily	
	dulaglutide (Trulicity)	0.75 and 1.5 mg 1x a week pen injector	
	semaglutide (Ozempic)†	0.5 and 1.0 mg 1x a week pen injector	
lixisenatide (Adlyxin)	(Only available in combo with glargine, IGLarLixi, see below)		



Diabetes Education SERVICES

CDE® Coach App – Download Success



Coach in your pocket.
Med Pocket Cards. Resources. Courses.

Get it on  **Google play**

Download on the  **App Store**





Loading...
SUCCESS!

Diabetes Education SERVICES


Bariatric Surgery

- ▶ Consider on diabetes pts w/ BMI >30, esp with comorbidities
- ▶ Remission (BG normalized)
 - ▶ rates range from 40 – 95%
 - ▶ Better results with newer diabetes (more beta cell mass)
 - ▶ Due to increase incretins (gut hormones)
- ▶ Still researching long term benefits, cost effectiveness and risk



Diabetes Education SERVICES

Natural History of Diabetes

 <p>Normal FBG <100 Random <140 A1c <5.7%</p>	 <p>Prediabetes FBG 100-125 Random 140 - 199 A1c ~ 5.7- 6.4% 50% working pancreas</p>	 <p>Diabetes FBG 126 + Random 200 + A1c 6.5% or + 20% working pancreas</p>
<p>← Yes! NO →</p>		
<p>← Development of type 2 diabetes happens over years or decades →</p>		

Signs of Diabetes



- ▶ Polyuria
- ▶ Polydipsia
- ▶ Polyphasia
- ▶ Weight loss
- ▶ Fatigue
- ▶ Skin and other infections
- ▶ Blurry vision
- ▶ Glycosuria, H₂O losses
- ▶ Dehydration
- ▶ Fuel Depletion
- ▶ Loss of body tissue, H₂O
- ▶ Poor energy utilization
- ▶ Hyperglycemia increases incidence of infection
- ▶ Osmotic changes



Diabetes Education
SERVICES

Diabetes Classifications

- ▶ Type 1
- ▶ Type 2
- ▶ Gestational
- ▶ Secondary



Diabetes Education
SERVICES

Case Study

1. Pt profile: 5'8", 192 lb male with history of schizophrenia

Diabetes 12 years, on insulin 3 yrs since starting on olanzapine (Zyprexa).



What type of DM and how do you know?

What is the link between olanzapine and glucose levels?



Diabetes Education
SERVICES

Schizophrenia and Diabetes - Link

- ▶ People with schizophrenia 3xs more likely to have diabetes.
- ▶ Links may be related to:
 - ▶ premature birth
 - ▶ low birth weight
 - ▶ Both are recognized as risk factors for both schizophrenia and diabetes.
- ▶ Raised levels of stress hormone, cortisol, is also a risk factor for diabetes.
- ▶ It is possible that the stress associated with developing schizophrenia, with increased cortisol levels, may also contribute to higher diabetes risk.



Diabetes Education SERVICES

Novel Antipsychotics Hyperglycemia, Wt gain, Cholesterol

- ▶ Zyprexa – olanzapine
 - ▶ Geodon - ziprasidone
 - ▶ Seroquel – quetiapine
 - ▶ Risperdal - risperadone
 - ▶ Clozaril - clozapine
 - ▶ Abilify – aripiprazole
- ▶ Check BG on initiation, with symptoms of hyper and yearly
- ▶ Meds tend to cause weight gain, hyperglycemia, high cholesterol, other metabolic problems.



Diabetes Monitoring – Schizophrenia (HEDIS)

Monitor	Numerator	Denominator
Diabetes Monitoring for People with Diabetes and Schizophrenia	Number of people who had both an LDL test and A1c test during measurement year	Number of people, ages 18 to 64 years, with schizophrenia and diabetes
Diabetes Screening for People with Schizophrenia, BPD Using Antipsychotic Med.	Number of people who had a diabetes screening test during measurement year	Number of people, ages 18 to 64 years, with schizophrenia or bipolar disorder, who were dispensed an antipsychotic medication



Diabetes Education SERVICES

Case Study

5'6", 108 lb female

On insulin 3u Regular before meals,
10u Lantus at bedtime

What type of DM and how do you know?

Is there anything else you are worried about?



Diabetes Education
SERVICES

Type 1 Rates Increasing Globally

- ▶ 23% rise in type 1 diabetes incidence from 2001-2009
- ▶ Why?
 - ▶ Autoimmune disease rates increasing over all
 - ▶ Changes in environmental exposure and gut bacteria?
 - ▶ Hygiene hypothesis
 - ▶ Obesity?



Diabetes Education
SERVICES

Incidence of Type 1 in Youth



- ▶ **General Pop 0.3%**
- ▶ **Sibling 4%**
- ▶ **Mother 2-3%**
- ▶ **Father 6-8%**
- ▶ Rate doubling every 20 yrs
- ▶ Many trials underway to detect and prevent (Trial Net)



Diabetes Education
SERVICES

Type 1 – 10% of all Diabetes
Genetics and Risk Factors

- ▶ Auto-immune pancreatic beta cells destruction
- ▶ Most commonly expressed at age 10-14
- ▶ Insulin sensitive (require 0.5 - 1.0 units/kg/day)

- ✱ Combo of genes and environment:
 - ✱ Autoimmunity tends to run in families
 - ✱ Higher rates in non breastfed infants
 - Ⓜ Viral triggers: congenital rubella, coxsackie virus B, cytomegalovirus, adenovirus and mumps.



Autoantibodies Assoc w/ Type 1

Panel of autoantibodies –

- ▶ GAD65 - Glutamic acid decarboxylase –
- ▶ ICA - Islet Cell Cytoplasmic Autoantibodies
- ▶ IAA - Insulin Autoantibodies



Type 1 Diabetes Associated with other immune conditions

- ▶ Celiac disease (gluten intolerance)
- ▶ Thyroid disease
- ▶ Addison's Disease
- ▶ Rheumatoid arthritis
- ▶ Other



Type 1 Summary

- ▶ Autoimmune pancreatic destruction
- ▶ Need insulin replacement therapy
- ▶ Often first present in DKA
- ▶ At risk for other autoimmune diseases
- ▶ Eval coping strategies



Diabetes Education
SERVICES



Patti LaBelle
"divabetic"
"I have diabetes, it
doesn't have me"

"I don't want diabetes
to steal one more life."
- Patti LaBelle


Join Patti LaBelle
to Stop
Diabetes®

Donate now and give hope





Diabetes Education
SERVICES

BMI Categories



Diabetes Education
SERVICES





Cardio Metabolic Risk - 5 Hypers -

- ▶ Hyperinsulinemia (resistance)
- ▶ Hyperglycemia
- ▶ Hyperlipidemia
- ▶ Hypertension
- ▶ Hyper"waistline"emia (35" women, 40" men)



Manifestations of Insulin Resistance

2. Classification and DM Diagnosis

- ▶ Pre Diabetes & Type 2- Screening Guidelines
- ▶ Start screening **at age 45 or for anyone** who is **overweight** (BMI \geq 25, Asians BMI \geq 23) with one or > additional **risk factor**:
 - ▶ First-degree relative w/ diabetes
 - ▶ Member of a high-risk ethnic population
 - ▶ Habitual physical inactivity
 - ▶ PreDiabetes
 - ▶ History of heart disease



Diabetes Education SERVICES

Diabetes 2 - Who is at Risk?

(ADA Clinical Practice Guidelines)

Risk factors cont'd



- ▶ HTN - BP > 140/90
- ▶ HDL < 35 or triglycerides > 250
- ▶ history of Gestational Diabetes
- ▶ Polycystic ovary syndrome (PCOS)
- ▶ Other conditions assoc w/ insulin resistance:
 - ▶ Severe obesity, acanthosis nigricans (AN)
 - ▶ Recheck every 3 years



Diabetes Education SERVICES

Acanthosis Nigricans (AN)

- ▶ Signals high insulin levels in bloodstream
- ▶ Patches of darkened skin over parts of body that bend or rub against each other
 - ▶ Neck, underarm, waistline, groin, knuckles, elbows, toes
 - ▶ Skin tags on neck and darkened areas around eyes, nose and cheeks.
- ▶ No cure, lesions regress with treatment of insulin resistance



Diabetes Education SERVICES

Diabetes Detectives Needed




- ▶ On average – takes 6.5 years to diagnose diabetes
- ▶ 1/4 of all people with diabetes don't know they have it
- ▶ 50% of Latino and Asians are undiagnosed



Diabetes Education SERVICES

Ominous Octet



SGLT2 Inhibitors- "Glucolectics"



- ▶ **Action:** "Glucolectic" decreases renal glucose reabsorption (resets renal threshold and increases glucosuria)
- ▶ Expensive

Class/Main Action	Name(s)	Daily Dose Range	Considerations
SGLT2 Inhibitors "Glucolectic"	Canagliflozin (Invokana)	100 - 300 mg 1x daily Don't start if GFR <45.	Side effects: hypotension, UTIs, increased urination, genital infections, ketoacidosis. Monitor GFR and other considerations: See package insert for dosing adjustment based on GFR.
• Decreases glucose reabsorption in kidneys	Dapagliflozin (Farxiga)	5 - 10 mg 1x daily Don't start if GFR <60.	- Canagliflozin increases risk of amputation. - Dapagliflozin, don't use in pts w/ bladder cancer.
	Empagliflozin (Jardiance)	10 - 25 mg 1x daily Don't start if GFR <45.	- Empagliflozin & canagliflozin decrease risk of death from CV disease.
	Ertugliflozin (Steglatro)	5 - 15 mg 1x daily Don't start if GFR <60.	Benefits: no hypo or weight gain. Lowers A1c 1.0%-2.0%. Lowers wt 1-3 lbs.



Diabetes Education SERVICES

Jardiance gets special FDA CV approval



Diabetes Education Services

Published by Beverly Thomassian (P) · December 2, 2016 ·

Jardiance decreases CV Mortality by 38%. The (FDA) has approved empagliflozin (Jardiance) for the new indication of improving survival in adults with type 2 diabetes and cardiovascular disease (CVD). Important info to share!



FDA Approves Empagliflozin for Reducing CVD Death

The new indication follows the landmark EMPA-REG trial, the first to show that a diabetes drug could reduce death as well as lower blood glucose.



Diabetes Education SERVICES

Comparison of Type 1 and Type 2

Feature	Type 1	Type 2
▶ Obesity	x	xxx
▶ Insulin dependence	xxx	30%
▶ Respond to oral agents	x	xxx
▶ Antibodies present	xxx	0
▶ Typical age of onset	puberty	40-65
▶ Insulin Resistance	x	xxx



Diabetes Education SERVICES

Gestational DM ~ 7% of all Pregnancies

- ▶ GDM prevalence increased by
 - ▶ ~10-100% during the past 20 yrs
- ▶ Native Americans, Asians, Hispanics, African-American women at highest risk
- ▶ Immediately after pregnancy, 5% to 10% of GDM diagnosed with type 2 diabetes
- ▶ Within 5 years, 50% chance of developing DM in next 5 years.



Diabetes Education SERVICES

Postnatal Health: Maternal Behavior

- ▶ Encourage breastfeeding for at least 6 mos
 - ▶ (Decreases risk of maternal diabetes 48%)
- ▶ Screening 6-12 weeks post partum using non-pregnant OGTT criteria (50%)
- ▶ Repeat at 3 yr intervals or signs of DM
- ▶ Encourage weight control and exercise
- ▶ Make sure connected with health care
- ▶ Preconception counseling



Diabetes Education SERVICES

Start Metformin therapy

- ▶ For women with PreDiabetes and History of GDM



Diabetes Education SERVICES

Metformin – New GFR Guidelines

Class/Main Action	Name(s)	Daily Dose Range	Considerations
Biguanides • Decreases hepatic glucose output • First line med at diagnosis of type 2	metformin (Glucophage)	500 - 2500 mg (usually BID w/ meal)	Side effects: nausea, bloating, diarrhea, B12 deficiency. To minimize GI Side effects, use XR and take w/ meals. Obtain GFR before starting. <ul style="list-style-type: none"> • If GFR <30, do not use. • If GFR <45, don't start Metformin • If pt on Metformin and GFR falls to 30-45, eval risk vs. benefit; consider decreasing dose. For dye study, if GFR <60, liver disease, alcoholism or heart failure, restart metformin after 48 hours if renal function stable. Benefits: lowers cholesterol, no hypo or weight gain, cheap. Approved for pediatrics, 10 yrs + Lowers A1c 1.0%-2.0%.
	Riomet (liquid metformin)	500 - 2500mg 500mg/5mL	
	Extended Release-XR (Glucophage XR) (Glumetza) (Fortamet)	(1x daily w/dinner) 500 - 2000 mg 500 - 2000 mg 500 - 2500 mg	



Biguanide derived from:
Goat's Rue *Galega officinalis*,
French Lilac



Diabetes Education SERVICES

Biguanides – Metformin (Glucophage)

- ▶ **Action:** decrease hepatic glucose (glycogen)
- ▶ **Names:**
 - ▶ Metformin (Glucophage)
 - ▶ Starting dose: 500 BID, max 2500mg daily
 - ▶ Metformin extended release (3 different versions)
 - ▶ Starting dose 500mg at dinner, max dose 2000 to 2500 mg daily
- ▶ **Efficacy:**
 - ▶ Decrease fasting plasma glucose 60-70 mg/dl
 - ▶ Reduce A1C 1.0-2.0%



Biguanides - Metformin

- ▶ **Benefits**
 - ▶ Decrease LDL cholesterol and triglycerides
 - ▶ No weight gain, possible modest weight loss
 - ▶ Cancer protective?
- ▶ **Concerns**
 - ▶ Diarrhea and abdominal discomfort – Use XR (may see pill shell in stool – okay)
 - ▶ Lactic acidosis if improperly prescribed
 - ▶ Watch for B12 deficiency
 - ▶ Special considerations for IV contrast dye studies. Resume when kidney function adequate.



If on Metformin and Sulfonylurea – A1c 8.4 - Pt struggling with weight



Other Causes of Hyperglycemia

- ▶ Steroids
- ▶ Agent Orange
- ▶ Tube feedings / TPN
- ▶ Transplant medications
- ▶ Cystic Fibrosis

Regardless of cause, requires treatment

- ▶ Insulin always works
- ▶ Sign of pancreatic malfunction



Diabetes Education SERVICES

Diabetes is also associated with

- ▶ Fatty liver disease
- ▶ Obstructive sleep apnea
- ▶ Alzheimer's
- ▶ Depression
- ▶ Cancer; pancreas, liver, breast



Diabetes Education SERVICES

DiaBingo

- B** Frequent skin and yeast infections
- B** A BMI of ____ or greater is considered overweight
- B** To reduce complications, control **A1c**, **B**lood pressure, **C**holesterol
- B** PreDiabetes – fasting glucose level of ____ to ____
- B** Erectile dysfunction indicates greater risk for ____
- B** Diabetes – fasting glucose level ____ or greater
- B** Type 1 diabetes is best described as an ____ disease
- B** People with diabetes are ____ times more likely to die of heart dx
- B** Elevated triglycerides, < HDL, smaller dense LDL
- B** Each percentage point of A1c = ____ mg/dl glucose
- B** At dx of type 2, about ____% of the beta cell function is lost
- B** Diabetes – random glucose ____ or greater



Diabetes Education SERVICES

Life Study – Mrs. Jones

Mrs. Jones is 62 years old, overweight and complaining of feeling tired and urinating several times a night. She is admitted with a urinary tract Infection. Her WBC is 12.3, glucose 237. She is hypertensive with a history of gestational diabetes. No ketones in urine.

- ▶ What are her risk factors, signs of diabetes
- ▶ What type of diabetes does she have?
- ▶ Does she have insulin resistance?



Strategies – One Step at a Time Everyone can be a Diabetes Advocate



Look for
“teaching moment”
opportunities



Diabetes Education
SERVICES

What Do You Say? Mrs. Jones asks you

- ▶ What is type 2 diabetes?
- ▶ Will this go away?
- ▶ Will I get complications?
- ▶ Will I need to take diabetes medication for the rest of my life?
- ▶ How come I got diabetes?
- ▶ Do I have to check my blood sugars?



Diabetes Education
SERVICES

Patient Centered Approach

- ▶ Support informed decision making
- ▶ Problem solving
- ▶ Active collaboration to improve clinical outcomes and quality of life
- ▶ Avoid judgmental words that increase feelings of shame and/or guilt
- ▶ Choose words and phrases that put people first
- ▶ Avoid shame and blame



Diabetes Education SERVICES

Overcoming barriers

- ▶ Focus on pt's strengths
- ▶ Offer pts evidence based hope message
- ▶ Frequent contact
- ▶ Paired glucose testing
- ▶ Ask pt, "Tell me 1 thing that is driving you crazy about your diabetes"
- ▶ Discuss medication beliefs

Bill Polonsky, PhD, CDE



Diabetes Education SERVICES

How will blood glucose testing help me?

- ▶ See if your treatment plan is working
- ▶ Make decisions regarding food and/or med adjustment when exercising
- ▶ Find out how that pizza affected your BG
- ▶ Avoid unwanted weight gain
- ▶ Enhanced athletic performance
- ▶ Find patterns
- ▶ Manage illness



Diabetes Education SERVICES

How Often Should I Check?

- ▶ Be realistic!!
- ▶ Type 2 on orals – Medicare covers 100 strips for 3 months
- ▶ Based on individual - Consider:
 - ▶ Types and timing of meds
 - ▶ Goals
 - ▶ Ability (physical and emotional)
 - ▶ Finances / Insurance



Diabetes Education SERVICES

Complications - Why?



- ▶ Degree of hyperglycemia “glucose toxicity”
- ▶ Duration of hyperglycemia
- ▶ Genes
- ▶ Multiple risk factors: smoking, vascular disease, dyslipidemia, hypertension, other



Diabetes Education SERVICES

Diabetes Complications

- ▶ Heart disease leading cause of death.
- ▶ CAD death rates are about 2 -4x's as high as adults without diabetes (it's not getting better)
- ▶ Risk of stroke is 2 - 4 times higher
- ▶ 60% - 65% of people with DM have HTN.
- ▶ DM accounts for 40% of new cases of ESRD
- ▶ 60 - 70% have mild - severe forms of neuropathy
- ▶ Diabetes is the leading cause of blindness
- ▶ Accounts for 50% of lower limb amputations



Diabetes Education SERVICES

Control Matters

- ▶ Prevention
- ▶ Trials
- ▶ Practice Recommendations



Diabetes Education SERVICES

Financial Advisor

- ▶ Mid 30s, friendly, he smiles to greet you and you notice his gums are inflamed. You'd guess a BMI of 26 or so, with most of the extra weight in the waist area.
- ▶ If you could give him some health related suggestions, what would they be?



Diabetes Education SERVICES

Can we stop pre diabetes from progressing?

3, 234 people w/ Pre-Diabetes randomized:

- ▶ Placebo
- ▶ Diet/Exercise or
- ▶ Metformin

over a three year period



Diabetes Prevention Program (DPP) 2001



Diabetes Education SERVICES

Diabetes Prevention Program

- ▶ Standard Group - 29% developed DM
- ▶ Lifestyle Results - 14% developed DM
 - ▶ 58% (71% for 60yrs +) Risk reduction
 - ▶ 30 mins daily activity
 - ▶ 5-7% of body wt loss
- ▶ Metformin 850 BID - 22% developed DM
 - ▶ 31% risk reduction (less effective with elderly and thinner pt's)




Diabetes Education SERVICES

Weight loss and Prevention

- ▶ For every 2.2 pounds of weight loss, risk of type 2 diabetes was reduced by 13%.



Diabetes Education SERVICES



Goals of Care



Diabetes Education
SERVICES

ABCs of Diabetes –


- ▶ **A**1c less than 7% (avg 3 month BG)
 - ▶ Pre-meal BG 80-130
 - ▶ Post meal BG <180
- ▶ **B**lood Pressure < 140/90
- ▶ **C**holesterol
 - ▶ DM and 40 yrs, start statin
 - ▶ HDL >40
 - ▶ Triglyceride < 150
- ▶ **E**xercise, Education
- ▶ **H**ealthy Eating



Diabetes Education
SERVICES

6. Glycemic Targets

- ▶ **Adult non pregnant A1c goals**
 - ▶ **A1c < 7%** - a reasonable goal for adults.
 - ▶ **A1c < 6.5%** - may be appropriate for those without significant risk of hypoglycemia or other adverse effects of treatment.
 - ▶ **A1c < 8%** - may be appropriate for patients with history of hypoglycemia, limited life expectancy, or those with longstanding diabetes and vascular complications.



Diabetes Education
SERVICES

A1c and Estimated Avg Glucose (eAG) 2008

A1c (%)	eAG
5	97
6	126
7	154
8	183
9	212
10	240
11	269
12	298

Order
teaching tool
kit free at
diabetes.org



eAG = 28.7 x A1c - 46.7 ~ 29 pts per 1%

Translating the A1c Assay Into Estimated Average Glucose Values – ADAG Study
Diabetes Care: 31, #8, August 2008



Diabetes Education
SERVICES

What are next steps?

- ▶ 72 yr old, thin, lives alone, A1c 7.3%. History of MI, stroke. DM for 12 yrs, “diet controlled”. Good Insurance. Creat 1.4.



Diabetes Education
SERVICES

DPP-4 Inhibitors – “Incretin Enhancers”

Januvia (sitagliptin) – Tradjenta (linagliptin)
Onglyza (saxagliptin) Nesina (alogliptin)

- ▶ **Action:**
 - ▶ Increase insulin release w/ meals
 - ▶ Suppress glucagon
- ▶ **Dosing:** Januvia – 100mg a day
Onglyza* – up to 5mg a day
Tradjenta – 5mg a day
Nesina* – up to 25 mg a day
- ▶ **Efficacy:** Decreases A1c by 0.6 -0.8%
- ▶ **Benefits/ Issues:** weight neutral, no hypo, few side effects. Expensive



Diabetes Education
SERVICES

DPP-IV Inhibitor Updates

- ▶ Can cause severe, disabling joint pain.
 - ▶ Contact Provider, Stop Medication
- ▶ Saxagliptin (Onglyza) and Alogliptin (Nesina) can increase risk of heart failure.
 - ▶ Notify provider for shortness of breath, edema, weakness, etc.
- ▶ Side effects: headache and flu-like symptoms
- ▶ Report signs of pancreatitis
- ▶ No wt gain or hypoglycemia
- ▶ Lowers A1c 0.6% - 0.8%



Diabetes Education SERVICES

Exercise Standards

- ▶ Adults – 150 min/wk moderate intensity
 - ▶ over 3 days a week.
 - ▶ Don't miss > 2 consecutive days w/out exercise
 - ▶ Get up every 30 mins - Reduce sedentary time
 - ▶ Flexibility and balance training 2-3 xs a week (Yoga and Tai Chi)
 - ▶ T1 and T2 – resistance training 2 -3 xs a week



Diabetes Education SERVICES

A hard truth


- ▶ Exercise alone doesn't cause weight loss
- ▶ But....
 - ▶ It helps keep weight off
 - ▶ Decreases visceral fat
 - ▶ Decreases CV Risk
- ▶ To combat obesity, we need to change the food environment
- ▶ "You cannot outrun a bad diet"

IT TAKES 524 BURPEES
TO BURN OFF 1 LARGE FRIES
BURPEES SUCK. SO CHOOSE WISELY.
@BIGHEALTH



Diabetes Education SERVICES

Where are your patients on this continuum?



Diabetes Education SERVICES

Good Exercise Info / Quotes

▶ “Passagiata” – take an after meal stroll

- ▶ Exercise decreases A1c 0.7%
- ▶ No change in body wt, but 48% loss in visceral fat
 - ▶ ADA PostGrad 2010



“Every minute of activity lowers blood sugar one point.”

“I don’t have time to exercise, I MAKE time.” Mike Huckabee

Each minute of activity lowers BG by 1 point

- ▶ Each minute of exercise lowers BG 1 point
 - ▶ 15 minutes of walking drops BG 15 points
 - ▶ 30 minutes of biking drops BG 30 points
 - ▶ 40 minutes of housework drops BG 40 points
 - ▶ 50 minutes of walking and window shopping lowers BG 50 points



Diabetes Education SERVICES

DiaBingo- G

- G ADA goal for A1c is less than ____%
- G People with DM need to see their provider at least every month
- G Blood pressure goal is less than
- G People with DM should see eye doctor (ophthalmologist) at least
- G The goal for triglyceride level is less than
- G Goal for my HDL cholesterol is more than
- G The goal for blood sugars 1-2 hours after a meal is less than:
- G People with DM should get this shot every year
- G People with DM need to get urine tested yearly for _____
- G Periodontal disease indicates increased risk for heart disease
- G The goal for blood sugar levels before meals is:
- G The activity goal is to do ___ minutes on most days



Diabetes Education SERVICES

Diabetes Care Guidelines- ADA

Test / Exam	Frequency
▶ A1c	At least twice a year
▶ B/P	Each diabetes visit
▶ Cholesterol (LDL, HDL, Tri)	Yearly (less if normal)
▶ Vaccinations	Flu yearly, pneumonia
▶ Weight	each diabetes visit
▶ Microalbumin/GFR/Creat	Yearly
● Eye exam	Yearly
● Dental Care	At least twice a year
● Comprehensive Foot Exam	Yearly (more if high risk)
● Physical Activity Plan	As needed to meet goals
● Preconception counseling	As needed



Diabetes Education SERVICES

Comprehensive Diabetes Screening HEDIS

Monitor	Numerator	Denominator
Comprehensive Diabetes screening – All Four Tests A1c, lipid profile, dilated eye exam, nephropathy monitor	Number of people who received at least one of each of the following tests: A1c test, cholesterol screening test, diabetes eye exam, and Medicaid attention for nephropathy	Number of people ages 18 to 75 with diabetes
Comprehensive Diabetes Care: Hemoglobin A1c (HbA1c) Poor Control (>9.0%)	Number of people whose most recent HbA1c level indicated poor control (>9.0 percent), was missing or did not have a HbA1c test	Number of people ages 18 to 75 with diabetes



Diabetes Education SERVICES

Mr. Jones - What are Your Recommendations?

Patient Profile

64 yr old with type 2 for 11 yrs. Hx of CVD.

Labs:

- ▶ A1c 9.3%
- ▶ HDL 37 mg/dl
- ▶ Triglyceride 260mg/dl
- ▶ Proteinuria - neg
- ▶ B/P 152/94

Self-Care Skills

- ▶ Walks dog around block 3 x's a week
- ▶ Bowls every Friday
- ▶ 3 beers daily
- ▶ *What meds?*
- ▶ *What referrals?*
- ▶ *My foot hurts*



Diabetes Education SERVICES

Using Alcohol Safely

- ▶ Women- 1 or fewer alcoholic drinks a day
- ▶ Men 2 or fewer alcoholic drinks a day
 - ▶ 1 alcoholic drink equals
 - ▶ 12 oz beer, 5 oz glass of wine, or 1.5 oz distilled spirits (vodka, gin etc)
- ▶ If drink, limit amount and drink w/ food.
- ▶ Ask HCP if safe for you to drink. Tell them your usual quantity and frequency.
- ▶ Can cause hypo and worsen neuropathy



Diabetes Education SERVICES

Hospitalization Prevention

Monitor	Domain	Numerator	Denominator
PQI # 1 (DM Short term complications) *AHRQ 4.4	ED & Hospitalization	Number of admissions with a principal diagnosis of diabetes with short-term complications (ketoacidosis, hyperosmolarity, or coma)	Number of people 18 years and older as of June 30 of measurement year

*Agency for Healthcare Research and Quality (AHRQ)


ACO - 36 | All-Cause Unplanned Admissions for Patients with Diabetes

Medicare's vision is that these measures will illuminate variation in hospital admission rates and incentivize Accountable Care Organizations to develop efficient and coordinated chronic disease management strategies that anticipate and respond to patients' needs and preferences.



Diabetes Education SERVICES

Foot Wounds




Diabetes Education SERVICES

No Bathroom Surgery



Diabetes Education SERVICES

5.07 monofilament = 10gms linear pressure



<http://www.hrsa.gov/leap/>



Diabetes Education SERVICES

Mr. Jones - What are Your Recommendations?

Patient Profile

64 yr old with type 2 for 11 yrs. Hx of CVD.



Current Status:

- ▶ A1c 9.3%
- ▶ On Metformin 500mg BID
- ▶ Partial foot amputation
- ▶ Lives alone
- ▶ What resources, teaching?



Diabetes Education SERVICES

Three Most Important Foot Care Tips

- ▶ Inspect and apply lotion to your feet every night before you go to bed.
- ▶ Do NOT go barefoot, even in your house. Always wear shoes!
- ▶ Every time you see your doctor, take off your shoes and show your feet.



Diabetes Education SERVICES

Psychological Insulin Resistance (PIR)

- ▶ 50% of providers in study threatened pts "with the needle".
- ▶ Less than 50% of providers realized insulins' positive effect on type 2 dm
- ▶ Most pts don't believe that insulin would "better help them manage their diabetes".
- ▶ Solutions: Find the root of PIR and address




Diabetes Attitudes, Wishes, Needs Study - Rubin



Diabetes Education SERVICES

Needle Size often a Barrier Size *Does* Matter




- ▶ Use more short needles – 4 mm
- ▶ Effective for pts with BMI of 24- 49
- ▶ Keeps it subq
- ▶ If pt thin, inject at angle
- ▶ To avoid leakage, count to 10 before withdrawing needle
- ▶ ½ the patients who could benefit from insulin are not using it due to needle phobias



Diabetes Education
SERVICES

Physiologic Insulin Secretion: 24-Hour Profile



Diabetes Education
SERVICES

Insulin Action Teams

- ▶ Bolus: lowers after meal glucose levels
 - ▶ Very Rapid Acting – Aspart (Fiasp)
 - ▶ Rapid Acting
 - ▶ Aspart, Lispro, Admelog, Glulisine, Afrezza
 - ▶ Short Acting - Regular
- ▶ Basal: controls glucose between meals, hs
 - ▶ Intermediate
 - ▶ NPH
 - ▶ Long Acting
 - ▶ Detemir (Levemir)
 - ▶ Glargine (Lantus, Basaglar)
 - ▶ Degludec (Tresiba)



Diabetes Education
SERVICES

Case Study



- ▶ 70 yr old, weighs 100kg
- ▶ History of CABG, tobacco
- ▶ A1c – 11.3%, BG 400-500 for past weeks
- ▶ Insulin – 100+ units Lantus at hs (solostar)
- ▶ Oral Meds: Metformin, Invokana
- ▶ What is a better insulin dosing strategy?
- ▶ Pt can't afford insulin pen – what other option
- ▶ [Diabetes Meds on a Budget - 2014](#) - provides practical and affordable strategies to manage hyperglycemia

Cost Per Vial in Northern CA

Per vial cost	Walmart	Walgreens	Costco
Regular Insulin	\$25*	\$92	\$99
NPH	\$25*	\$92	\$99
70/30	\$25*	\$92	\$101
Humalog	\$200	\$220	\$178
Novolog	\$197	\$217	\$178
Apidra	\$180	\$246	\$178
Levemir	\$300	\$300	\$300
Lantus	\$226	\$221	\$206

Bolus Insulins (½ of total daily dose ÷ meals)

Name	Onset	Peak Action
▶ Aspart (Fiasp)	2.5 min	1 hour
▶ Aspart (NovoLog)	15-30 min	1-1.5 hrs
▶ Lispro (Humalog, Admelog)		
▶ Glulisine (Apidra)		
▶ Afrezza (Inhaled)		
▶ Regular	30 mins	2-4 hrs

Aspart (Fiasp)


- ▶ New Aspart formulation, which includes the addition of niacinamide (vitamin B3) to increase absorption speed
- ▶ Appears in blood in ~ 2.5 mins.
- ▶ Faster onset *and* offset.
- ▶ Starts working within minutes
- ▶ Can be taken as long as 20 minutes after starting a meal.
- ▶ Fiasp available in Flex Touch Pens and 10mL vials.



Diabetes Education SERVICES

Lispro (Admelog) - bolus Glargine (Basaglar) – basal

- ▶ Can't use the term generics for *large* molecule biologicals because they are manufactured in living organisms (bacteria and yeast)
- ▶ Each batch may be slightly different
- ▶ Correct term is "biosimilar"
- ▶ Currently - Pharmacist to contact Provider before switching to biosimilar
 - ▶ Future – may be same as generics



Insulin – Large Molecule



Aspirin – Small Molecule



Diabetes Education SERVICES

Bolus Insulin Summary

- ▶ Regular, aspart, lispro, glulisine,
- ▶ Starts working fast (15-30 mins)
- ▶ Gets out fast (3-6 hours)
- ▶ Post meal BG reflects effectiveness
- ▶ Should comprise about ½ total daily dose
- ▶ Covers food or hyperglycemia.
- ▶ 1 unit
 - ▶ Covers ≈ 10 -15 gms of carb
 - ▶ Lowers BG ≈ 30 – 50 points



Diabetes Education SERVICES

Bolus Insulin Timing

- ▶ How is the effectiveness of bolus insulin determined?
 - ▶ 2 hour post meal (if you can get it)
 - ▶ Before next meal blood glucose
- ▶ Glucose goals (ADA) – may be modified by provider/pt
 - ▶ 1-2 hours post meal <180
 - ▶ Before next meal – 80 - 130



Pattern Management –AKA

How to think like a pancreas



Pattern Management

- ▶ Safety 1st!! - Evaluate 3 day patterns
- ▶ **Hypo:** eval 1st and fix:
 - ▶ If possible, decrease medication dose
 - ▶ Timing of meals, exercise, medications
- ▶ **Hyperglycemia:** evaluate 2nd
 - ▶ Identify patterns
 - ▶ Before increase insulin, make sure not missing something (carbs, exercise, omission)



Bolus – Insulin Sliding Scale

Starts at 150, 2 units for every 50 mg/dl >150

	Break	Lunch	Dinner	HS
Day 1	94 no insulin	212 4 uR	148 no insulin	254 6 uR
Day 2	243 4uR	254 6 uR	201 4uR	199 no insulin
Day 3	189 2uR	243 4uR	162 2uR	244 4uR
Day 4	66 No insulin	287 6uR	144 none	272 6uR



Diabetes Education SERVICES

Basal Insulins

(½ of total daily dose)

Intermediate Acting	Peak Action	Duration
▶ NPH	4-12 hrs	12-24

Long Acting	Peak Action	Duration
▶ Detemir (Levemir)	No Peak	20 hrs
▶ Glargine (Lantus)		24 hrs
▶ Glargine (Basaglar)		24 hrs
▶ Degludec (Tresiba)		42 hrs

Fasting BG reflects efficacy of basal



Diabetes Education SERVICES

Degludec

▶ Degludec (Tresiba)

- ▶ An ultra long acting insulin - lasts up to 42 hours
- ▶ Takes 3-4 days to reach steady state
- ▶ Available in u-100 and u-200 pens
- ▶ Seems to cause less hypo
- ▶ Adjust dose every 3-4 days
- ▶ Wait at least 8 hours between doses
- ▶ Good at room temp for 8 wks



▶ Ryzodeg 70/30

- ▶ mixture of insulin degludec and aspart



Diabetes Education SERVICES

Basal Insulin Summary

- ▶ NPH, Levemir, Lantus, Degludec
- ▶ Covers in between meals, through night
- ▶ Starts working slow (4 hours)
- ▶ Stays in long (12-24 hours)
 - ▶ NPH 12 hrs
 - ▶ Levemir, Lantus 20-24 hrs
 - ▶ Degludec – 42 hours
- ▶ Fasting blood glucose reflects effectiveness



Type 2 Metformin 2000 mg day, Lantus 40 units hs. Weighs 80kg. A1c 8.7%

▶ Blood Sugars

	AM	Lunch	Dinner	HS
Day 1	137	178	203	193
Day 2	96	154	267	242
Day 3	73	127	253	269
Day 4	61	193	333	282
Day 5?				



Next Steps

- ▶ When is it too much basal insulin?
 - ▶ If basal insulin is >0.5 units/kg day, advance to combination injectable therapy
 - ▶ Add bolus, switch premixed 70/30 or to Basal + GLP-RA
- ▶ Pt is at max basal dose
 - ▶ $80 \times 0.5 = 40$ units



24u 70/30 am, 16 u 70/30 pm
 Patterns? Changes needed?

	Break	Lunch	Dinner	HS
Day 1	102	63	92	181
Day 2	112	67	106	195
Day 3	98	56	112	201
Day 4	99	71	132	211



Diabetes Education SERVICES

What Medications Cause Hypoglycemia?

- ▶ Insulin
- ▶ Sulfonylureas
- ▶ Meglitinides
- ▶ Or any combo medication that includes these



Diabetes Education SERVICES

Sulfonylureas - Squirts

- ▶ Action: Increase endogenous insulin secretion throughout day
- ▶ Efficacy:
 - ▶ Decrease FPG 60-70 mg/dl
 - ▶ Reduce A1C by 1.0-2.0%
- ▶ Side Effects:
 - ▶ Weight gain, hypoglycemia
- ▶ Benefits:
 - ▶ Cheap, effective



Diabetes Education SERVICES

Hypoglycemia = "Limiting Factor"

- ▶ Defined as glucose of 70mg/dl or below
- ▶ 50% of episodes occur during the night
- ▶ Higher mortality rate with severe hypoglycemia secondary to sulfonylureas
 - ▶ Especially (glyburide) Micronase®, Diabeta®
- ▶ Blood glucose levels don't describe severity, response is individual



Diabetes Education SERVICES

Hypoglycemic Symptoms

- ▶ Autonomic
 - ▶ Anxiety
 - ▶ Palpitations
 - ▶ Sweating
 - ▶ Tingling
 - ▶ Trembling
 - ▶ Hypoglycemic Unawareness
- ▶ Neuroglycopenia
 - ▶ Irritability
 - ▶ Drowsiness
 - ▶ Dizziness
 - ▶ Blurred Vision
 - ▶ Difficulty with speech
 - ▶ Confusion
 - ▶ Feeling faint



Diabetes Education SERVICES

Treatment of Hypoglycemia

- ▶ If blood glucose **70mg/dl** or below:
 - 10-15 gms of carb to raise BG 30 - 45mg/dl
- Ⓞ Retest in 15 minutes, if still low, treat again, even without symptoms
- Ⓞ Follow with usual meal or snack
- Ⓞ If non responsive, give D50 IV or glucagon Emergency Kit
- Ⓞ Figure out how to prevent in future



Diabetes Education SERVICES

15 - 20 Gms Carb Sources

- ⦿ 4 ounces apple juice
- ⦿ 3 - 4 Glucose Tablets
- ⦿ 8 - 10 Lifesavers candy
- ⦿ 8 - 10 Hard candies
- ⦿ 2 Tablespoons Raisins
- ⦿ 4 - 6 oz's Nondiet soda
- ⦿ 4 - 6 oz's Fruit Juice
- ⦿ 8 oz Milk (non fat)



Diabetes Education SERVICES



Diabetes Education SERVICES

Basal Bolus – What Adjustments?

Pt weighs 80kg

	Break	Lunch	Dinner	HS
Day 1	69 7H	79 5H	245 8H	190 22u Det
Day 2	81 7H	87 5H	170 8H	133 22u Det
Day 3	73 7H	94 5H	194 8H	110 22u Det
Day 4	62 7H	83 5H	211 8H	127 22u Det



Diabetes Education SERVICES

**Intensive Diabetes Therapy
Insulin Dosing Strategy**

50/50 Rule

- ▶ 0.5-1.0 units/kg day
- ▶ Basal = 50% of total
 - Glargine QD
 - NPH or Detemir BID
- Bolus = 50% of total
 - usually divided into 3 meals

Example

- ▶ Wt 50kg x 0.5 = 25 units of insulin/day
- ▶ Basal dose: 13 units
 - Glargine 13 units QD
 - NPH/Detemir 6u BID
- ▶ Bolus dose: 12 units
 - ▶ 4 units NovoLog, Apidra Humalog, Regular each meal



Diabetes Education SERVICES

**Intensive Diabetes Therapy
Insulin Dosing Strategy**

50/50 Rule

- ▶ 0.5-1.0 units/kg day
- ▶ Basal = 50% of total
 - Glargine QD
 - NPH or Detemir BID
- Bolus = 50% of total
 - usually divided into 3 meals

Example – You Try

- ▶ Wt 60 kg x 0.5 = ____ units of insulin/day
- ▶ Basal dose: ____ units
 - Glargine ____ QD
 - NPH/Detemir __ BID
- ▶ Bolus dose: ____ units
 - ____units NovoLog, Apidra Humalog, Reg each meal



Diabetes Education SERVICES

**Intensive Diabetes Therapy
Insulin Dosing Strategy**

50/50 Rule

- ▶ 0.5-1.0 units/kg day
- ▶ Basal = 50% of total
 - Glargine QD
 - NPH or Detemir BID
- Bolus = 50% of total
 - usually divided into 3 meals

Example – You Try

- ▶ Wt 60kg x 0.5 = 30 units of insulin/day
- ▶ Basal dose: 15 units
 - Glargine 15 QD or
 - NPH/Detemir 7u BID
- ▶ Bolus dose: 15 units
 - ▶ 5 NovoLog, Apidra, Humalog, Reg each meal



Diabetes Education SERVICES

Insulin Teaching Keys

- ▶ Bolus insulin with meals
- ▶ Basal 1-2xs daily
- ▶ Abdomen preferred injection site
- ▶ Stay 1" away from previous site
- ▶ Don't re-use ultra fine syringes
- ▶ Keep unopened insulin in refrigerator
- ▶ Toss opened insulin vial after 28 days
- ▶ Proper disposal
- ▶ Review patients ability to withdraw and inject.
- ▶ Side effects include hypoglycemia/wt gain
- ▶ Insulin pens –
 - ▶ Prime needle to assure accurate insulin dose given
 - ▶ Hold needle in for 5 seconds after injection
 - ▶ Roll 70/30 pens



Diabetes Education SERVICES

Sharps Disposal: Product and Info

- ▶ Look in the Government section white pages for a household hazardous waste listing for your city or county.
- ▶ Call 1-800-CLEANUP (1-800-253-2687)
- ▶ Search for collection centers on the California Integrated Waste Management Board (CIWMB) Web site:
<http://www.ciwmb.ca.gov/HHW/HealthCare/Collection/>




Diabetes Education SERVICES

Diabetes Vacations

“The greatest glory in living lies not in never falling, but in rising every time we fall.”

—Nelson Mandela



Diabetes Education SERVICES

Your health can only get better



Diabetes Education SERVICES

Bacterial Cells Outnumber Human Cells 10 to 1



Diabetes Education SERVICES

Poll Question

- ▶ How much does your gut bacteria weigh?
 - A. 24 ounces
 - B. 3 pounds
 - C. Less than 1 pound
 - D. 1.5 pounds
 - E. Not sure



Diabetes Education SERVICES

3 lbs of Microbes in our Gut


- ▶ This community of bacteria can be thought of as an extra 'organ' "microbiome".
- ▶ We have evolved together with our microbiome over millions of years.
- ▶ Ratios of these communities has changed over the past 30 years
- ▶ Mirrors global spikes in obesity, diabetes, allergic and inflammatory diseases
- ▶ What are we doing to change these bacteria?



Diabetes Education SERVICES

Gut Microbiome

- ▶ Part of endocrine axis
- ▶ Stabilized by 3 years of age
- ▶ Influenced by:
 - ▶ Birth method
 - ▶ Breast fed
 - ▶ Early Antibiotic use
 - ▶ Environment
 - ▶ Travel
- ▶ Help us
 - ▶ utilize energy
 - ▶ fight off invaders



Diabetes Education SERVICES

C-Section – Consider Gauze in Vagina

- ▶ Eat a healthy, balanced diet during pregnancy.
- ▶ Avoid unnecessary antibiotics.
- ▶ Breast-feed baby for the first 12 months.
- ▶ If possible, avoid a C-section delivery.
- ▶ If you need a C-section, try the "gauze-in-the-vagina technique."
- ▶ early research by Dr. Maria Gloria Dominguez-Bello, an associate professor in the Human Microbiome Program at the NYU School of Medicine. She is testing a fast and easy work-around called the "gauze-in-the-vagina technique."



Diabetes Education SERVICES

Human Intestine Friends

▶ The majority belong 2 major phyla:

▶ Firmicutes

▶ includes *Clostridium*, *Enterococcus*, *Lactobacillus* and *Ruminococcus*

▶ Bacteroidetes

▶ includes *Bacteroides* and *Prevotella*

in proportions determined in part by birth, breastfeeding, diet



Diabetes Education SERVICES

Weight and Gut Bacteria New and Early Research

▶ Leaner people appear to have more bacterial diversity and a higher proportion of **bacteroidetes**

▶ Gut bacteria less efficient at converting food to calories

▶ Obese people appear to have higher levels of **firmicutes**

▶ Gut bacteria very efficient at calorie extraction

▶ Bacteria tend to run in families



Diabetes Education SERVICES

Getting to Better Gut Bacterial Health

Eat more PREbiotics

▶ Foods with indigestible fibers that nourish the good bacteria:

▶ High fiber foods like, whole grains, fruits, veggies, nuts

▶ High in prebiotic fibers include: Jerusalem artichokes, onions, kale, Brussels sprouts, bananas, dandelion greens & more

PRObiotics

▶ These foods contain healthy bacteria like *Bifidobacterium* and *Lactobacillus*.


▶ Yogurt, Kefir – look for “live or active cultures”

▶ Fermented foods like: Sauerkraut, Kimchi, Miso soup, kombucha



Diabetes Education SERVICES

Follow Your Gut – Dr. Rob Knight



Check out Dr. Knight's:

- ▶ TED Talk
- ▶ Website – AmericanFoodProject.org
- ▶ Articles in Nature and all over



Diabetes Education SERVICES

Take Home Message

- ▶ Get Dirty
- ▶ Limit Unnecessary C-Sections
- ▶ Breastfeed if possible
- ▶ Limit early antibiotics
- ▶ Eat a wide variety of fiber foods



Diabetes Education SERVICES

Medical Nutrition Therapy – ADA

- ▶ Focus on the Individual
- ▶ Maintain pleasure of eating
- ▶ Provide positive messages about food
- ▶ Limit food choices only when backed by science
- ▶ Provide practical tools
- ▶ Refer to a RD and Diabetes Education – Lowers A1c by 1-2%



Diabetes Education SERVICES



United States: The Revis family of North Carolina. Food expenditure for one week: \$341.98. Favorite foods: spaghetti, potatoes, sesame chicken. Peter Menzel, from the book, "Hungry Planet: What the World Eats."



Diabetes Education SERVICES



Guatemala: The Mendozas of Todos Santos - Food expenditure for one week: 573 Quetzales or \$75.70. Family Recipe: Turkey... [VIEW MORE](#) Peter Menzel, from the book, "Hungry Planet: What the World Eats."

19 of 27



Diabetes Education SERVICES

Approach Depends on Patient

- New Type 2
 - Portion Control
 - Plate Method
 - Record Keeping
 - Education
- On Insulin?
 - Carb counting
 - Post prandial checks



Diabetes Education SERVICES

Healthy Eating Patterns

- ▶ Mediterranean Diet
- ▶ DASH Diet
- ▶ Plant based eating
- ▶ Diabetes Plate Method
- ▶ Weight Watchers or other groups




Diabetes Education SERVICES

Move toward the Tomato



Diabetes Education SERVICES

Mediterranean Diet Pyramid



Diabetes Education SERVICES

The Mediterranean diet emphasizes:

- ▶ Eating primarily plant-based foods, such as fruits and vegetables, whole grains, legumes and nuts
- ▶ Replacing butter with healthy fats such as olive oil and canola oil
- ▶ Using herbs and spices instead of salt to flavor foods
- ▶ Limiting red meat to no more than a few times a month
- ▶ Eating fish and poultry at least twice a week
- ▶ Enjoying meals with family and friends
- ▶ Drinking red wine in moderation (optional)
- ▶ Getting plenty of exercise



Diabetes Education SERVICES

USDA www.myplate.gov

Balancing Calories

- ▶ Enjoy your food, but eat less.
- ▶ Avoid oversized portions.

Foods to Increase

- ▶ Make half your plate fruits and vegetables.
- ▶ Make at least half your grains whole grains.
- ▶ Switch to fat-free or low-fat (1%) milk.

Foods to Reduce

- ▶ Compare sodium in foods like soup, bread, and frozen meals — and choose the foods with lower numbers.
- Drink water instead of sugary drinks.



Diabetes Education SERVICES

10 SuperFoods

- ▶ Beans
- ▶ Dark Green Leafy Veggies
- ▶ Citrus Fruit
- ▶ Sweet Potatoes
- ▶ Berries
- ▶ Tomatoes
- ▶ Fish High in Omega-3 Fatty Acids
- ▶ Whole Grains
- ▶ Nuts
- ▶ Fat-Free Milk and Yogurt



As posted on diabetes.org website



Diabetes Education SERVICES

Choose Healthy Carbs

- o Carbs have fiber, vitamins, minerals and phytonutrients
- o 25 gms of fiber a day
- o Power Carbs include:
 - o Beans
 - o Veggies
 - o Fruits
 - o Whole grain foods



Diabetes Education SERVICES

Another plate example

Mi planificador de plato Una comida saludable sabe buenísima



El Método del Plato es una manera simple de planificar las comidas para usted y su familia. No necesita contar nada ni usar largas listas de alimentos. Todo lo que necesita es un plato de 9 pulgadas.

3/4 de proteína, 1/4 de almidón, 1/2 de vegetales.

Plato de 9 pulgadas



Diabetes Education SERVICES

Successful weight loss strategies include

- ▶ Weekly self-weighing
- ▶ Eat breakfast
- ▶ Reduce fast food intake.
- ▶ Decrease portion size
- ▶ Increase physical activity
- ▶ Use meal replacements
- ▶ Eat healthy foods
- ▶ Drink Water
- ▶ Sleep



Diabetes Education SERVICES

Diabetes Prevention Program Focus on fat = wt loss success

To help you lose weight and improve your health, stay as close as possible to your fat and calorie goals. Find your starting weight below. Your fat and calorie goals are in the same row. Circle your fat and calorie goals.


Weight (lb)	Fat Goal (grams)	Calorie Goal
120-174	33	1,200
175-219	42	1,500
220-249	50	1,800
>250	55	2,000

<http://www.cdc.gov/diabetes/prevention/recognition/curriculum.htm>



Diabetes Education SERVICES

How nutrients affect blood sugar



Diabetes Education SERVICES

Carbohydrate Needs for Most Adults

	Grams	Servings
Each Meal	45-60 gm	3 - 4
Snacks	15-30 gm	1- 2



Carbs affect Post Meal Blood Glucose



Diabetes Education SERVICES

Ms. Gonzales' Daily Meal plan


Break	Lunch	Dinner	Night
5 corn tortillas, 1/2 c. beans, salsa, peppers, egg beaters	Sandwich, low fat potato chips, 1c. juice, 2-4 lowfat cookies	Lg bowl low salt soup, 1c. rice, BBQ meat, salad & cooked vevs 1 glass wine	1 bowl of cereal
Avg BG 120's	Avg BG 200's	Avg BG 200's	Avg BG 180's



Diabetes Education
SERVICES

Carb Counting - Starch


Each Food has:
80 Calories
15 grams carb



Diabetes Education
SERVICES

Carb counting- fruit

Each Food has:
60 Calories
15 grams carb



Diabetes Education
SERVICES

Carb Counting - Milk

Each Food has:
90-150 calories
12-15 grams carb

1 slice bread

- 8 oz buttermilk
- 1 packet diet hot cocoa
- 8 oz milk
- 8 oz soy milk
- 6 oz plain yogurt
- 6 oz light fruit yogurt

Diabetes Education SERVICES

Carb Counting - Sweets

Each Food has:
Calories vary
15 grams carb

1 slice bread

- 2 inch square cake or brownie, unfrosted
- 1/2 cup diet pudding
- 1/2 cup regular jello
- 2 tsp light syrup
- 2 small cookies
- 1/2 cup ice cream or frozen yogurt
- 1/2 cup sherbet
- 1/4 cup sorbet
- 1 tbsp syrup, jam, jelly, table sugar, honey

Diabetes Education SERVICES

Give the gift of Non-Judgment

out beyond ideas
of wrongdoing
and rightdoing,
there is a field.
i'll meet you there
-rumi

Diabetes Education SERVICES
