

Glucagon Treatment for Diabetes-Related Hypoglycemia



Name / Delivery	Supplied	Dose Range		Age / Route / Storage
		Adults	Peds / Age Wt Dosing	
Glucagon Emergency Kit Injection requires mixing glucagon powder + diluent filled syringe	1mg / 1mL vial + syringe	1 mg	0.03mg/kg or < 6yr or < 25 kgs 0.5mg ≥ 6yr or > 25kgs 1mg	All ages approved SubQ or IM admin Expires in 2 years at room temp.
Gvoke Injectable liquid stable glucagon solution	0.5mg/1.0ml prefilled syringe or 0.5mg/1.0ml HypoPen auto-injector	1 mg	< 2yr: not recommended 2- 12 yrs < 45kg 0.5mg ≥ 45kg 1mg 12 years or older 1mg	Approved Age 2+ SubQ admin in arm, thigh, abdomen Expires in 2 years at room temp (keep in foil pouch).
Baqsimi Nasal glucagon powder	3 mg intranasal device	3 mg	< 4 yrs: not recommended 3 mg dose for 4 years or older	Approved Age 4+ Nasal admin Expires ~ 2 yrs at room temp (keep in shrink-wrapped tube)

**All raise BG 20+ points. Can cause nausea, vomiting. After admin, roll person on side. Seek medical help. If no response after 1st dose, give 2nd dose in 15 mins. When awake, give oral carbs ASAP when safe to swallow. Please consult package insert for detailed info.*

Copyright Diabetes Ed Services 2020.

www.DiabetesEd.net

All PocketCard content is for educational purposes only. Please consult prescribing information for detailed guidelines.

© 2020