



# Insulin Calculation Workshop- For Pumps and Beyond 2022

Beverly Thomassian, RN, MPH, BC-ADM, CDCES  
President, Diabetes Education Services



# DiabetesEd.net Website Orientation

Diabetes Education SERVICES

Advancing Your Career in Diabetes Education

20th Anniversary

Download FREE Medication PocketCards

HOME EXAM PREP ABOUT US CONFERENCES ONLINE COURSES BOOKS + STUDY FREE RESOURCES BLOG BYTES CONTACT US

Start your journey in Diabetes Education

- Online University & Live Seminars
- Certification Tools & Resources
- Accredited Continuing Education

GET STARTED

Coach Beverly

PASS THE CDCES EXAM! (FORMERLY CDE) — 10 STEPS TO SUCCEED

We are here to help!

[www.DiabetesEd.net](http://www.DiabetesEd.net) | [info@diabetesed.net](mailto:info@diabetesed.net) | 530-893-8635

## PocketCards



NEW Accordion 2-sided PocketCards

### Glucagon Treatment for Diabetes-Related Hypoglycemia

Name / Delivery	Supplied	Dose Range	
		Adults	Peds / Age Wt Dosing
Glucagon Emergency Kit Injection requires mixing glucagon powder + diluent filled syringe	1mg / 1mL vial + syringe	1 mg	0.03mg/kg or < 6yr or < 25 kg   0.5mg ≥ 6yr or > 25kg   1mg
Gvoke Injectable liquid stable glucagon solution	0.5mg/1.0mg prefilled syringe or 0.5mg/1.0mg HypoPen auto-injector	1 mg	< 2yr: not recommended 2-12 yrs < 45kg   0.5mg ≥ 45kg   1mg 12 years or older   1mg
Baqsimi Nasal glucagon powder	3 mg intranasal device	3 mg	< 4 yrs: not recommended 3 mg dose for 4 years or older

\*All raise BG 20+ points. Can cause nausea, vomiting. After admin, roll person on side. Seek medical help. If no response after 1st dose, give 2nd dose in 15 mins. When awake, give oral carbs ASAP when safe to swallow. Please consult package insert for detailed info.

All PocketCard content is for educational purposes only. Please consult prescribing information for detailed guidelines.

www.DiabetesEd.net Copyright Diabetes Ed Services 2020. © 2020

Free on CDCES Coach App  
Constantly updated  
App users receive priority notification  
Download on Website



Diabetes Education SERVICES

# Bryanna is here to Help!



Bryanna Sabourin, Director of  
Operations and Customer Happiness

If you have questions,  
you can chat with  
Bryanna at  
[www.DiabetesEd.net](http://www.DiabetesEd.net)  
or call 530 / 893-8635  
or email at  
[info@diabetesed.net](mailto:info@diabetesed.net)





ONLINE  
UNIVERSITY

## 2022 Live Webinar Updates

Previously recorded & available for on-demand viewing

- [The Impact of Adverse Childhood Experiences on Health](#) | 1.0 CE

All courses air at 11:30 a.m. (PST)

- July 19, 2022 | [3 Steps to Save Feet; Assess, Screen, and Report](#) | 1.0 CEs
- August 9, 2022 | [Solving Glucose Mysteries for Type 1](#) | 1.5 CEs
- August 11, 2022 | [Solving Glucose Mysteries for Type 2](#) | 1.5 CEs
- August 23, 2022 | [Insulin Calculation Workshop – From Pumps & Beyond](#) | 1.5 CEs
- August 25, 2022 | [Basal Bolus Therapy in Hospital](#) | 1.5 CEs
- September 15, 2022 | [Type 2 Diabetes Intensive](#) | 2 CEs
- October 26, 2022 | [Cancer & Diabetes](#) | 1.25 CEs
- November 9, 2022 | [Getting to the Gut \(& Skin\)](#) | 1.0 CEs

Topics included in Level 4 Bundle:



Level 4 | Advanced Level &  
Specialty Topic Courses | 13+  
CEs

\$ 139.00

Quantity

1


Add to Cart



Diabetes Education  
SERVICES

# Conflict of Interest and Resources

- ▶ Coach Bev has no conflict of interest
- ▶ Technology field is rapidly changing
- ▶ Photos in slide set are from Pixabay – not actual clients
- ▶ **Resources**
  - ▶ ADCES Practice Paper- [Continuous Subcutaneous Insulin Infusion \(CSII\) Without and With Sensor Integration 2021](#)
  - ▶ Company web sites – virtual demo
  - ▶ ADCES – DANA **Diabetes Advanced Network Access** [www.diabeteseducator.org](http://www.diabeteseducator.org) Need to be AADE Member to access
  - ▶ [ADA Consumer Guide \(diabetes.org\)](http://diabetes.org)
  - ▶ Pumping Insulin by John Walsh, PA, CDCES – Diabetes Mall
  - ▶ Gary Scheiner, MS, CDCES – Integrated Diabetes Services





# Insulin Calculation Workshop

- ▶ 1. Describe critical teaching content before starting insulin pump therapy
- ▶ 2. Discuss strategies to determine insulin pump basal rates.
- ▶ 3. Discuss how to determine and evaluate bolus rates including coverage for carbs and hyperglycemia.
- ▶ 4. State important safety measures to prevent hyperglycemic crises.
- ▶ 5. List inpatient considerations for insulin pump therapy and CGMs
- ▶ 6. Describe 3 essential steps for emergency preparedness.



# Pump Candidates: Lifestyle Indications and Attributes

- ▶ Erratic schedule
- ▶ Varied work shifts
- ▶ Frequent travel
- ▶ Desire for flexibility
- ▶ Tired of MDI
- ▶ Athletes
  - ▶ Temporary basal adjust
  - ▶ Disconnect options
  - ▶ Waterproof options



ADCES Practice Paper 2021 - [Continuous Subcutaneous Insulin Infusion \(CSII\) Without and With Sensor Integration](#)



# LifeStyle Indications for Candidate or Parents of Pump Wearer

- ▶ Parents and caretakers must have a thorough understanding and willingness and time to understand the pump and work with team to problem solve
- ▶ Willingness to work with healthcare provider during pre-pump training
- ▶ Adequate insurance benefits or personal resources





# LifeStyle Indications for Candidate or Parents of Pump Wearer

- ▶ Physical ability
  - ▶ View pump
  - ▶ Fill and replace insulin cartridge
  - ▶ Insert an infusion set
  - ▶ Wear the pump
  - ▶ Perform technical functions
- ▶ Emotional stability and adequate emotional support from family or others



# Caregiver education about pumps

- ▶ Key Topics
  - ▶ Hypo detection /treatment
  - ▶ Hyperglycemia trouble shooting
  - ▶ Basic bolus procedure
  - ▶ Cartridge set change process
  - ▶ Understand what alarms mean
  - ▶ History recall



# Pre Pump Knowledge / Education

- ▶ Establishment of Goals
- ▶ Competence in Carb counting
- ▶ Insulin Carb Ratios (ICR) & Correction or sensitivity factor (CF)
- ▶ Ability to manage hyper and hypoglycemia
- ▶ Self-adjust insulin
  - ▶ Carbs
  - ▶ Correction
  - ▶ Physical activity
  - ▶ Alcohol intake



# Pre Pump Knowledge / Education

- ▶ Ability to fill and insert cartridge/reservoir and insert and change infusion sets
- ▶ Ability to detect infusion set and site issues
- ▶ Manage sick days, exercise and travel
- ▶ Trouble shoot and ability to solve pump issues
- ▶ Understand BG Data
- ▶ Hypo prevention and treatment
- ▶ Basic of basal bolus therapy and how to switch back to injections if needed



# Poll Question 1

- ▶ Teenagers benefit from insulin pump therapy for the following reason.
- ▶ A. Can increase insulin rate to cover for alcohol intake.
- ▶ B. Decreased risk of glucose emergencies
- ▶ C. Greater dependence on parents
- ▶ D. Match insulin to hormone swings





# Toddlers to Teens Benefit



- ▶ Delayed bolusing for fussy eaters
- ▶ Dosing precision 10ths 20ths and 40ths of a unit
- ▶ Reduced hypo risk
- ▶ Lockout features

## ▶ Teens

- ▶ Basal patterns for hormonal swings
- ▶ Historical data records/ downloading / app sharing
- ▶ Easy snack coverage
- ▶ Greater independence
- ▶ Technical coolness



# Written Plan for Pump Use

- ▶ Blood glucose checks or CGM Checks
- ▶ Record keeping of BG, Carbs, insulin, activity and other issues
- ▶ Site-change guidelines
- ▶ Restart injections if needed
- ▶ When to check ketones and action to take
- ▶ Hypoglycemia and Hyperglycemia treatment guidelines



# CGM Time in Range Recommendations


- ▶ For most with type 1 or type 2 diabetes
  - > 70% of readings within BG range of 70-180mg/dL
  - < 4% of readings < 70 mg/dL
  - < 1% of readings < 54 mg/dL
  - < 25% of readings > 180 mg/dL
  - < 5% of readings > 250 mg/dL
- ▶ For under 25 years, with  $A_{1c}$  goal is < 7.5%, time-in-range target is set to about 60%.



## 6. Glycemic Targets: *Standards of Medical Care in Diabetes—2022*

FREE

American Diabetes Association Professional Practice Committee



*Diabetes Care* 2022;45(Supplement\_1):S83–S96

<https://doi.org/10.2337/dc22-S006>



# Time in Range | Older Adults

- ▶ For older adults or those at high risk for hypoglycemia (ie, hypoglycemic unawareness, cognitive impairment, or comorbidities):

> 50% of BG within 70-180 mg/dL

< 1% of readings < 70 mg/dL

< 10% of readings > 250 mg/dL



6. Glycemic Targets: *Standards of Medical Care in Diabetes—2022*

FREE

American Diabetes Association Professional Practice Committee



*Diabetes Care* 2022;45(Supplement\_1):S83–S96

<https://doi.org/10.2337/dc22-S006>



# Time in Range | Pregnancy

- ▶ For those with type 1 diabetes and pregnant:
  - > 70% of BG readings within 63-140 mg/d
  - < 4% of readings < 63 mg/dL
  - < 1% of readings < 54 mg/dL
  - < 25% of readings > 140 mg/dL



6. Glycemic Targets: *Standards of Medical Care in Diabetes—2022*

FREE

American Diabetes Association Professional Practice Committee



*Diabetes Care* 2022;45(Supplement\_1):S83–S96


<https://doi.org/10.2337/dc22-S006>





# AGP Report: Continuous Glucose Monitoring

## Time in Ranges Goals for Type 1 and Type 2 Diabetes



**Test Patient** DOB: Jan 1, 1970

**14 Days: August 8–August 21, 2021**

**Time CGM Active: 100%**

## Glucose Metrics


**Average Glucose** ..... **175 mg/dL**  
Goal: <154 mg/dL

**Glucose Management Indicator (GMI)** ..... **7.5%**  
Goal: <7%

**Glucose Variability** ..... **45.5%**  
Defined as percent coefficient of variation  
Goal: ≤36%

## Ambulatory Glucose Profile (AGP)

AGP is a summary of glucose values from the report period, with median (50%) and other percentiles shown as if they occurred in a single day.



6. Glycemic Targets: *Standards of Medical Care in Diabetes—2022*

FREE


American Diabetes Association Professional Practice Committee

[Check for updates](#)

Diabetes Care 2022;45(Supplement\_1):S83–S96

<https://doi.org/10.2337/dc22-S006>


# Let's practice calculating basal rates





# Initial Calculations for CSII

TDD: total daily dose



- Start with 1 basal rate, adjust according to glucose trends over 2-3 days
- Adjust to maintain stability in fasting state (between meals & sleep)
- Add additional basals according to diurnal variation (dawn phenomenon)

- Adjust based on low-fat meals with known carbohydrate content
- Acceptable 2-h post-prandial rise is ~60mg/dL above pre-prandial BG
- Adjust carb ratio in 10%-20% increments based on post-prandial BG

- Sensitivity Factor is correct if BG is within 30 mg/dL of target range within 2 hours after correction
- Make adjustments in 10%-20% increments if 2-hr post-correction BGs are consistently above or below target

Hypoglycemia patients – start at lower value of method 1 & 2


Hyperglycemic, elevated A1C or pregnant – start at higher value of method 1 & 2

Consensus Statement by AACE/ACE insulin pump management task force. Endocr Pract. 2014 May; 20(5):463-89.



# Initial Calculations for CSII

TDD: total daily dose



- Start with 1 basal rate, adjust according to glucose trends over 2-3 days
- Adjust to maintain stability in fasting state (between meals & sleep)
- Add additional basals according to diurnal variation (dawn phenomenon)

- Adjust based on low-fat meals with known carbohydrate content
- Acceptable 2-h post-prandial rise is ~60mg/dL above pre-prandial BG
- Adjust carb ratio in 10%-20% increments based on post-prandial BG

- Sensitivity Factor is correct if BG is within 30 mg/dL of target range within 2 hours after correction
- Make adjustments in 10%-20% increments if 2-hr post-correction BGs are consistently above or below target

Hypoglycemia patients – start at lower value of method 1 & 2  
Hyperglycemic, elevated A1C or pregnant – start at higher value of method 1 & 2

Consensus Statement by AACE/ACE insulin pump management task force. Endocr Pract. 2014 May; 20(5):463-89.


TDD = Total Daily Dose

TDI = Total Daily Insulin

# TDD insulin practice – TDD 30 units / 70kg

## Initial Calculations for CSII

TDD: total daily dose



Hypoglycemia patients – start at lower value of method 1 & 2

Hyperglycemic, elevated A1C or pregnant – start at higher value of method 1 & 2

Consensus Statement by AACE/ACE insulin pump management task force. Endocr Pract. 2014 May; 20(5):463-89.

### ► Method 1 (TDD)

► TDD x 0.75

► 30 units x 0.75 = 22.5

### ► Method 2 (Wt)

► Pt wt kg x 0.50

► 70kg x 0.50 = 35

### ► Final daily dose

► A1c 6.3% - Method 1

► A1c 9.2% - Method 2

► A1c 7.5% - Take avg 1 & 2

**Avg 22.5 + 35 = 57.5 / 2 = 28.75 or 29**





# Initial Calculations for CSII

TDD: total daily dose

**METHOD 1**  
Pre-pump TDD

Pre-Pump TDD x 0.75

**STARTING Pump TDD**  
Average method 1 & 2

**METHOD 2**  
Patient Weight

Wt (kg) x 0.50

**BASAL RATE**  
 $\text{Pump TDD} \times 0.5 / 24$

- Start with 1 basal rate, adjust according to glucose trends over 2-3 days
- Adjust to maintain stability in fasting state (between meals & sleep)
- Add additional basals according to diurnal variation (dawn phenomenon)

**CARB RATIO**  
 $450 / \text{Pump TDD}$

- Adjust based on low-fat meals with known carbohydrate content
- Acceptable 2-h post-prandial rise is ~60mg/dL above pre-prandial BG
- Adjust carb ratio in 10%-20% increments based on post-prandial BG

**CORRECTION RATIO**  
 $1700 / \text{Pump TDD}$

- Sensitivity Factor is correct if BG is within 30 mg/dL of target range within 2 hours after correction
- Make adjustments in 10%-20% increments if 2-hr post-correction BGs are consistently above or below target

Hypoglycemia patients – start at lower value of method 1 & 2

Hyperglycemic, elevated A1C or pregnant – start at higher value of method 1 & 2

Consensus Statement by AACE/ACE insulin pump management task force. Endocr Pract. 2014 May; 20(5):463-89.

# Example – LS weighs 80 kg, TDD 50 units, A1c 8.2%

## Method 1 – Based on TDD

- ▶  $50 \times 0.75 = 37.5$  units total daily dose
- ▶  $37.5 \times 0.5 = 18.75$  units for basal
- ▶  $18.75$  divided by  $24$  hrs =  $0.78$  units/hr (Basal rate)

## ▶ Method 2 – Based on body wt

- ▶  $80\text{kg} \times 0.5 = 40$  units
- ▶  $40 \times 0.5 = 20$  units for basal
- ▶  $20$  divided by  $24$  hours =  $0.83$  units/hr (Basal rate)



Which method would you use? **Method 2**



# Example – KL weighs 40 kg, TDD 20 units, A1c 6.2%

## Method 1 – Based on TDD

- ▶  $20 \times 0.75 = \underline{\hspace{1cm}}$  units total daily dose
- ▶  $15 \times 0.5 = \underline{\hspace{1cm}}$  units for basal
- ▶ 7.5 divided by 24 hrs =  $\underline{\hspace{1cm}}$  units/hr (basal rate)

## ▶ Method 2 – Based on body wt

- ▶  $40\text{kg} \times 0.5 = \underline{\hspace{1cm}}$  units
- ▶  $20 \times 0.5 = \underline{\hspace{1cm}}$  units for basal
- ▶ 10 divided by 24 hours =  $\underline{\hspace{1cm}}$  units/hr (basal rate)

Which method would you use?



# Example – KL weighs 40 kg, TDD 20 units, A1c 6.2%

## Method 1 – Based on TDD

- ▶  $20 \times 0.75 = 15$  units total daily dose
- ▶  $15 \times 0.5 = 7.5$  units for basal
- ▶  $7.5 \text{ divided by } 24 \text{ hrs} = .31 \text{ units/hr}$  (basal rate)

## ▶ Method 2 – Based on body wt

- ▶  $40\text{kg} \times 0.5 = 20$  units
- ▶  $20 \times 0.5 = 10$  units for basal
- ▶  $10 \text{ divided by } 24 \text{ hours} = .416$  (.42) units/hr (basal rate)




Which method would you use? **Method 1**



# Example – JR weighs 70 kg, TDD 30 units, A1c 7.5%

Which method would you use?

Figure out the average first



## Method 1 – Based on TDD

- ▶  $30 \times 0.75 = 22.5$  units total daily dose

## ▶ Method 2 – Based on body wt

- ▶  $70\text{kg} \times 0.5 = 35$  units

- ▶ Average =  $22.5 \text{ units} + 35 \text{ units} / 2$   
= 28.7 or 29 units

- ▶  $29 \times 0.5 = 14.5$  units for basal

- ▶ 14.5 divided by 24 hours = 0.6 units/hr (Basal rate)





# Basal insulin

- ▶ Drip of rapid insulin very few minutes
- ▶ If basal rate is set correctly, stable BG between meals and hs
  - ▶ Can skip delay meals
- ▶ Delivered auto on 24 hour cycle
- ▶ Temporary adjustments may include:
  - ▶ lower basal insulin during exercise
  - ▶ increase during sick days



# Basal insulin feedback

- ▶ Keep glucose steady
  - ▶ On average, 5 different basal segments needed
- ▶ Basal insulin rate not correct
  - ▶ Glucose rises or falls even when not eating
  - ▶ Fasting glucose is elevated or low
  - ▶ Correction bolus does not get glucose to target
  - ▶ To prevent hypoglycemia, not covering for snacks
  - ▶ If person is eating to cover for in-between meal hypoglycemia



# Poll Question 2

LS wears an insulin pump and uses lispro insulin. LS has an average basal rate of 0.6 units and hour, a 1:15 carb ratio and a 1:50 correction ratio. Based on the ambulatory glucose profile, LS is experiencing elevated glucose levels from 4am to 7am. To get glucose to target, what is the best next step?



- A. Add basal insulin glargine to prevent Somogyi effect
- B. Make sure LS isn't consuming carbohydrates after 10pm
- C. Ask LS to double check their CGM insertion site
- D. Increase the basal rate to prevent glucose elevations




# Basal Insulin Needs

- ▶ Dawn phenomena
  - ▶ Higher needs from 3-7am for adults
  - ▶ Kids from Midnight to 7am
  - ▶ Basal rate can be adjusted to match sleep and work schedule
- ▶ Traveling – change clock in pump to match new time



# “Typical” Basal Needs

- Growth years: extended peak, evening & overnight
- Adolescent needs >> childhood needs
- Post-growth years: dawn phenomenon
- Senior needs << young adult needs



# Basal Insulin Dosing – Beyond Basics

- ▶ Active, healthy
  - ▶ 35-45% of total daily insulin
- ▶ Less active, lower carb intake
  - ▶ 45-55% of total daily insulin
- ▶ Percentage may increase during puberty
- ▶ Tends to decrease with advanced age
- ▶ Sleep and growth patterns have major influence





# Adjusting basal rates – think ahead

Takes time for basal rate to affect glucose

- ▶ For children: change in basal rate **1 hour prior** to rising or falling glucose
- ▶ For adults: change in basal rate **2 hour prior** to rising or falling glucose
- ▶ Repeat basal test after adjustment

	Current basal level (units /hr)		
	0.0 – 0.45	0.5 – 1.2	>1.2
Modest Rise/Fall (30-60 mg/dl)	.05	0.1	0.2
Large Rise/Fall (>60 mg/dl)	0.1	0.15	0.3

*Gary Scheiner, MS, CDE*



# Poll Question 3

RT is 33 years old and has had diabetes for the past 20 years. RT uses an insulin pump and CGM and works hard to keep A1cs less than 7%. Their most recent A1c increased to 7.9% and RT sets up an appointment with the diabetes specialist for help. After downloading the report, the specialist thinks they have discovered the reason behind the increasing A1c. Which of the following would most likely explain the A1c increase?



- A. Carbohydrate bolus insulin omissions
- B. Basal insulin rate set too high
- C. Bolus insulin given 15 minutes before meal
- D. CGM sensor malfunction




# Bolus Rate Calculations are next

- ▶ I:C
- ▶ Sensitivity
- ▶ Timing
- ▶ Considerations



# Initial Calculations for CSII

TDD: total daily dose



Hypoglycemia patients – start at lower value of method 1 & 2

Hyperglycemic, elevated A1C or pregnant – start at higher value of method 1 & 2

Consensus Statement by AACE/ACE insulin pump management task force. Endocr Pract. 2014 May; 20(5):463-89.



# Bolus Rates - Same for each meal to start

- ▶ CHO Ratio
  - ▶ Start with 1:15 or
  - ▶  $450 \text{ divided by TDD} = \text{I:C Ratio}$
- ▶ Correction/sensitivity
  - ▶  $1700 \text{ divided by TDD}$
- ▶ Active insulin/insulin On Board
  - ▶ 3-6 hours
- ▶ Time in Range target: 70-180 mg/dl
- ▶ Or Target of 120



## Insulin to Carb Ratio I :C 450 / Total Daily Dose

### ► 450 Rule I:C 450/TDD

- 450 divided by total daily insulin dose.
- Equals Gms of carb covered by 1unit insulin.
- Example:  
Ind takes 45 units daily.  
 $450 / 45 = 10$
- 1 unit for 10 grams carb

You try

- JR TDD is 90 units
- 
- 1 unit for \_\_\_\_\_ gms carb

You try

- ML TDD is 15 units
- 
- 1 unit for \_\_\_\_ gms carb





## Insulin to Carb Ratio I :C 450 / Total Daily Dose

### ► 450 Rule I:C 450/TDD

- 450 divided by total daily insulin dose.
- Equals Gms of carb covered by 1unit insulin.
- Example:  
Takes 45 units daily.  
 $450 / 45 = 10$
- 1 unit for 10 grams carb

You try

- JR TDD is 90 units
- $450 / 90 = 5$
- 1 unit for 5 gms carb

You try

- ML TDD is 15 units
- $450/15 = 30$
- 1 unit for 30 gms carb



## Example – JR injects 30 TDD, A1c 6.7%

- ▶ Average =  $22.5 \text{ units} + 35 \text{ units} / 2$   
= 28.7 or 29 units a day
  - ▶  $29 \times 0.5 = 14.5$  units for basal
  - ▶ 14.5 divided by 24 hours = 0.6 units/hr (Basal rate)
- 



### What is his I:C ratio?

- ▶  $450 / 29 = 15.5$
- ▶ I:C Ratio = 15.5



# Insulin /Carb Ratio - How does that work?

TDD 40 units, A1c 8.2%

- ▶ Method 2 – Based on body wt
  - ▶  $80\text{kg} \times 0.5 = 40$  units
  - ▶  $40 \times 0.5 = 20$  units for basal
  - ▶ 20 divided by 24 hours = 0.83 units/hr (Basal rate)



## Calculate Insulin to Carb Ratio

Use 450 rule

$450 / \text{TDD}$

$450 / 40 = 11.25$

(round down to 11)

1 unit Humalog for each 11 gms of carb

Insulin/Carb Ratio I:C 1:11

- ▶ Uses Humalog insulin
- ▶ Dinner
  - ▶ 4 ounces steak
  - ▶ 1 dinner roll
  - ▶ 1 cup mashed potatoes
  - ▶ Few sprigs broccoli
  - ▶ Glass of white wine

How much bolus for this meal?

What if she ate 60 gms?

BG is 220 – Target is 120

# Insulin /Carb Ratio - How does that work?

TDD 40 units, A1c 8.2%

- ▶ Method 2 – Based on body wt
  - ▶  $80\text{kg} \times 0.5 = 40$  units
  - ▶  $40 \times 0.5 = 20$  units for basal
  - ▶ 20 divided by 24 hours = 0.83 units/hr (Basal rate)

## Calculate Insulin to Carb Ratio

Use 450 rule

$450 / \text{TDD}$

$450 / 40 = 11.25$

(round down to 11)

1 unit Humalog for each 11  
gms of carb

Insulin/Carb Ratio I:C 1:11



- ▶ Uses Humalog insulin
- ▶ Dinner
  - ▶ 4 ounces steak
  - ▶ 1 dinner roll (15)
  - ▶ 1 cup mashed potatoes (30)
  - ▶ Few sprigs broccoli
  - ▶ Glass of white wine

How much bolus for this meal?

$45 / 11 = 4.1$  units

What if she ate 60 gms?

$60 / 11 = 5.5$

BG is 220 – Target is 120

# Covering Carbs with Insulin

- ▶ Dose based on:
  - ▶ Grams of carb in meal
  - ▶ Insulin carb ratio or fixed dose?
- ▶ Right dose?
  - ▶ Brings glucose to prebolus glucose level within 3-4 hours
  - ▶ If BG rises more than 60 - 80 points 2 hours post meal, needs adjustment
  - ▶ If BG falls more than 30 points 2 hours post meal, may need adjustment
  - ▶ Adjust in small increments (10-20% ideal)

← ← ← If glucose rising post-meal ← ← ←

1:3 1:4 1:5 1:6 1:7 1:8 1:9 1:10 1:12 1:14 1:16 1:18 1:20 1:25 1:30 1:35 1:40 1:50

→ → → If glucose dropping post-meal → → →



# But wait... what about correction insulin for current glucose level? $1700/\text{TDD} - \text{Target } 120$

TDD = 40 units  
BG target is 120.  
Current BG is 220.  
Based on her current BG, how much correction insulin does she need to get to target?



- ▶ Correction/sensitivity
  - ▶ 1700 divided by TDD
  - ▶  $1700 / 40 = 42.5$  or 43
  - ▶ **Correction: 1 unit of insulin lowers BG 43 points.**

$$220 - 120 = 100 \text{ over target}$$

$$100 / 43 = 2.3 \text{ units to correct for hyperglycemia}$$

What if her BG is 320?

$$320 - 120 = \underline{\hspace{2cm}} \text{ over target}$$

$$\underline{\hspace{2cm}} \text{ units to correct for hyperglycemia}$$



# But wait... what about correction insulin for current glucose level? $1700/\text{TDD} - \text{Target } 120$

TDD = 40 units  
BG target is 120.  
Current BG is 220.  
Based on her current BG, how much correction insulin does she need to get to target?



- ▶ Correction/sensitivity
  - ▶ 1700 divided by TDD
  - ▶  $1700 / 40 = 42.5$  or 43
  - ▶ **Correction: 1 unit of insulin lowers BG 43 points.**

$$220 - 120 = 100 \text{ over target}$$

$$100 / 43 = 2.32 \text{ units to correct for hyperglycemia}$$

What if her BG is 320?

$$320 - 120 = 200 \text{ over target}$$

$$200/43 =$$

$$4.65 \text{ units to correct for hyperglycemia}$$

# Correction Insulins Example

Correction Factor Fine-Tuning


Mathematical Approach

The lower the TDI = more insulin sensitive

Correction Scale / TDI	Sensitivity mg/dl 30 units	Sensitivity mg/dl 40 units	Sensitivity mg/dl 50 units
Aggressive (1500) 1500 / TDI	50 ?	38	? 30
Common (1700) 1700 / TDI	57	43	34
Conservative (2000) 2000 / TDI	67	? 50	? 40



# Stretch Break




*REMEMBER:*  
*Stretch*  
*Often!*

---



# Initial Calculations for CSII

TDD: total daily dose



Hypoglycemia patients – start at lower value of method 1 & 2

Hyperglycemic, elevated A1C or pregnant – start at higher value of method 1 & 2

Consensus Statement by AACE/ACE insulin pump management task force. Endocr Pract. 2014 May; 20(5):463-89.



# But wait, what about IOB?

- ▶ Method 2 – Based on body wt
  - ▶  $80\text{kg} \times 0.5 = 40$  units
  - ▶  $40 \times 0.5 = 20$  units for basal
  - ▶ 20 divided by 24 hours = 0.83 units/hr (Basal rate)

Insulin to Carb Ratio I:C

450 / TDD

$450 / 40 = 11$

I:CR = 1:11

Correction/sensitivity

1700 divided by TDD

$1700 / 40 = 42.5$

Correction: 1:43 points.





# Active Insulin time - IOB

- ▶ How much “insulin on board” IOB to prevent stacking and hypoglycemia
- ▶ Typical active insulin time is 3-5 hours
  - ▶ Average about 4 hours
- ▶ Action time shorter in leaner, young, active individuals in hot climates
- ▶ Action time is longer, 6-8 hours, for those with renal disease or using regular insulin
- ▶ Careful monitoring or CGM to eval if bolus rates set correctly





# Pump Bolus Estimate Features

- ▶ Based on glucose and carb data entered by user

## Bolus Estimate Details

Total	8.1 U
Food intake	75 gms
BG	220
Food Dose	6.8 U
Correction Dose	2.3 U
Insulin-On-Board	1.0 U
(Based on BG and Carbs entered by user.)	

ICR 1:11 gms

Correction 1 unit for 43

Target BG 120

Active insulin on board (IOB)  
subtracted from the correction

$75 \text{ gms carb} / 11 = 6.8 \text{ units}$

$\text{Correction } 220 - 120 = 100 / 43 = 2.3 \text{ units}$

$\text{IOB} = 1 \text{ unit}$

$6.8 + 2.3 = 9.1 - 1 \text{ units} = 8.1 \text{ units}$

Bolus delivery of 8.1 units



# What bolus would this person need?

- ▶ Plans to eat 75 gms Carb Snack
- ▶ BG is 68

## Bolus Estimate Details

Total	U
Food intake	75 gms
BG	68
Food Dose	? .0 U
Correction Dose	? .0 U
Insulin-On-Board	2.0 U
(Based on BG and Carbs entered by user.)	

ICR 1:15 gms

Correction 1 unit for 50

Target BG 100

Active insulin on board (IOB)  
subtracted from the correction

75 gms carb/15 = \_\_\_\_\_ ?units

Correction \_\_\_\_ -100 = \_\_\_\_ /50 ?units


IOB = 2 unit

Total insulin = \_\_\_\_\_ ?units



# Poll Question 4

- ▶ For case study, how much bolus insulin?
  - ▶ A. 3.6 units
  - ▶ B. 2.4 units
  - ▶ C. 4 units
  - ▶ D. Determine activity first



# What bolus would this person need?

- ▶ Plans to eat 75 gms Carb meal
- ▶ BG is 68

## Bolus Estimate Details

Total	2.4 U
Food intake	75 gms
BG	68
Food Dose	5.0 U
Correction Dose	-.64 U
Insulin-On-Board	2.0 U
(Based on BG and Carbs entered by user.)	

ICR 1:15 gms

Correction 1 unit for 50

Target BG 100

Active insulin on board (IOB)  
subtracted from the correction

$75 \text{ gms carb} / 15 = 5 \text{ units}$

$\text{Correction } 68 - 100 = -32 / 50 = -.64 \text{ units}$

$\text{IOB} = 2 \text{ unit}$

$\text{Total insulin} = 2.4 \text{ units}$





# Not using insulin/carb bolus ratios?

- ▶ Fixed dosing
  - ▶ Take half of total daily dose, divide by number of meals to get fixed dose per meal
  - ▶ Calculate insulin sensitivity correction factor
    - ▶  $1700 \div$  by total daily insulin
  - ▶ No target BG – choose acceptable target range



40 units x 0.5 for basal and bolus  
20 units/24 for basal = 0.83 hr  
20 units for bolus  
20 units/3 meals  
7, 6, 7 units per meal plus correction  
Correction  $1700/40$  units = 1:43



# Advanced Pump Features

- ▶ Prolonged bolus for
  - ▶ Gastroparesis, amylin, GLP-1 Receptor Agonists
- ▶ Advanced Basal Features
  - ▶ Temporary basal rates
  - ▶ Secondary, tertiary programs
- ▶ Custom alerts examples
  - A1c of 13% - Alarm at 70
  - A1c of 8% - Alarm 70 – 300
  - A1c of 7 % - Alarm 70-250
- ▶ Data downloads



# Prolonged bolus

## ▶ Standard bolus

- ▶ Delivered within a few minutes
  - ▶ Peaks in one hour
  - ▶ Lasts for 4 hours
- 

## ▶ Prolonged bolus

- ▶ Delivered over a couple of hours
- ▶ Peak delay
- ▶ Duration extended

## ▶ Purpose

- ▶ Match insulin to absorption of food
- ▶ Works well with slowly digested food


## ▶ Applications

- ▶ Large portions
- ▶ Slow consumption
- ▶ Gastroparesis
- ▶ Use of incretin mimetics



# Prolonged bolus

- ▶ Square/extended
  - ▶ None of the bolus is delivered up front
  - ▶ Common timing is 1-2 hours after start of meal
  - ▶ Can last for up to 8 hours
- ▶ Dual/combo/combination bolus
  - ▶ 30% delivered up front, the rest of bolus over the next several hours.
  - ▶ Lasts about 5 hours



# Insulin coverage for protein?

- ▶ Most of time, protein won't affect glucose
- ▶ If person on low carb diet, protein may start impacting blood glucose levels
  - ▶ Bolus for 50% of protein grams
- ▶ If large protein portion consider extended bolus





# Problem solving

- ▶ Prevent missed boluses
  - ▶ 1 missed meal bolus over a month raises A1c 0.5%
  - ▶ Get in habit of pre-bolusing – 15 minutes before meal works best
  - ▶ Use reminder alerts on pumps
- ▶ If basal or bolus is more than 65% of total daily dose, usually indicates need to recalculate ratios



# Disconnecting from Pump

- ▶ BG rises about 1 mg/dl a minute when disconnected
- ▶ Avoid extended disconnection since can lead to ketones and hyperglycemia
- ▶ Strategies
  - ▶ Short term disconnection < 1 hour
    - ▶ Bolus to replace missed basal insulin
  - ▶ Long term >1 hour and bolus missed basal insulin hourly
  - ▶ Protective caps usually not necessary



With pump therapy, there is no background insulin on board



# Safety guidelines

- ▶ Review signs and treatment of hypo
- ▶ If frequent lows, may want to set pump alarm at 90
  - ▶ Try not to suspend pump when low, unless no treatment available
- ▶ Diabetes Ketoacidosis
  - ▶ Those with negative c-peptide at higher risk
  - ▶ Insulin pump interruption for 2-3 hours can lead to DKA
  - ▶ Provide education to prevent, detect and reverse



# Poll Question 5

- ▶ AL is on an insulin pump. Her BG at 10am is 108, at 11am, 219 and noon 298. She has not eaten anything since breakfast. What is best action?
- ▶ A. Program insulin pump to deliver 3 units bolus stat
- ▶ B. Increase basal rate starting at 8am
- ▶ C. Go to emergency room
- ▶ D. Check for ketones




# Prevent DKA and Hyperglycemia

- ▶ Eval sites for malabsorption, make sure to change site and infusion sets every 2-3 days
- ▶ Protect insulin from overheating
- ▶ Tubing or infusion set clogs – change site
- ▶ Check for leaks, smell for insulin, use angled sets
- ▶ Make sure to purge air bubbles before priming tube
- ▶ Inspect daily for dislodgement
- ▶ Correct priming technique when changing infusion set
- ▶ Extended pump suspension or disconnect?
- ▶ Limit suspension to one hour, always have back-up syringes

Figure 1. MiniMed 530G System



# Action in Case of Hyperglycemia for Pump Users





# Ketone Testing Options

- ▶ Urine ketostix or diastix
  - ▶ More than 15 mg/dl = positive ketones
- ▶ Blood sampling
  - ▶ Novamax or Precision Xtra blood meter
  - ▶ More than 0.5 mmol/l  $\beta$  – hydroxybutyrate indicates action and insulin needed




<https://www.novabio.us/nova-max-plus/>



Diabetes Education  
SERVICES

## Keeping connected - Pump Users need to contact clinical staff if:

- ▶ Severe or repeated hypo
- ▶ Ketosis
- ▶ Signs of infection
- ▶ Call pump company if technical difficulties
- ▶ See pumper in 1-2 weeks, download device, troubleshooting
- ▶ At 3-4 weeks review more advanced features




# Hospital Stay for Insulin Pump Users

- ▶ Staff to assess:
  - ▶ How long using pump?
  - ▶ Who adjusts pump settings?
  - ▶ What type of insulin is used?
  - ▶ How much insulin is in pump now?
  - ▶ When is next site change? Who does it?
  - ▶ Basal rates? I:C ratios? Correction?
  - ▶ Have your supplies?
  - ▶ When usually check BG or CGM?



### Patient With Insulin Pump Admitted to Hospital



### Changes to Pump Therapy With Imaging Studies

X-ray/CT	Pump should be covered by lead apron
MRI	Pump and metal infusion set should be removed
Ultrasound	No need to remove pump but transducer should not be pointed directly at the pump
Cardiac catheterization	Pump should be covered by lead apron
Pacemaker/automatic implantable cardioverter defibrillator (AICD)	Pump should be covered by lead apron
Colonoscopy/EGD	Pump can remain in place
Laser surgery	Pump can remain in place

# Hospital Stay - Need orders

- ▶ Backup plan in case pump can't be used
- ▶ Don't stop pump without administering rapid insulin first (or IV insulin).
- ▶ Designate surrogate programmer(s)
- ▶ Specify frequency and carb count for meals/snacks
- ▶ Keep pump and programmer outside room during MRI, CT Scan, Xray.
- ▶ Don't aim Echo/US transducer at pump
- ▶ CGM - Remove infusion set and sensor for MRI
- ▶ Hospital meter to determine BG levels



Refer to individual tech user manual for more detailed info

# Pumpers Responsibility in Hospital

- ▶ Provide own pump (and sensor) supplies
- ▶ Change pump reservoirs and infusion sets
- ▶ Provide staff with SMBG and insulin doses
- ▶ Notify staff of adjustments to standard doses
- ▶ Respond to alarms





# Backup Plan if pump isn't working

- ▶ Immediate basal insulin injection
- ▶ Mealtime rapid insulin injection
- ▶ Keep written log of I:C ratios, correction and meal boluses
- ▶ Keep log of off-pump activity
- ▶ Resume pump when basal insulin wears off



# Poll Question 6

- ▶ JL is on an insulin pump and CGM and asks the diabetes educator how to best prepare for emergency situations. What is the most critical step to take in case of an emergency evacuation?
- ▶ A. Have back up energy source
- ▶ B. Keep insulin on ice
- ▶ C. Know the CDCs info line number
- ▶ D. Alert local emergency responders of status



# MY DIABETES EMERGENCY PLAN

AN IMPORTANT  
CHECKLIST FOR  
PEOPLE WITH  
DIABETES.




# EARTHQUAKE



# Medical Diabetes Identification

- ▶ Speaks when you cannot
- ▶ Necklace, bracelet or watch band
- ▶ A wallet card is additional identification only





# Prepare A Portable Emergency Kit

# MY DIABETES EMERGENCY PLAN

**Prepare a portable, insulated and waterproof diabetes emergency kit that contains the following items:**

- ☐ List of the following information:

  - *Type of diabetes*
  - *All of your medical conditions, allergies and prior surgeries*
  - *All medications (include pharmacy contact information, active prescription information and eligible refills)*
  - *Previous diabetes medications and reason for discontinuation*
  - *Contact information for all your health care professionals and for at least two emergency contacts*

☐ Letter from your diabetes healthcare professionals with most recent diabetes medication regimen (especially if taking insulin), health insurance card, living will, healthcare power of attorney, etc.

☐ Most recent laboratory results (especially A1C, kidney and liver tests)

☐ If possible, a 30-day supply of all medications taken by mouth or injection for diabetes as well as all other medical conditions

  - *Include insulin and a severe hypoglycemia emergency (e.g., glucagon) kit—if prescribed (always check expiration date)*

☐ Blood glucose testing supplies including, if possible, 2 glucose meters with extra batteries

☐ A cooler with room for 4 freezeable gel packs, insulin and unused injectable medications to be added when ready to go

  - *Note: Do not use dry ice and avoid freezing the medication*

☐ 14-day supply of diabetes medications (for example, glucose tablets, 6 oz. juice boxes, glucose gel, regular soda, sugar, honey or hard candy)

☐ A 2-day supply of nonperishable food (for example, peanut butter or cheese crackers, meal replacement shakes or bars, etc.)

☐ At least a 3-day supply of bottled water

☐ Pen/pencil and notepad to record blood sugar, other test results and any new signs/symptoms suggesting medical problems

☐ First aid supplies such as bandages, cotton swabs, dressings and topical medications (antibiotic ointments or creams)

**Other recommendations:**

☐ Wear shoes and socks while awake and examine your feet often for cuts, sores, red spots, swelling, blisters, calluses and infected toenails or any unusual condition

☐ Make sure that all vaccinations, including tetanus, are up-to-date

☐ Pack extra comfortable clothing, including undergarments

☐ Take a mobile phone with an extra charger or extra batteries for you and family members

☐ Choose a designated meeting place in case you are separated from your family and/or significant others and are unable to reach them by phone



# www.diabetesdisasterresponse.org

## DO YOU OR A LOVED ONE HAVE DIABETES AND USE INSULIN?

### Make a plan to stay healthy during natural disaster or emergency

Managing diabetes can be even harder when you are dealing with a major storm, loss of electricity, and possible evacuation from your home. Building a "diabetes kit" now can save a lot of worry and time when a disaster strikes. A checklist template is included for your use.



**Your diabetes kit can be stored in an easy-to-carry waterproof bag or container** to hold the documents, information, and supplies that you will want to have with you.

### Important Information to Keep In Your Kit - Write down or copy the following:

- Type of diabetes you have
- Other medical conditions, allergies, and previous surgeries
- Current medications, doses, and time you take them. Include your pharmacy name, address and phone number.
- Previous diabetes medications you have taken
- A letter from your diabetes care team with a list of your most recent diabetes medications, if possible.
- A copy of your most recent laboratory result, like A1C results
- Make, model and serial number of your insulin pump or CGM. Include pump manufacturer's phone number in case you need to replace your device.
- Doctor's name, phone number, and address
- Phone numbers and email addresses for your family, friends, and work. Include out-of-town contacts.
- A copy of your health insurance card
- A copy of your photo ID
- Cash



Please check out this Diabetes Disaster Response Resource Page.

Let's help get people ready for the worst.





# Disaster Readiness

- ▶ **American Red Cross**

**Shelters:** Contact the American Red Cross directly at 1-800-RED-CROSS.

- ▶ **Resource For Health Care Providers:**

- ▶ **Insulin Supply Hotline:** During a disaster, call the emergency diabetes supply hotline 314-INSULIN (314-467-8546) if you know of diabetes supply shortages in your community (i.e. shelter, community center). Hotline is for health care providers only.



# Disaster Readiness



- ▶ **Have an Emergency Diabetes Kit Ready:**
- ▶ People with Diabetes can download the Diabetes Disaster Response Coalition's (DDRC) [Diabetes Preparedness Plan](#).
- ▶ **Stay Updated:** Visit [JDRF Disaster Relief Resources](#) and Diabetes Disaster Response Coalitions [Facebook page](#) with information on how to access medical support, shelters, and open pharmacies during time of disaster.
- ▶ **Know where to get help:**
- ▶ Call 1-800-DIABETES (800-342-2383).
- ▶ American Diabetes Association Center is open, MON.-FRI. 9 a.m. TO 7 p.m. ET.
- ▶ Representatives regularly updated with information on how to access medical support, shelters, pharmacies



# Join us for Level 4 Courses & ReVive 5

- ▶ Solving Glucose Mysteries – Type 2
- ▶ Insulin Calculation workshop
- ▶ Type 2 Diabetes Intensive
- ▶ Basal Bolus in Hospital
- ▶ Getting to the Gut
- ▶ Cancer and Diabetes



## REViVE 5 Diabetes Training Program

Unlocking Hidden Barriers to Diabetes Self-Management

- ▶ Assess diabetes distress and other barriers to self-management.
- ▶ Identify negative self-talk and explore a more positive inner conversation.
- ▶ Develop skills to foster a new narrative using self-compassion.
- ▶ Optimize glucose self-management—"find the expert within."
- ▶ Create a plan for next steps based on different choices & individual values.

Starts November 1<sup>st</sup>  
4 Interactive Sessions



# Thank You



► Please email us with any questions.

[bev@diabetesed.net](mailto:bev@diabetesed.net)

► [www.diabetesed.net](http://www.diabetesed.net)



*Diabetes Education*  
SERVICES

# Extra info for further reading



# Travel Suggestions from Diabetes.org


- ▶ Review TSA's website for travel updates
- ▶ Download [My TSA Mobile App](#)
- ▶ Whenever possible, bring prescription labels for medication and medical devices (while not required by TSA, making them available will make the security process go more quickly)
- ▶ Consider printing out and bringing an optional [TSA Disability Notification Card](#).





# What about diabetes Tech and Security?

- ▶ Refer to training manual for each manufacturer
- ▶ To be safe, ask for pat down if wearing pump, CGM or both




# Travel Suggestions from Diabetes.org

- ▶ Pack medications in a separate clear, sealable bag. Bags that are placed in your carry-on-luggage need to be removed and separated from your other belongings for screening.
- ▶ Keep a quick-acting source of glucose to treat low blood glucose as well as an easy-to-carry snack such as a nutrition bar
- ▶ Carry or wear medical identification and carry contact information for your physician



# Travel: What items allowed?

- ▶ Insulin and insulin loaded dispensing products (vials or box of individual vials, jet injectors, biojectors, epipens, infusers and preloaded syringes)
- ▶ Unlimited number of unused syringes when accompanied by insulin or other injectable medication
- ▶ Lancets, blood glucose meters, blood glucose meter test strips, alcohol swabs, meter-testing solutions
- ▶ Insulin pump and insulin pump supplies (cleaning agents, batteries, plastic tubing, infusion kit, catheter and needle)—insulin pumps and supplies must be accompanied by insulin



# Travel: What items allowed?

- ▶ Glucagon emergency kit, Urine ketone test strips
- ▶ Unlimited number of used syringes when transported in Sharps disposal container or other similar hard-surface container
- ▶ Sharps disposal containers or similar hard-surface disposal container for storing used syringes and test strips
- ▶ Liquids (to include water, juice or liquid nutrition) or gels
- ▶ Continuous blood glucose monitors
- ▶ All diabetes related medication, equipment, and supplies

