

Cholesterol Medications

LDL Lowering Medications

Class / Action	Generic / Trade Name	Usual Daily Dose Range	LDL % Lowering	Considerations
“Statins” HMG- CoA Reductase Inhibitors Inhibits enzyme that converts HMG-CoA to mevalonate - limits cholesterol production	Atorvastatin / Lipitor*	10 – 80 mg	20- 60	Lowers TGs 7-30% Raise HDL 5-15% Take at night. Side effects: weakness, muscle pain, elevated glucose levels. Review package insert for specific dosing adjustments based on drug, food interactions (ie grapefruit).
	Fluvastatin / Lescol* Lescol XL	20 – 80 mg 80 mg	20- 35	
	Lovastatin* Mevacor Altoprev XL	20 - 80 mg 10 - 60 mg	20- 45	
	Pravastatin / Pravachol*	10 - 80 mg	20- 45	
	Rosuvastatin / Crestor	5 – 40 mg	20- 60	
	Simvastatin / Zocor*	20 – 80 mg	20- 55	
	Pitavastatin / Livalo	2 – 4 mg		
Bile Acid Sequestrants Action: Bind to bile acids in intestine, decreasing cholesterol production. Secondary action – raise HDL	Cholestyramine/ Questran*	4 to 16 g per day powder – 1 scoop 4g	Lower LDL by 15-30%	May raise TG levels. Raise HDL 3-5%. Avoid taking in same timeframe w/ other meds – may affect absorption (see package insert). Side effects: GI in nature
	Colesevelam / Welchol Lowers A1c 0.5%	3.75 x 1 daily 1.875 x 2 daily (625mg tablets)		
	Colestipol / Colestid	2 - 16 gms per day tabs Powder – 1 scoop = 5g 5 to 20 gm per day Mix w/ fluid		
Cholesterol Absorption Inhibitors	Ezetimibe / Zetia	10 mg – 1x daily	15-20%	Usually used in combo w/statin. Headache, rash.
Plant Stenols	Benecol	3 servings daily	14%	Well tolerated
Plant Sterols	Take Control	2 servings daily	17%	

Triglyceride Lowering / HDL Raising Medications

If TG > 500, lower TG first, then reduce LDL.

Class / Action	Generic / Trade	Usual Daily Dose Range	Lowers TG	Considerations
Fibrates or Fibric Acids Reduces liver lipogenesis	Fenofibrate/ Tricor Multiple brand formulations	48-145 mg 1x daily Please refer to individual package insert for dosing	20-50%	Lowers LDL 5-20% Raise HDL 10-20% GI side effects, myopathy Avoid w/ severe renal or hepatic disease
	Gemfibrozil / Lopid*	600mg 2x daily		
Nicotinic Acid Raise HDL/Lower TG Inhibits mobilization of free fatty acid	Niacin (immediate release)*	1.5- 3 gms	20-50%	Raise HDL 15-35% Flushing, hyperglycemia, hepatotoxicity – monitor liver enzymes. Can take w/aspirin to < flushing
	NiaSpan (extended release) Niacin (sustained release)	1-2 gms		
Omega 3 Fatty Acid	Omega 3 Acid/ Lovaza	4 gm a day	45%	Raise HDL 9% - Primary use for TG > 500

Combination Medications

Vytorin	Zetia + Simvastatin	Observe precautions of each component drug
Juvisync	discontinued	

*indicates medication is available in generic form.

Antihypertensive Medications

ACE and ARBs are preferred therapy for diabetes with hypertension and albuminuria – If B/P not at goal with either of these agents, add a diuretic or other class. Do not use during pregnancy or in persons w/ renal or hepatic dysfunction. Start w/ low dose, gradually increase. If one class is not tolerated, the other should be substituted. For those treated with an ACE inhibitor, angiotensin receptor blocker, or diuretic, serum creatinine/estimated glomerular filtration rate and serum potassium levels should be monitored at least annually. ADA Standards CV Disease Risk Management

Class / Action	Generic / Trade Name	Usual Daily Dose Range	Frequency	Considerations
ACE Inhibitors Angiotensin Converting Enzyme Action - Block the conversion of AT-I to AT-II. Also stimulates release of nitric oxide causing vasodilation.	benazepril / Lotensin [†]	10 – 40 mg	1 x a day	Try to take same time each day. Effects seen w/in 1 hr of admin, max effects in 6 hrs. Side effects: Can cause cough (due to increased bradykinin) – can try different med in same class. Also can cause fatigue, dizziness, hypotension. [†] These meds are also available as a combo w/ low dose HCTZ (hydrochlorothiazide). [‡] These meds are also available as a combo w/ CCB (calcium channel blocker) usually amlodipine
	captopril /Capoten* [†]	12.5 - 100 mg	2-3 x a day	
	Enalapril/ Vasotec* [†]	2.5 - 40 mg	1-2 x a day	
	Fosinopil / Monopril [†]	10- 40 mg	1 x a day	
	Lisinopril * [†] Prinivil Zestril	10 – 40 mg 10 - 40 mg		
	Ramipril / Altace* [†]	2.5 – 10 mg		
	Moexipril / Univasc [†]	3.75 - 15 mg		
	Perindopril/Aceon [‡]	2-16 mg		
	Perindopril/ Indapamide combo (Coversyl)	2 - 8 mg 0.625 - 2.5 mg		
	Quinapril /Accupril [†]	5 – 40 mg		
Trandolapril/ Mavik	1.0 – 4 mg			
Trandolapril/ Verapamil combo (TARKA)	1-4 mg 180 to 240 mg			
ARBs -Angiotensin Receptor Blockers Action -Block AT-I receptor which reduces aldosterone secretion and vasoconstriction	Azilsartan/Edarbi	40 - 80 mg	1 x daily	Try to take same time each day Side effects- Can cause dizziness, drowsiness, diarrhea, hyperkalemia, hypotension. [†] These meds are also available as a combo w/ low dose HCTZ (hydrochlorothiazide). [‡] These meds are also available as a combo w/ CCB (calcium channel blocker) usually amlodipine
	Azilsartan/ Chlorthalidone combo (Edarbyclor)	40 mg 12.5 - 25 mg		
	Candesartan/Atacand [†]	8 – 32 mg		
	Eprosartan/Teveten [†]	400 - 600 mg		
	Irbesartan/ Avapro [†]	75 – 300 mg		
	Losartan / Cozaar* [†]	25 – 100 mg		
	Olmesartan / Benicar ^{†‡} Tribenzor (triple combo)	20 – 40 mg		
	Telmisartan / Micardis	20 – 80 mg		
Valsartan / Diovan ^{†‡} Exforge HCT (triple combo)	80 – 320 mg			
Valsartan/ Nebivolol combo (Byvalson)	80 mg 5 mg			

*indicates medication is available in generic form.

Class / Action	Generic / Trade Name	Usual Daily Dose Range	Frequency	Considerations
DRIs - Direct Renin Inhibitors -	Aliskiren / Tekturna†	150 – 160 mg	1 x daily	Generally well tolerated. †These meds are also available as a combo w/ low dose HCTZ (hydrochlorothiazide).

Beta Blockers are commonly prescribed as an add-on to other B/P meds for people with DM. Beta Blockers are beneficial for persons w/ concurrent cardiac problems and prevention of recurrent MI and heart failure. Caution in DM since Beta Blockers can cause hyperglycemia and mask hypoglycemia induced tachycardia (but do not block hypoglycemia related dizziness and sweating). Monitor B/P, heart rate, lipids and glucose.

Beta Blockers <i>β1- Selective</i> Action: Blockade β1 receptors & reduce cardiac output & kidney renin activation.g	Acebutolol / Sectral*	200 – 800 mg	2 x daily	Side Effects: Usually CNS related including sedation, dizziness, lightheaded . Watch for bradycardia, hypotension, depression and sexual dysfunction. Check heart rate each visit, adjust dose if HR <50. Can cause heart block – review package insert for drug-drug interactions. Watch for exercise intolerance. When stopping beta blockers, taper dose gradually. Use cautiously at lowest dose. †These meds are also available as a combo w/ low dose HCTZ (hydrochlorothiazide).
	Atenolol / Tenormin*	25 – 100 mg	1 x daily	
	Atenolol with Chlorthalidone/ Tenoretic	50 -100 mg 25 mg	1 x daily 1 x daily	
	Betaxolol / Kerlone	5 – 10 mg	2 x daily	
	Bisoprolol/ Zebeta†	2.5 – 10 mg	1 x daily	
	Metoprolol tartate/Lopressor*†	25 – 100 mg		
	Metoprolol succinate / Toprol XL	25 - 100 mg		
Beta Blockers <i>Non Selective</i> Action: Blockades β1 & β2	Nebivolol/Bystolic	5 to 40 mg	1 x daily	
	Nebivolol with Valsartan/ Byvalson	5 mg 80 mg		
	Nadolol / Corgard*	40 - 120 mg		
	Nadolol with Bendroflumethiazide	40-80 mg 5 mg		
	Penbutolol / Levatol	10 - 40 mg		
	Pindolol / Visken	10 – 40 mg		
Combined α- and β- Blockers	Propranolol / Inderal*	40 – 160 mg	2 x daily	
	Inderal LA (extended)	60 – 180 mg	1 x daily	
	Timolol / Blocadren*	10 – 60 mg	2 x daily	
Combined α- and β- Blockers	Corvedilol / Coreg	6.25 – 50 mg	2 x daily	
	Coreg CR	20 – 80 mg	1 x daily	
	Labetalol / Normodyne*	100 – 2400 mg	2 x daily	

Diuretics are often used as adjunct therapy. Obtain baseline B/P, electrolytes, uric acid, glucose and lipids prior to starting and periodically. May require supplementation w/ magnesium and potassium.

Class / Action	Generic / Trade Name	Usual Daily Dose Range	Considerations
Thiazide Diuretics Action: cause diuresis and decrease vascular resistance. (Many meds combined with this class)	Hydrochlorothiazide (HCTZ)* HydroDIURIL Microzide	12.5 – 25 mg Most frequently prescribed	1 x daily in am with or w/out food Side effects: lyte imbalances; hypokalemia, hypomagnesemia, hyperuricemia, hyperglycemia, hyperlipidemia and hyper/hypocalcemia. S/S include muscle cramps, fatigue, dizziness and cardiac arrhythmias .
	Chlorthalidone / Clorpres*	12.5 – 25 mg	
	Metolazone / Zaroxolyn*	2.5 – 20 mg	
	Indapamide / Lozol*	1.2 – 2.5 mg	
Class / Action	Generic / Trade Name	Usual Daily	Considerations

*indicates medication is available in generic form.

		Dose Range		
Loop Diuretics (resistant HTN)	Furosemide/Lasix*	20 – 600 mg 2x day	Side Effects as above, but more intense. Need K ⁺ supplement. Used if GFR < 30 or if greater diuresis is needed	
	Torsemide / Demadex*	2.5 – 200 mg 1x day		
	Bumetanide / Bumex*	0.5 – 10 mg 2 x day		
Potassium Sparing Diuretics	Amiloride / Midamor	5 – 20 mg	1 x day	Usually combined with thiazide diuretic to balance serum potassium. Alone, they do little to lower BP.
	Triamterene / Dyrenium	37.5 – 75 mg	1 x day	
	Spironolactone / Aldactone*	25 – 100 mg	1-2 x day	
	Eplerenone / Inspra	50 - 100 mg	1 -2 x day	

Calcium Channel Blockers are usually second or third line BP med for diabetes, since they have less impact on CVD. They may also be used for those who can't tolerate ACE or ARB Therapy.

Class / Action	Generic / Trade Name	Usual Daily Dose Range	Frequency	Considerations
Calcium Channel Blocker <i>Nondihydropyridine</i> Relaxes coronary blood vessels to decrease heart rate and cardiac output.	Diltiazem immediate release*	30 – 360 mg	4 x day	Monitor BP, heart rate, liver enzymes and cardiac function a baseline and periodically. Take at the same time each day (with meals if possible). Take in evening if experience drowsiness. Side Effects: Watch for cardiac conduction abnormalities, bradycardia, CHF and edema. Can cause peripheral edema and constipation. Metabolized through CYP3A4, so review package insert for drug and food interactions (ie grapefruit).
	Diltiazem extended release*			
	Cardizem CD	120 – 480 mg	1 x day	
	Tiazac	120 – 540 mg	1 x day	
	Dilacor, Diltia	180 – 540 mg	1 x day	
	Verapamil immediate release*			
	Calan	80 -320 mg	3 x day	
Verapamil sustained release*				
Calan SR, Verelan	120 mg – 480 mg	1 -2 x day		
Verapamil extended release*				
Covera-HS	120 – 480 mg	1 x day		
Verelan PM	100 – 400 mg			
Calcium Channel Blocker – <i>Dihydropyridine</i> Causes vasodilation and decreases peripheral vascular resistance.	Amlodipine/Norvasc	2.5 – 10 mg	1 x day	
	Felodipine / Plendil	2.5 – 10 mg	1 x day	
	Isradipine controlled release DynaCirc CR	2.5 – 10 mg	1 x day	
	Nicardipine sustained release / Cardene SR	30 – 60 mg	2 x day	
	Nifedipine long-acting* Adalat CC /Procardia XL	30 – 120 mg	1 x day	
	Nisoldipine / Sular	10 – 40 mg	1 x day	

α1 – Receptor Blockers - Often used for pts with DM & benign prostatic hypertrophy (BPH).

α1 – Receptor Blockers Vasodilation	Doxazosin/Cardura*	1 – 8 mg	1 x day	Take at hs and low dose to reduce risk of postural hypotension/syncope.
	Prazosin / Minipress*	2 – 20 mg	2 - 3 day	
	Terazosin/ Hytrin*	1 – 10 mg	1 – 2 day	

α2 agonists- Not usually first line due to side effects. Effective in pts w/ renal disease, since does not compromise renal function.

α2 agonists – Centrally act to block influence of norepinephrine on the heart and lower B/P	Clonidine / Catapres*	0.1 to 0.8 mg	2 x day	Administer w/ diuretic. Side effects: sedation, dry mouth, bradycardia orthostatic hypotension, impotence. Do not stop abruptly, can cause hypertensive crisis.
	Methyldopa / Aldomet*	250 – 1000 mg	2-3 x day	

*indicates medication is available in generic form.

New Lipid Lowering Medications

Contributor: Diana Isaacs, PharmD, BCPS, BCACP, BC-ADM, CDCES, FADCES, FCCP 2022

PCSK9 Inhibitors Lipid Medications		
Proprotein convertase subtilisin/kexin type 9		
	Alirocumab (Praluent)	Evolocumab (Repatha)
FDA-approved indications	<ul style="list-style-type: none"> Primary hyperlipidemia (HLD) Homozygous familial hypercholesterolemia (HoFH) Secondary prevention of cardiac events 	
Dosing	<ul style="list-style-type: none"> HoFH: 150 mg SC q2 weeks HLD or secondary cardiac prevention: 75 mg SC q2 weeks or 300 mg SC q4 weeks; if adequate LDL response not achieved, may increase to max of 150 mg q2 weeks 	<ul style="list-style-type: none"> HoFH: 420 mg SC q4 weeks; may increase to 420 mg q2 weeks if meaningful response not achieved in 12 weeks HLD or secondary cardiac prevention: 140 mg q2 weeks or 420 mg q4 weeks
Dosage forms	<ul style="list-style-type: none"> Auto-injector 75 mg/mL or 150 mg/mL 	<ul style="list-style-type: none"> Repatha Sure Click (auto-injector) 140 mg/mL Repatha Pushtonex System (single use infusor with pre-filled cartridge) 420 mg/3.5 mL – administered over 9 minutes
Storage	<ul style="list-style-type: none"> Store in refrigerator in outer carton until used Once used, keep at room temperature, use within 30 days 	
Injection clinical pearls	<ul style="list-style-type: none"> Do not shake or warm with water Administer by SC injection into thigh, abdomen, or upper arm Rotate injection site with each injection 	
Drug interactions	<ul style="list-style-type: none"> No known significant interactions 	
Monitoring parameters	<ul style="list-style-type: none"> Lipid panel before initiating therapy, 4-12 weeks after initiating, and q3-12 months thereafter 	
Side effects	<ul style="list-style-type: none"> Injection site reaction (4-17%) Hypersensitivity reaction (9%) Influenza (6%) Myalgia (4-6%) Diarrhea (5%) 	<ul style="list-style-type: none"> Nasopharyngitis (6-11%) Upper respiratory tract infection (9%) Diabetes mellitus (9%) Influenza (8-9%) Injection site reaction (6%) Myalgia (4%)

Adenosine Triphosphate-citrate Lyase - ACL Inhibitor

Indicated for adults with heterozygous familial hypercholesterolemia or established ASCVD who require additional LDL lowering. Use with maximally tolerated statins for further LDL reduction. Available in a combination pill with ezetimibe. The effects on CV morbidity and mortality have not yet been determined.

Class / Action	Generic / Trade Name	Daily Dose	Frequency	Considerations
Lowers LDL cholesterol by inhibiting production in the liver.	Bempedoic acid / Nexletol	180 mg	Once daily	May increase uric acid levels- use caution in gout. May take with or without food. No dose adjustment for renal or hepatic impairment.
	Bempedoic acid/ezetimibe (Nexlizet)	180 mg /10mg		

*indicates medication is available in generic form.

Page 5

Diabetes Education Services ©2024 - www.DiabetesEd.net For educational purposes only, see package insert for prescribing info.