

## Effective

Action		Insulin Name	Onset	Peak	Duration	Considerations
Bolus	Very Rapid Acting Analogs	Aspart (Fiasp)	16 - 20 min	1 - 3 hrs	5 - 7 hrs	Bolus insulin lowers after-meal glucose. Post meal BG reflects efficacy.  Basal insulin controls BG between meals and nighttime. Fasting BG reflects efficacy.
		Lispro-aabc (Lyumjev)	15 - 17 min	2 - 3 hrs	5 - 7 hrs	
	Rapid Acting Analogs	Aspart (Novolog)	20 - 30 min	1 - 3 hrs	3 - 7 hrs	
		Lispro (Humalog*/ Admelog)	30 min	2 - 3 hrs	5 - 7 hrs	
		Glulisine (Apidra)	15 - 30 min	1 - 3 hrs	3 - 4 hrs	
Short Acting	Regular*	30 - 60 min	2 - 4 hrs	5 - 8 hrs		
Basal	Intermediate	NPH	2 - 4 hrs	4 - 10 hrs	10 - 16 hrs	Side effects: hypoglycemia, weight gain.  Typical dosing range: 0.5-1.0 units/kg body wt/day.
	Long Acting	Glargine (Lantus*/Basaglar/Semglee/Rezvoglar)	2 - 4 hrs	No Peak	20 - 24 hrs	
		Degludec (Tresiba)*	~ 1 hr		< 42 hrs	
Basal + Bolus	Intermediate + short	Combo of NPH + Reg 70/30 = 70% NPH + 30% Reg 50/50 = 50% NPH + 50% Reg	30 - 60 min	Dual peaks	10 - 16 hrs	Discard most open vials after 28 days. For pen storage guidelines, see package insert.
	Intermediate + rapid	Novolog® Mix - 70/30 Humalog® Mix - 75/25 or 50/50	5 - 15 min		24 hrs	

# Concentrated & Inhaled Insulins

Name/Concentration	Insulin/Action	Considerations
Humulin Regular U-500 <ul style="list-style-type: none"> <li>500 units insulin/mL</li> <li>KwikPen or Vial</li> </ul>	Regular Bolus / Basal	Indicated for those taking 200+ units daily. 3 mL pen holds 1,500 units. Max dose 300 units. Once opened, good for 28 days. 20 mL vial holds 10,000 units. Max dose 250 units using U-500 syringe. Once opened, good for 40 days.
Humalog KwikPen U-200 200 units insulin/mL.	Lispro (Humalog) Bolus	3 mL pen holds 600 units. Max dose 60 units. Once opened good for 28 days.
Lyumjev KwikPen U-200 200 units insulin/mL.	Lispro (Lyumjev) Bolus	3 mL pen holds 600 units. Max dose 60 units. Once opened good for 28 days.
Toujeo Solostar U-300 Pen 300 units insulin/mL.	Glargine (Lantus) Basal	1.5 mL pen holds 450 units. Max dose 80 units. 3 mL Max Solostar pen holds 900 units. Max dose 160 units. Once opened good for 56
Tresiba FlexTouch U-200 Pen	Degludec (Tresiba)	3 mL pen holds 600 units. Max dose 160 units. Once opened good for 56 days.

**All concentrated insulin pens and the U-500 syringe automatically deliver correct dose (in less volume). No conversion, calculation or adjustments required. For example,** if order reads 30 units, dial the concentrated pen to 30 units or draw up 30 units on the U-500 syringe. **Important – never withdraw concentrated insulin from the pen using a syringe.**

## Inhaled Insulins

Action	Insulin Name	Dose Range	Onset	Peak	Duration	Considerations
Bolus – Rapid-acting	Afrezza Inhaled regular human insulin	4, 8, and 12 unit cartridges before meals	~ 12 min	35 - 45 mins	1.5 - 3 hrs	Assess lung function. Avoid in lung disease — bronchospasm risk. Side effects: hypo, cough, throat irritation.