



Diabetes Essentials: Building Blocks of Person-Centered Care

Beverly Thomassian, RN, MPH, BC-ADM, CDCES

Pronouns: She, her and hers

Founder - www.DiabetesEd.net


Coach Bev has no Conflict of Interest

- ▶ She's not on any speaker's bureau
- ▶ Does not invest or have any financial relationships with diabetes related companies.
- ▶ Gathers information from reading package inserts, research and articles
- ▶ The ADA Standards of Medical Care is main resource for course content

Diabetes Essentials: Building Blocks of Person-Centered Care

Objectives

- ▶ Identify the key elements of the standards that improve clinical care for people with diabetes.
- ▶ Review and discuss appropriate use of the latest medications and that address hyperglycemia and cardiorenal health.
- ▶ Describe strategies to incorporate lifestyle changes into diabetes self-management.
- ▶ Share practical approaches to assess and address diabetes distress in clinical care.



CDC Announces



35% of
Americans will
have Diabetes
by 2050

Boyle, Thompson, Barker, Williamson

2010, Oct 22:8(1)29

www.pophealthmetrics.com

Poll Question 1

- ▶ What percent of total people in the U.S. are living with undiagnosed and diagnosed type 2 diabetes?
- ▶ A. About 30%
- ▶ B. 11.3%
- ▶ C. 16.8%
- ▶ D. 25.6%




Type 2 Diabetes in America 2025

- ▶ 16.8% with Diabetes
 - ▶ 11% don't know they have it
- ▶ 38% with Prediabetes – 97 million adults

Figure 3. Age-adjusted, county-level prevalence of diagnosed diabetes among adults aged 20 years or older, United States, 2004, 2012, and 2019

<https://www.cdc.gov/nchs/data/databriefs/db516.pdf>




Data sources: US Diabetes Surveillance System; Behavioral Risk Factor Surveillance System.

NCHS Data Brief ■ No. 516 ■ November 2024

Diabetes Prevalence by Ethnic Group

- ▶ For adults, diabetes prevalence highest among:
 - American Indians and Alaska Natives (14.5%),
 - Non-Hispanic Blacks (12.1%),
 - People of Hispanic origin (11.8%),
 - Non-Hispanic Asians (9.5%)


Figure 2. Age-adjusted estimated prevalence of diagnosed diabetes by race/ethnicity group and sex for adults aged 18 years or older, United States, 2018–2019



Lived Experiences & Advocacy



Equality vs Equity



© 2017 Robert Wood Johnson Foundation


Design and deliver diabetes care with goal of **health equity** across all populations.

<https://coveragetoolkit.org/health-equity/defining-health-equity/>

Address Barriers to Self Management

- **Barriers exist** within health system, payer, health care professional & individual.
- **Address barriers** through innovation, including community health workers, telehealth, other digital health solutions.
- **Consider social determinants of health** in the target population when designing care.

What Goes Into Your Health?



Source: Institute for Clinical Systems Improvement, Going Beyond Clinical Walls: Solving Complex Problems (October 2014)

<https://coveragetoolkit.org/health-equity/defining-health-equity/>

Social Determinants of Health


- ▶ SDOH are defined as the economic, environmental, political, and social conditions in which people live and are responsible for a major part of health inequality worldwide.



1. Improving Care and Promoting Health in Populations:
Standards of Care in Diabetes—2025 **FREE**
American Diabetes Association Professional Practice Committee

Greater exposure to adverse SDOH over the life course results in poor health. Use quality data to identify inequities & take action.

Now, let's get to the Nitty Gritty



- Beta – insulin - 60%
- Alpha – glucagon 30%
- Delta –somatostatin 10%

LIVER

SMALL
INTESTINE

PANCREAS

DUCT

ENZYME-PRODUCING
CELL


INSULIN-
PRODUCING
CELL

WHITE
BLOOD CELL

ISLET OF
LANGERHANS

BLOOD
VESSEL

1 Surgeons remove the pancreas
from a human cadaver.



Hormones Effect on Glucose

<u>Hormone</u>	<u>Effect</u>
▶ Glucagon (pancreas)	↑
▶ Stress hormones (kidney)	↑
▶ Epinephrine (kidney)	↑
▶ Insulin (pancreas)	↓
▶ Amylin (pancreas)	↓
▶ Gut hormones - incretins (GLP-1) released by L cells of intestinal mucosa, beta cell has receptors)	↓

Pre Diabetes & Type 2- Screening Guidelines (ADA 2025 Clinical Practice Guidelines)

1. Start screening all people at age 35.
2. Screen at any age if BMI ≥ 25 (Asians BMI ≥ 23) plus one or $>$ additional **risk factor**:

- ▶ First-degree relative w/ diabetes
- ▶ Member of a high-risk ethnic population
- ▶ Habitual physical inactivity
- ▶ History of heart disease
- ▶ Check more frequently if taking high risk meds; antiretrovirals, 2nd generation antipsychotics or steroids, thiazide diuretics, statins
- ▶ History of pancreatitis, prediabetes, GDM, periodontitis



Diabetes 2 - Who is at Risk?

(ADA 2024 Clinical Practice Guidelines)



Screen using A1C, Fasting Blood Glucose or OGTT.

Repeat screening at least every 3 years if negative.

*If prediabetes or on high risk meds, recheck yearly

Risk factors cont'd

- ▶ HTN - BP > 130/80
- ▶ HDL < 35 or triglycerides > 250
- ▶ History of Gestational Diabetes Mellitus
- ▶ Polycystic ovary syndrome (PCOS)
- ▶ Other conditions associated w/ insulin resistance:
 - ▶ Elevated BMI, acanthosis nigricans (AN)

Diabetes Screening Guidelines

(ADA 2025 Clinical Practice Guidelines – Cheat Sheet)

RECOMMENDATIONS FOR DIAGNOSIS AND CLASSIFICATION OF DIABETES – 2025

CRITERIA FOR SCREENING FOR DIABETES AND PREDIABETES IN ASYMPTOMATIC ADULTS – TABLE 1

DIABETES TYPE	RISK FACTORS and FREQUENCY OF SCREENING and TESTING FOR DIABETES
<i>Type 1</i>	Screen those at risk for presymptomatic type 1 diabetes, by testing autoantibodies to insulin, GAD, islet antigen 2 or ZnT8. Also test antibodies for those with type 1 phenotypic risk (younger age, weight loss, ketoacidosis , etc.)
<ol style="list-style-type: none"> Test all adults starting at age 35 for prediabetes and diabetes using Fasting Plasma Glucose, A1C or OGTT. Perform risk-based screening if BMI ≥ 25 or BMI ≥ 23 in Asian Americans 10yrs+ with 1 or more risk factors: <ul style="list-style-type: none"> History of cardiovascular disease Physical inactivity First or second degree relative with diabetes HDL ≤ 35 mg/dl or triglyceride ≥ 250 mg/dl High risk ethnicity or ancestry Hypertension $\geq 130/80$ or on therapy for HTN Other conditions associated with insulin resistance (PCOS, Acanthosis Nigricans, Steatosis) If results normal, repeat test at a minimum of 3-year intervals or more frequently based on risk status. Test Yearly if A1C $\geq 5.7\%$ or Impaired Fasting Glucose or History of GDM (test at least every 1- 3 years) <p>Closely monitor high-risk groups (before taking 2nd generation antipsychotics, steroids, thiazide diuretics, statins, HIV meds <i>and</i> after initiating therapy) with history of pancreatitis, or periodontal disease.</p>	


2. Diagnosis and Classification of Diabetes: Standards of Care in Diabetes—2025 FREE
American Diabetes Association Professional Practice Committee

Poll Question 1

- ▶ Which of the following level is considered pre-diabetes range?
 - a. Fasting BG of 62
 - b. A1c of 5.9 %
 - c. After meal BG of 137
 - d. A1c of 7.1 %



Natural History of Diabetes




Healthy

FBG <100

Random <140

A1c <5.7%

Yes!



Prediabetes

FBG 100-125

Random 140 - 199

A1c ~ 5.7- 6.4%

**~ 50% working
pancreas**

NO



Diabetes

FBG 126 +

Random 200 +

A1c 6.5% or +

**~ 20% working
pancreas**

Development of type 2 diabetes happens over years or decades

PreDiabetes is FREAKING ME OUT

- ▶ 96 million people in US
- ▶ 80% don't know they have it
- ▶ In 3-5 years, about 30% of predm will get diabetes
- ▶ Associated with higher rates of heart attack, stroke, neuropathy and vessel disease



Do I look like I am freaking out?

3. Prevention or Delay of Diabetes and Associated Comorbidities:
Standards of Care in Diabetes—2025 **FREE**

American Diabetes Association Professional Practice Committee


Poll Question 2

- ▶ What best describes prediabetes in the U.S.?
 - a. Prediabetes affects 18-20% of people above the age of 20.
 - b. The prevalence of prediabetes and diabetes are almost equal.
 - c. Most people with BMI of 30 or greater have prediabetes.
 - d. Prediabetes is associated with increased risk of CV disease




3. Detecting PreDiabetes Matters

- ▶ Given the cost-effectiveness of lifestyle behavior modification programs for diabetes prevention:
 - ▶ Offer diabetes prevention programs to adults at high risk of type 2 diabetes
 - ▶ Prescribe effective eating patterns
 - ▶ Address inconsistencies in access – leverage technology
- ▶ Screening guidelines for people with Type 1



Get About 7 Hours of Quality Sleep to Prevent Diabetes

- ▶ Poor sleep quality was associated with a 40–84% increased risk of developing type 2 diabetes in a meta-analysis.
- ▶ Chronotype preference has been linked with many chronic diseases, including type 2 diabetes.
- ▶ Night owls have 2.5 higher odds ratio for diabetes risk (i.e., going to bed late and getting up late) than early birds.
 - ▶ Independent of sleep duration and sleep sufficiency



3. Prevention or Delay of Diabetes and Associated Comorbidities:
Standards of Care in Diabetes—2025 **FREE**

American Diabetes Association Professional Practice Committee

The composition of the gut microbiome may also affect the likelihood of developing type 2 diabetes.

3. Interventions for Prediabetes

- ▶ Use more intensive approach for high-risk individuals:
 - ▶ BMI of 35+
 - ▶ If A1C is ~6.0 or FPG is 110
 - ▶ History of GDM
- ▶ No FDA approved med for prevention (off label)
- ▶ Consider Metformin Therapy for Prediabetes
 - ▶ Monitor B12 level (esp with neuropathy or anemia)

- ▶ CV Risk Mitigation important.
- ▶ Statin can increase BG, stop if notice elevation
- ▶ Consider low dose pioglitazone (Actos) if history of stroke.



Common Oral Diabetes Meds



Class/Main Action	Name(s)	Daily Dose Range	Considerations
Biguanides <ul style="list-style-type: none"> Decreases hepatic glucose output First line med at diagnosis of type 2 	metformin (Glucophage)	500 - 2550 mg (usually BID w/ meal)	Side effects: nausea, bloating, diarrhea, B12 deficiency. To minimize GI Side effects, use XR and take w/ meals. Obtain GFR before starting. <ul style="list-style-type: none"> If GFR <30, do not use. If GFR <45, don't start Meformin If pt on Metformin and GFR falls to 30-45, eval risk vs. benefit; consider decreasing dose. For dye study, if GFR <60, liver disease, alcoholism or heart failure, restart metformin after 48 hours if renal function stable. Benefits: lowers cholesterol, no hypo or weight gain, cheap. Approved for pediatrics, 10 yrs + Lowers A1c 1.0%-2.0%.
	Riomet (liquid metformin)	500 - 2550 mg 500mg/5mL	
	Extended Release-XR (Glucophage XR) (Glumetza) (Fortamet)	(1x daily w/dinner) 500 – 2000 mg 500 – 2000 mg 500 – 2500 mg	

Biguanide derived from:
 Goat's Rue *Galega officinalis*,
 French Lilac
 Does NOT harm kidneys
 \$10 for 3-month supply from
 Walmart & other pharmacies

GOAT'S RUE
(GALEGA OFFICINALIS)

Used for

- Diabetes

Potential uses

- Cancer
- Ovarian cysts

Uses under investigation

- Parkinson's
- Neuron growth



Indications for Insulin Sensitizers

Pioglitazone (Actos)

- ▶ **Action:** decrease insulin resistance by making muscle and adipose cells more sensitive to insulin. Decrease free fatty acids
- ▶ **Names:**
 - ▶ pioglitazone (Actos) – bladder cancer warning
 - ▶ Dosing: 15-45 mg daily
 - ▶ Consider adding low dose if history of stroke or have steatosis
 - ▶ rosiglitazone Dosing: 4-8 mg daily

Class/Main Action	Name(s)	Daily Dose Range	Considerations
Thiazolidinediones "TZDs" • Increases insulin sensitivity	pioglitazone (Actos) rosiglitazone	15 – 45 mg daily 4 – 8 mg daily	Black Box Warning: TZDs may cause or worsen CHF. Monitor for edema and weight gain. Increased peripheral fracture risk. Actos may increase risk of bladder cancer. Lowers A1c 0.5% – 1.0%

- ▶ **Efficacy/ Considerations**
 - ▶ Reduce A1C ~0.5-1.0%
 - ▶ 6 weeks for maximum effect
 - ▶ Actos \$5 a month
 - ▶ Can cause fluid retention, not indicated w/ CHF



Poll question 3

- ▶ JR is started on Metformin 500mg BID. Which of the following is true?
- a. Hold metformin if blood glucose below 90 mg/dl.
 - b. Evaluate B12 levels before starting medication.
 - c. Metformin is considered weight neutral
 - d. Metformin can cause kidney damage, so increase fluid intake



Medication Taking Behaviors

- ▶ Adequate medication taking is defined as 80%
- ▶ 23% of time, if A1c, B/P, lipids above target - due to med taking behavior
- ▶ Assess for barriers
- ▶ If taking meds 80% of time and goals not met, consider medication intensification



Barriers include:

Forgetting to fill Rx, forgetting to take, fear, depression, health beliefs, med complexity, cost, knowledge gap, system factors, etc.

**Work on targeted approach
for specific barrier**

Diabetes Admit for Hyperglycemia

- ▶ JR is admitted for hyperglycemia because he stopped taking his diabetes meds.
- ▶ HCP says, “Don’t you realize you are going to get complications, like kidney disease or amputation if you don’t take your medications?”
- ▶ Door Closed – No Connection made

How Does JR Feel?

- ▶ Embarrassed
- ▶ Ashamed
- ▶ Defeated
- ▶ Angry
- ▶ Unheard



How does HCP feel?

- ▶ Frustrated
- ▶ Defeated
- ▶ Worried

Diabetes Visit – Let's Go *through*

A small adjustment can make a **BIG** Difference


- ▶ HCP says, “JR, I am worried about you and your elevated blood glucose. Can you share what is going on in your life?”
- ▶ Door Open – Connection made

How Does JR Feel?

- ▶ Heard & Seen
- ▶ Recognized
- ▶ Connected
- ▶ Engaged

How does the HCP feel?

- ▶ Connected
- ▶ Concerned
- ▶ Collaborative



Create a Judgement Free Zone – Roll out the Carpet of Acceptance

There are no bad or good blood glucose numbers.

There is no such thing as cheating.
You are not failing at your diabetes.
It is not your fault you have diabetes.
Thank you for showing up today.



5 M's and Judgement Free Zone



Mood



Meals



Movement




Medicines




Minutes

Let's meet people where they are at.



Type 1 ~ Immune Mediated 5-10% of Diabetes



Screening is offered at no cost to eligible individuals to evaluate their personal risk of developi... See more

An infographic with a blue background and a large white question mark. On the left, there is a photo of a smiling woman. The text reads: 'DID YOU KNOW? The risk for people in the general population (no T1D family history) is about 1 in 300. For those who have a family member with T1D, the risk is 1 in 20.' The Type 1 Diabetes TrialNet logo is in the top right corner.

DID YOU KNOW

The risk for people in the general population (no T1D family history) is about 1 in 300. For those who have a family member with T1D, the risk is 1 in 20.



1.5 Million people have type 1 in U.S.

Prevalence increasing:

2001 – 1.48 per 1000 youths diagnosed with diabetes

2017 - 2.15 per 1000 youths diagnosed with diabetes

Incidence & Prevalence increasing

Highest incidence in Finland or Northern Europe.

ADCES In Practice - March 2024

Recent Advances in Type 1 Diabetes: Teplizumab (Tzeild®)

Karen S. Fiano, PHARMD, BCACP, Devada Singh-Franco, PHARMD, CDCES, Young M. Kwon, BS, PHD

Type 1 – 10% of all Diabetes

- Auto-immune pancreatic beta cells destruction
- Most commonly expressed at age 10 - 14
- Insulin sensitive (require 0.5 - 1.0 units/kg/day)
- Expression due to a combo of genes and environment:
 - Autoimmunity tends to run in families
 - Exposure to virus or other environmental factors
- Signs can include:
 - Increased thirst and hunger
 - Frequent urination or new bed-wetting at hs
 - Unintended weight loss
 - Fatigue and irritability



Type 1 Diabetes Features?



- ▶ For JR, a 28 admitted to the ICU with a blood glucose of 476 mg/dl, pH of 7.1, anion gap of 15. Recently lost 13 pounds.

Type 1 Most Discriminative Features

- Younger than 35 years at diagnosis
- Lower BMI (<25 kg/m²)
- Unintentional weight loss
- Ketoacidosis
- Glucose 360 mg/dl or greater.


Misdiagnosis is common and can occur in ~40% of adults with new type 1 diabetes

Pharmacologic Intervention to Delay Symptomatic Type 1 (in Stage 2)

- ▶ Teplizumab-Tzielid (CD3-monoclonal antibody)
- ▶ 14-day infusion can delay the onset of symptomatic type 1 diabetes (stage 3)
- ▶ An option in selected individuals aged ≥ 8 years with stage 2 type 1 diabetes.
- ▶ In a single trial, 44 individuals received 14-day course of teplizumab vs 32 placebo.
- ▶ The median time to stage 3 diagnosis of type 1
 - ▶ 48.4 months in tep group
 - ▶ 24.4 months placebo
- ▶ Cost: \$193,000
- ▶ Provention Bio has financial assist programs.

Type 1 (stage 2) Delayed with Teplizumab by 2 years www.DiabetesTrialNet.org

► How to get families linked to screening?



The screenshot shows the homepage of the Type 1 Diabetes TrialNet website. At the top left is the logo for Type 1 Diabetes TrialNet. To the right are navigation links for Researchers, Publications, Contact Us, FAQs, and Terminology, along with a search bar. Below the navigation is a horizontal menu with links for Our Research, T1D Facts, Participate, Our Families, TrialNet Locations, About Us, News & Events, and COVID-19. The main content area features a large grid of photos showing diverse families and individuals. In the center of this grid is a white box with the text: "Join the TrialNet #T1Dfamily" in blue and red, followed by "Detect future risk of T1D and advance important research!". Below the grid, on the left, is the heading "Imagine a future without type 1 diabetes" and a paragraph of text describing the organization's mission. On the right, under the heading "GET STARTED", are two buttons: "Signup to be screened!" and "Find a location near me".

Type 1 Diabetes TrialNet

English Español

Researchers Publications Contact Us FAQs Terminology Search

Our Research T1D Facts Participate Our Families TrialNet Locations About Us News & Events COVID-19

Join the TrialNet #T1Dfamily
Detect future risk of T1D and advance important research!

Imagine a future without type 1 diabetes

TrialNet is an international network of leading academic institutions, endocrinologists, physicians, scientists and healthcare teams at the forefront of type 1 diabetes (T1D) research. We offer risk screening for relatives of people with T1D and innovative clinical studies testing ways to slow down and prevent disease progression. Our goal: a future without T1D!

GET STARTED


Signup to be screened!

Find a location near me

Miracle of Insulin



Patient J.L., December 15, 1922



February 15, 1923



Patti LaBelle

"divabetic"

"I have diabetes, it
doesn't have me"




Signs of Diabetes

- ▶ Polyuria
- ▶ Polydipsia
- ▶ Polyphasia
- ▶ Weight loss
- ▶ Fatigue
- ▶ Skin and other infections
- ▶ Blurry vision



Visceral Fat and Subcutaneous Fat



What is Type 2 Diabetes?

► Complex metabolic disorder

(Insulin resistance and deficiency)

with social, behavioral and environmental risk factors unmasking the effects of genetic susceptibility.

New Diagnosis?

Call 800 – DIABETES to request

“Getting Started Kit”

www.Diabetes.org



Life Study – Mrs. Jones

Mrs. Jones is 62 years old, with a BMI of 36 and complains of feeling tired and urinating several times a night. She has an urinary tract infection. Her A1c is 8.3%, glucose 237.

She is hypertensive with a history of gestational diabetes. No ketones in urine.

- ▶ What are her risk factors and signs of diabetes?
- ▶ You find a few moments to teach and she asks you some questions.



Mrs. Jones asks you What Do You Say?

- ▶ What is diabetes?
- ▶ They say I am a diabetic because I am obese?
- ▶ How am I going to control this?
- ▶ What is a normal blood sugar?
- ▶ Do I have to test my blood sugars?
- ▶ My doctor told me to stay away from white foods. Is that true?




Mrs. Jones asks you What Do You Say?

- ▶ You are wondering if your weight caused your diabetes?
- ▶ You can manage your diabetes and improve your health at the same time.
- ▶ For people without diabetes, fasting blood sugar is less than 100 and A1c is less than 5.7%
- ▶ Checking blood sugars can help you figure out if the plan is working.




Look Beyond Diabetes


- ▶ ACE – Adverse Childhood Experiences
- ▶ Feelings around their diabetes
- ▶ Cultural traditions, family system.
- ▶ Social, religious and employment influences
- ▶ Personal factors: attitudes, cognitive factors, literacy, learning styles, health beliefs
- ▶ Social Determinants of health



The Noxious Nine – Pathophysiology T2D



Signs of Hypercortisolism



▶ Other signs

- ▶ muscle weakness,
- ▶ high blood pressure,
- ▶ diabetes,
- ▶ excessive hair growth,
- ▶ acne,
- ▶ bone loss and
- ▶ mood changes like irritability and depression.

Hypothalamus → CRH → Pituitary → ACTH → Adrenal → Cortisol

- ▶ Cortisol effects: gluconeogenesis, insulin resistance, fat redistribution

SGLT2 Inhibitors- “Glucoretics”

- ▶ **Action:** decreases renal reabsorption of glucose proximal tubule of kidneys (reset renal threshold)
- ▶ **Preferred** diabetes treatment for people with heart and kidney failure. Decreases BG & CV Risk.
- ▶ **AWP:** ~\$650 a month



Class/Main Action	Name(s)	Daily Dose Range	Considerations
SGLT2 Inhibitors “Glucoretic” • Decreases glucose reabsorption in kidneys	Canagliflozin* (Invokana) Dapagliflozin*† (Farxiga) Empagliflozin*† (Jardiance) Ertugliflozin (Steglatro) Bexagliflozin (Brenzavvy)	100 - 300 mg 1x daily 5 - 10 mg 1x daily 10 - 25 mg 1x daily 5 – 15 mg 1x daily 20 mg 1x daily	Side effects: hypotension, UTIs, genital infections, increased urination, weight loss, ketoacidosis. Heart Failure, CV & Kidney Protection: 1st line therapy for Heart Failure (HF), Kidney Disease (CKD), Cardiovascular Disease, before or with metformin Considerations: If GFR ≥ 20, use SGLT-2 to reduce CVD, Heart Failure and Chronic Kidney Disease. Limited BG lowering effect if GFR <45. See package insert for GFR cut-offs and dosing. Benefits: SGLT-2s* reduce BG, CV death & HF, slow CKD. †Approved for peds, 10 yrs +. Lowers A1C 0.6% to 1.5%.

SGLT-2i Indications Summary

Drug	Lower BG	Reduce CV Risk?	Use to treat Heart Failure?	Slow renal disease?
Dapagliflozin (Farxiga)	Yes	Yes	Yes +/- Diabetes	Yes
Empagliflozin (Jardiance)	Yes	Yes	Yes +/- Diabetes	Yes
Canagliflozin (Invokana)	Yes	Yes	Yes w/ Diabetes	Yes
Ertugliflozin (Steglatro)	Yes	No	Yes w/ Diabetes	Yes
Bexagliflozin (Brenzavvy)	Yes	NA	NA	NA

“Getting diabetes saved my life.”

~ Sherri Sheperd

PLAN

D

How to

**LOSE WEIGHT
AND BEAT**

DIABETES

(EVEN IF YOU DON'T HAVE IT)

**SHERRI
SHEPHERD**

Emmy Award-Winning Cohost of *The View*

WITH BILLIE FITZPATRICK


READ BY THE AUTHOR



Sherri Shepard decided to embrace diabetes and use it as a motivator to improve her health.


Other Specific Types of DM

- ▶ Medications such as:
steroids, protease inhibitors
and Prograf®
- ▶ Secondary to Agent Orange
- ▶ Liver failure
- ▶ TPN or tube feedings
- ▶ **Diabetes Type 3c**
 - ▶ Cystic fibrosis, **pancreatitis**
 - ▶ Pancreatic cancers or
removal
 - ▶ Hemochromatosis



Pancreatitis

- ▶ People with diabetes 2xs risk of acute pancreatitis
- ▶ After episode of pancreatitis, one third of people will get prediabetes or diabetes
 - ▶ About 25% to 80% of people with chronic pancreatitis develop Type 3c diabetes.
- ▶ Pancreatitis is an exocrine dysfunction:
 - ▶ Disrupts global architecture or physiology of pancreas
 - ▶ Results in both exocrine and endocrine dysfunction.



2. Diagnosis and Classification of Diabetes: Standards of Care in Diabetes—2025 FREE
American Diabetes Association Professional Practice Committee

Regardless of the cause, hyperglycemia needs to be treated.



DiaBingo

- ~~B Frequent skin and yeast infections~~
- B A BMI of _____ or greater indicates increased pre/diabetes risk?
- B To reduce complications, control **A1c**, **B**lood pressure, **C**holesterol
- B PreDiabetes – fasting glucose level of ____ to _____
- B Erectile dysfunction indicates greater risk for _____
- B Diabetes – fasting glucose level _____ or greater
- B Type 1 diabetes is best described as an _____ disease
- B People with diabetes are _____ times more likely to die of heart dx
- B Elevated triglycerides, < HDL, smaller dense LDL
- B Each percentage point of A1C = _____ mg/dl glucose
- B At dx of type 2, about __% of the beta cell function is lost
- B Diabetes – random glucose _____ or greater

Sulfonylureas - Secretagogues or “Squirters”

- ▶ Mechanism: Stimulate beta cells to release insulin
- ▶ Dosed 1-2x daily before meals
- ▶ Adverse effects
 - ▶ Hypoglycemia, Weight gain, watch renal function
- ▶ Low cost, \$12 for 3 months supply
- ▶ Can help with glucose toxicity, lowers A1C 1-2%



Sulfonylureas • Stimulates sustained insulin release	glyburide: (Diabeta) (Glynase PresTabs)	1.25 – 20 mg 0.75 – 12 mg	Can take once or twice daily before meals. Low cost generic. Side effects: hypoglycemia and weight gain. Eliminated via kidney. Caution: Glyburide most likely to cause hypoglycemia. Lowers A1c 1.0% – 2.0%.
	glipizide: (Glucotrol) (Glucotrol XL)	2.5 – 40 mg 2.5 – 20 mg	
	glimepiride (Amaryl)	1.0 – 8 mg	

Reducing Hypoglycemia

▶ Which are the only diabetes meds that directly cause hypoglycemia?



- ❑ Insulin
- ❑ Secretagogues
(sulfonylureas, glitinides)

Hypoglycemia – A Big Deal

Hypoglycemia (Low Blood Glucose)

Some Symptoms:

Causes: Too little food or skipping a meal; too much insulin or diabetes pills; more active than usual.

Onset: Often sudden.



SHAKY



FAST
HEARTBEAT



SWEATING



DIZZY



ANXIOUS



HUNGRY



BLURRY VISION



WEAKNESS OR FATIGUE



HEADACHE



IRRITABLE

Hypoglycemia (Glucose) Alert Values

- ▶ **BG <70mg/dl – Level 1**
- ▶ Follow 15/15 rule and contact provider to make needed changes. At increased hypo risk.
- ▶ **BG < 54mg/dl – Level 2**
- ▶ Indicates serious hypo. Reassess BG Goals. Consider med decrease. Predictive of Level 3 Hypo. Needs Glucagon Emergency Kit
- ▶ **Severe Hypoglycemia – Level 3**
- ▶ Altered mental, physical functioning.
- ▶ Requires external assistance – no threshold



Hypoglycemia: Identify, Treat, & Prevent

PocketCards are updated twice yearly.
Scan QR code to download or
order the latest version.



Step 1

Identify your signs of hypoglycemia or low blood sugar:

- Sweaty
- Shaky
- Hungry
- Can't think straight
- Headache
- Irritated, grouchy
- Other



Step 2

If have signs of hypo, treat with carbs until glucose reaches 70+, then eat usual meal.

- Sugary drink, 4–8oz
- Piece of fruit
- Raisins, handful
- Glucose tabs, 4+
- Honey/glucose gel
- Skittles candy, 15+



Step 3

Have glucagon rescue meds available.

In case of severe hypo, identify someone (ahead of time) who can get medical help & give a glucagon rescue medication.

Notify your provider of low blood sugar events.

Hypoglycemia Levels:

Level 1 – Glucose less than 70

Level 2 – Glucose less than 54

Level 3 - Severe, needs assistance


Identify Causes of Hypo & Problem Solve to Prevent Future Episodes

- » Low carb meal
- » Extra activity
- » Drinking alcohol

- » Delayed, missed meal
- » Too much insulin/meds
- » Insulin timing

AGP Report: Continuous Glucose Monitoring

Time in Ranges Goals for Type 1 and Type 2 Diabetes



Test Patient DOB: Jan 1, 1970

14 Days: August 8-August 21, 2021

Time CGM Active: 100%

Glucose Metrics

Average Glucose..... **175 mg/dL**

Goal: <154 mg/dL

Glucose Management Indicator (GMI)..... **7.5%**


Goal: <7%

Glucose Variability..... **45.5%**

Defined as percent coefficient of variation
Goal: <36%

Ambulatory Glucose Profile (AGP)

AGP is a summary of glucose values from the report period, with median (50%) and other percentiles shown as if they occurred in a single day.



Glucagon Rescue Medications for Diabetes-Related Hypoglycemia

Name/Delivery	Supplied	Dose Range		Age / Route / Storage
		Adult	Peds / Age WT Dosing	
Glucagon Emergency Kit Injection requires mixing glucagon powder	1mg / 1mL vial + syringe	1mg	0.03mg/kg or < 6yrs or < 25 kgs 0.5mg ≥ 6yrs or > 25kgs 1mg	All ages approved SubQ or IM admin Expires in 2 years at room temp.
Baqsimi Nasal glucagon powder	3 mg intranasal device	3 mg	< 4 yrs: not recommended 4 yrs or older 3mg dose	Approved Age 4+ Nasal admin Expires ~ 2 years at room temp (keep in shrink-wrapped tube).
Gvoke Injectable liquid stable glucagon solution	0.5mg or 1.0mg in -Prefilled syringe -HypoPen auto-injector -Kit with vial and syringe	1 mg	< 2yrs: not recommended 2- 12 yrs < 45kg 0.5mg ≥ 45kg 1mg 12 yrs or older 1mg	Approved Age 2+ SubQ admin in arm, thigh, abdomen Expires in 2 years at room temp (keep in foil pouch).
Dasiglucagon (Zegalogue) Stable liquid glucagon analog	0.6mg/0.6mL Prefilled syringe Autoinjector	0.6mg	< 6yrs: not recommended 6 yrs or older 0.6mg	Approved Age 6+ SubQ in abdomen, buttocks, thigh outer upper arm Expires in 1 year at room temp. (store in red protective case).

***All raise BG 20+ points. Can cause nausea, vomiting. After admin, roll person on side. Seek medical help. If no response after 1st dose, give 2nd dose in 15 mins. When awake, give oral carbs ASAP when safe to swallow. Please consult package insert for detailed info.**

All PocketCard content is for educational purposes only. Please consult prescribing information for detailed guidelines.



Poll Question 4



- ▶ JL is 78 and drinks a “few cocktails” every night. Lives with partner and takes basal insulin at night and bolus insulin as needed. **Has had a few low blood glucose levels in past week of 62, 49 and 51.** What is the most important recommendation?
- ▶ A. Decrease alcohol intake
- ▶ B. Check BG at least 4 times a day.
- ▶ C. Double check injection sites.
- ▶ D. Get glucagon rescue medication.



Weight is a Heavy Issue



Health Behavior Change: Shifting Focus

Health at Every Size (HAES) Principles

- ▶ Weight Inclusivity
- ▶ Health Enhancement
- ▶ Eating for Well-being
- ▶ Respectful Care
- ▶ Life-Enhancing Movement

“Lots of people are fat and fit—many avid dancers, runners, lifters, and sports team members are big to start with and stay that way. They tend to be far healthier than thin people who don't move around much or eat a nutritious mix of foods.”

"People might think they can tell who's fit and who's not by looking at them, but in fact, it's trickier than that."

Health at Every Size: The Surprising Truth About Your Weight. Bacon holds a Ph.D. in physiology with a focus on nutrition and weight regulation.

"Health at Every Size is about taking care of your body without worrying about whether you're 'too' big or small."

Get a Tape Measure & Other Assessments



- ▶ WHO defines Obesity as: *abnormal or excessive fat accumulation that presents a risk to health*
- ▶ BMI poor indicator for "excessive fat" and health risk

- Overall - assess individual's
 - adipose tissue mass
 - using waist circumference
 - 35" woman, 40" man
 - waist-to-hip ratio
 - Waist smaller than hips
 - waist-to-height ratio
 - Waist < half height
 - presence of associated health or well-being consequences: metabolic, physical, or psychological well-being

Medical Nutrition Therapy Works

- ▶ MNT is effective and beneficial to people with diabetes.
- ▶ When delivered by an RDN, MNT is associated with A1C absolute decreases of
 - ▶ 1.0–1.9% for people with type 1 diabetes and
 - ▶ 0.3–2.0% for people with type 2 diabetes



Healthy Eating Patterns/Approaches

Eating Patterns:

Total Foods Consumed


- ▶ Mediterranean Diet
- ▶ Plant based eating
- ▶ DASH (Dietary Approaches to Stop Hypertension)
- ▶ Low Carbohydrate

Eating Approach:

Tools for developing an eating pattern

- ▶ Diabetes Plate Method
- ▶ Carbohydrate Counting
- ▶ Individualized behavioral approaches

Use Integrative food-based approach.
“People eat food, not nutrients”.



STANDARDS OF CARE | DECEMBER 09 2024

5. Facilitating Positive Health Behaviors and Well-being to Improve Health Outcomes: Standards of Care in Diabetes—2025 **FREE**

American Diabetes Association Professional Practice Committee

Plant-based, Vegetarian or Vegan Diet Patterns

Plan Your Portions

Portion Guide

- Asparagus
- Broccoli
- Brussels sprouts
- Cabbage (with kale)
- Cauliflower
- Cucumbers
- Eggplant
- Eggplant
- Mushrooms
- Onion
- Peas
- Peppers
- Pumpkin
- Spinach
- Tomatoes
- Zucchini

Water or no-calorie drinks

- Corn
- Custard
- Fruit
- Nuts
- Whole grains
- White rice
- Beans, lentils and peas
- Milk and yogurt
- Cheese
- Eggs
- Nut butter
- Oils
- Tofu
- Tofu

Use a smaller plate. This is a 9-inch plate to help guide you.

9 inches

American Diabetes Association

© 2017 American Diabetes Association

vega

Medical Nutrition Therapy – ADA Macronutrient Distribution

- ***“No one-sized-fits-all eating pattern for individuals with diabetes”***
 - no ideal percent of calories from protein, carbohydrate and fat.
- ▶ **Macronutrient distribution based on *individualized assessment***
- ▶ **Consider personal preferences**
 - ▶ tradition, culture, religion, health beliefs and goals, economics
 - ▶ metabolic goals and comorbidities



Healthcare team members should complement MNT, providing guidance on healthy food choices for the individual and behavioral support

Limit Highly Processed Carbs and Added Sugars

Eat more HIGH Fiber foods:

- Choose High fiber carbs loaded with vitamins, minerals and phytonutrients
- “Power Carbs” include:
 - Beans/Lentils
 - Veggies
 - Whole Fruits
 - Low-fat, low sugar milk/yogurt
 - Whole Grain foods
 - as culturally appropriate



Reading the Food Label

1. Check the **Serving size** first. All the numbers on this label are for a 2/3-cup serving.
2. This package has **8 servings**. If you eat the whole thing, you are eating 8 times the amount of calories, carbs, fat, etc., shown on the label.
3. **Total Carbohydrate** shows you types of carbs in the food, including sugar and fiber.
4. Choose foods with **more fiber, vitamins, and minerals**.
5. Choose foods with **lower calories, saturated fat, sodium, and added sugars**. Avoid *trans* fat.

Nutrition Facts	
8 servings per container	
Serving size	2/3 cup (55g)
Amount per serving	
Calories	230
% Daily Value*	
Total Fat 8g	10%
Saturated Fat 1g	5%
Trans Fat 0g	
Cholesterol 0mg	0%
Sodium 160mg	7%
Total Carbohydrate 37g	13%
Dietary Fiber 4g	14%
Total Sugars 12g	
Includes 10g Added Sugars	20%
Protein 3g	
Vitamin D 2mcg	10%
Calcium 260mg	20%
Iron 8mg	45%
Potassium 235mg	6%

* The % Daily Value (DV) tells you how much a nutrient in a serving of food contributes to a daily diet. 2,000 calories a day is used for general nutrition advice.

Fiber – the New “F” Word

- ▶ Goal: minimum
 - ▶ 14 gms / 1000 calories, ~ 30 gms a day
- ▶ How?
 - ▶ Avoid highly processed foods
 - ▶ Choose > 3 gm fiber per serving
 - ▶ Foods: Whole intact grains, legumes, fruits, veggies, nuts/seeds, avocados
- ▶ Why?
 - ▶ Lower all cause mortality and reduced risk of type 2 diabetes
 - ▶ Increased microbiome diversity

Nutrition Facts

Serving Size 1 cup (236g)
Servings Per Container about 2

Amount Per Serving

Calories 260 Calories from Fat 130

% Daily Value*

Total Fat 14g **22%**

Saturated Fat 5g **25%**

Trans Fat 0g

Cholesterol 35mg **12%**

Sodium 990mg **41%**

Total Carbohydrate 19g **6%**

Dietary Fiber 3g **12%**

Sugars 4g

Protein 15g **29%**

Vitamin A 10% • Vitamin C 0%

Calcium 4% • Iron 8%


*Percent Daily Values are based on a 2,000 calorie diet.



Eating Patterns: Key Nutrition Principles

Until there is more evidence:

- ▶ Emphasize non-starchy vegetables in a rainbow of colors
- ▶ “Power carbs”: fruit, legumes, whole grains, nuts and seeds, lean proteins, low-fat dairy
- ▶ Minimize red meat, added sugars, sugary beverages, refined grains and ultra-processed foods



Any approach should consider:

Individual needs: “health status, personal and cultural preferences, ability to sustain recommendations, food access and nutrition security”

Carbs and Lowering Glucose

- ▶ Reducing carb intake has significant evidence for improved glycemia
- ▶ **Low Carb Definitions**
 - ▶ Very Low = < 26% of kcals
 - ▶ Ketogenic: 20-50 gm carb, also high fat.
 - ▶ Most people consume 44-46% of Cals from carb
- ▶ **HOW:** Focus on key nutrition principles, food quality, and choose minimally processed foods and high fiber foods.



Systematic reviews and RCT found:




- Very low carbohydrate diet effectively reduced A1c at 6 months, less difference beyond 1 year.
- Ketogenic Diet increased LDL and no sig. difference in A1c compared with low-carb Mediterranean diet.

Very Low Carb Meal Plan Not Recommended for:

- ▶ Women who are pregnant or lactating or children
- ▶ People with or at risk for disordered eating
- ▶ People who have kidney disease
- ▶ Avoid ketogenic diets if taking SGLT-2 Inhibitor due to high risk of ketoacidosis
 - ▶ Educate on prevention, signs of DKA, how to measure ketones.



How Nutrients Affect Blood Glucose



If type 1/MDI, may need additional insulin or change in dosing strategy for high fat/protein meals

Carbs affect Post-Meal Glucose

- Starch
- Fruit
- Milk
- Desserts



Poll Question 4

- Based on the Food list for Diabetes: which of the following food choices equals ~ 15 gms of carbohydrate? (multiple)
- A. ½ bagel, 4 oz
 - B. 1 ¼ cup strawberries
 - C. 1 cup of milk
 - D. ½ cup of cooked rice
 - E. 1 oz trail mix with fruit



Estimating Carbohydrate for Meals Example (Not ADA Standard)

	<u>Grams</u>	<u>Servings</u>
Each Meal	45-60 gm	3 - 4
Snacks	15-30 gm	1- 2



Carbs affect Post-Meal Glucose
RDA – at least 130 grams of Carb a day
PWD get about 45% of Cals from carbs

Calories	20%	30%	40%	50%
1500	75 g	113 g	150 g	188 g
2000	100 g	150 g	200 g	250 g


Carb Counting - Starch



1/2 cup
cooked beans



1 small ear of corn or
1/2 cup corn



1 slice bread



1/3 cup
cooked pasta



3/4 cup cold
cereal




1/3 cup
cooked rice




6" tortilla



6" roti



5-6 small crackers




1 small potato



1/2 English muffin

Each Food has:
80 Calories
15 grams carb

Carb Counting- Fruit



1 small fresh fruit



1/2 cup fruit juice



1/2 banana, 4"




1/2 cup unsweetened apple sauce



17 small grapes

Each Food has:
60 Calories
15 grams carb



1 cup melon

1/4 cup dried fruit




2 tbsp raisins




1 1/4 cup strawberries



Carb Counting - Milk



Carb Counting - Sweets



Average American Consumes 17 teaspoons added sugar per day

- ▶ WHO/DGA and AHA – Goals
< ~ 5 - 10 teaspoons/day
- ▶ 1 tsp = 4 gms sugar (15 Cals)
- ▶ 15 cals x 19 teaspoons a day
=
 - ▶ 285 cals a day just from added sugars
- ▶ 12oz. soda = 39 gm carb/10
tsps added sugar



Total Carbohydrate 39g	14%
Total Sugars 39g	
Includes 39g Added Sugars	78%

Reduce Refined Carbs, Added Sugars - ADA

- ▶ To manage wt, reduce CVD risk and fatty liver disease
- ▶ ADA strongly discourages consumption of:
 - ▶ Sugar sweetened beverages
 - ▶ Processed “low-fat” or “non-fat” foods with high amounts of refined grains & added sugar



Sugary and processed foods can displace healthier, more nutrient dense food choices

Water is recommended

Non-Nutritive Sweeteners

- ▶ Use in moderation and short term to reduce overall calorie/carbohydrate.
- ▶ Encourage decrease in both sweetened and non-nutritive sweetened beverages.
- ▶ Emphasize water intake.

How: add lemon, lime, or cucumber slices to water, choose no calorie waters



STANDARDS OF CARE | DECEMBER 09 2024

5. Facilitating Positive Health Behaviors and Well-being to Improve Health Outcomes: Standards of Care in Diabetes—2025 **FREE**

American Diabetes Association Professional Practice Committee

New Consult for Diabetes Ed

- ▶ JR, 63 y/o, BMI 32, Waist circumference: 43", A1c 9.3%, Type 2 diabetes 10+ years. On Metformin 2000mg daily + Glipizide 20mg. Low income and

Strategies?

What is JR doing right?

Plate method?

Increase fruit / veggie intake – canned or frozen

Add in beans, chili, soups (lower sodium)

Include, Yogurt, Nuts, oatmeal, wheat toast

Refer to RD / RDN

- ▶ Dinner – Sandwich or burger
- ▶ Evening – Snacks, nuts, crackers, boiled egg
- ▶ Beverages – Diet sodas, tea, water



Poll Question 5

- ▶ A person with diabetes presents with unexplained weight loss, what are some possible causes?
 - A. Taking less insulin than needed
 - B. Disordered eating
 - C. Finances
 - D. Poor dentition
 - E. All the above



Disordered Eating

- ▶ For people with type 1
 - ▶ insulin omission causing glycosuria to lose weight is the most reported disordered eating behavior.
 - ▶ Have high rates of diabetes distress and fear



- ▶ For people with type 2
 - ▶ bingeing episodes with an accompanying sense of loss of control most reported.
 - ▶ If treated with insulin, intentional omission is also frequently reported.

People with diabetes and diagnosable eating disorders have high rates of other psychiatric disorders

Poll Question 6

- ▶ Which of the following is true about alcohol and diabetes based on ADA Standards?
 - A. Only white wine decreases blood sugars
 - B. Men less than 2 drinks and women less than one drink a day
 - C. Alcohol increases risk of hyperglycemia
 - D. 6 ounces of wine is considered one serving.



Incretins: GLP & GIP Receptor Agonists




GLP-1: glucagon like peptide 1

GIP: glucose-dependent insulinotropic polypeptide

GLP-1 Effects in Humans

Understanding the Natural Role of Incretins



Adapted from Flint A, et al. *J Clin Invest*. 1998;101:515-520
Adapted from Larsson H, et al. *Acta Physiol Scand*. 1997;160:413-422
Adapted from Nauck MA, et al. *Diabetologia*. 1996;39:1546-1553
Adapted from Drucker DJ. *Diabetes*. 1998;47:159-169

**GLP-1 degraded by
DPP-4 w/in minutes**

Pocket Card: GLP-1 & GIP RA

GLP-1 & GIP Receptor Agonists

Class/Main Action	Name	Dose Range	Considerations
GLP-1 RA - Glucagon Like Peptide Receptor Agonist "Incretin Mimetic" <ul style="list-style-type: none"> Increases insulin release with food Slows gastric emptying Promotes satiety Suppresses glucagon 	exenatide (Byetta)	5 and 10 mcg BID	Side effects: nausea, vomiting, weight loss, injection site reaction. Report signs of acute pancreatitis or intestinal blockage (ileus) and stop med. Black box warning: Thyroid C-cell tumor warning (avoid if family history of medullary thyroid tumor). *Significantly reduces risk of CV death, heart attack, and stroke. §Approved to reduce risk of CKD †Approved for pediatrics 10-17 yrs Lowers A1C 0.5 – 1.6% Weight loss: 4-6% body weight loss.
	exenatide XR† (Bydureon)	2 mg 1x a week Pen injector - Bydureon BCise	
	liraglutide*† (Victoza)	0.6, 1.2 and 1.8 mg daily	
	dulaglutide*† (Trulicity)	0.75, 1.5, 3.0 and 4.5 mg 1x a week pen injector	
	semaglutide*§ (Ozempic)	0.25, 0.5, 1.0 and 2.0 mg 1x a week pen injector	
	(Rybelsus) Oral tablet	3, 7, 14 mg - Original dosing. 1.5, 4, 9 mg - New dosing. AM dose, pre-food, w/ water sip	
GLP-1 & GIP Receptor Agonist Activates receptors for GLP-1 (see above) & Glucose-dependent Insulinotropic Polypeptide (GIP).	Tirzepatide (Mounjaro)	2.5, 5.0, 7.5, 10, 12.5 and 15 mg 1x a week injection Single dose via prefilled pen or vial. Adjust dose based on shared decision making and individual goals.	Side effects: nausea, diarrhea, injection site reaction. Report pancreatitis, signs of intestinal blockage. Black box warning: Avoid if family history of medullary thyroid tumor. Lowers A1C ~ 1.8 - 2.4% Weight loss: 7-13% body weight loss at max dose.

Counseling Points: GLP-1 RA & GLP-1/GIP

- ▶ Avoid if personal or family history of medullary thyroid cancer
- ▶ Avoid in combo with DPP-4 inhibitors
- ▶ Watch for intestinal obstruction
- ▶ Use of non-FDA *compounded* products not recommended
- ▶ Avoid with history pancreatitis
- ▶ If on tirzepitide, use back up contraception for first 4 weeks
- ▶ Ask about recent eye exam
 - ▶ Potential increase in diabetes retinopathy



STANDARDS OF CARE | DECEMBER 09 2024
9. Pharmacologic Approaches to Glycemic Treatment: Standards of Care in Diabetes—2025 **FREE**
American Diabetes Association Professional Practice Committee



Sudden discontinuation of semaglutide and tirzepitide results in regain of one-half to two-thirds of the weight loss within 1 year. Consider trying lowest effective dose, using intermittent therapy, or stopping medication followed by close weight monitoring.



Indication Chart for GLP/GIP Receptor Agonists - Diabetes, Weight, CVD and Others

Drug	Type 2 Diabetes	Weight Loss Indication	CV Indication	Other Indication
Exenatide IR (Byetta)	Yes	No	No	
Exenatide ER (Bydureon)	Yes, 10 yrs and older	No	No	
Dulaglutide (Trulicity)	Yes, 10 yrs and older	No	Yes	
Liraglutide (Victoza)	Yes, 10 yrs and older	No	Yes	
Liraglutide (Saxenda)	No	Yes, 12 yrs and older	No	
Semaglutide (Ozempic)	Yes	No	Yes	CKD
Semaglutide (Wegovy)	No	Yes, 12 yrs and older	Yes	MASH
Oral Semaglutide (Rybelsus)	Yes	No	CV benefit, indication pending	
Tirzepatide (Mounjaro)	Yes	No	No	
Tirzepatide (Zepbound)	No	Yes	No	Sleep Apnea

Where are we on this continuum?



Exercise Standards

- ▶ Adults – 150 min/wk moderate intensity
 - ▶ over 3 days a week.
 - ▶ Don't miss > 2 consecutive days w/out exercise
 - ▶ Get up every 30 mins - Reduce sedentary time
 - ▶ Flexibility and balance training 2-3 xs a week (Yoga and Tai Chi)
 - ▶ T1 and T2 – resistance training 2 -3 x's a week



A hard truth

▶ Exercise alone doesn't cause weight loss

▶ But....

- ▶ It helps keep weight off
- ▶ Decreases visceral adiposity
- ▶ Decreases CV Risk

IT TAKES 524 BURPEES

TO BURN OFF 1 LARGE FRIES

BURPEES SUCK, SO CHOOSE WISELY!

@IG.HEALTH



- ▶ To combat the rise in body weight, we need to change the food environment
- ▶ “You cannot outrun an unhealthy diet”.

Good Exercise Info / Quotes



- ▶ **“Passagiata” – take an after meal stroll**
- ▶ Exercise decreases A1C 0.7%
- ▶ No change in body wt, but 48% loss in visceral fat.

“Every minute of activity lowers blood sugar one point.”


“I don’t have time to exercise, I MAKE time.”

6. Glycemic Goals & Hypo

A1C

Blood Pressure

Cardiovascular risk
reduction



6. Glycemic Targets for Non-Pregnant Adults

- ▶ **A1c < 7%** - a reasonable goal for adults.
- ▶ **A1c < 6.5%** - for those without significant risk of hypoglycemia
- ▶ **A1c < 8%** - for those with history of hypoglycemia, limited life expectancy, or those with longstanding diabetes and vascular complications.
- ▶ **A1c Check Frequency:**
 - ▶ If meeting goal - At least 2 times a year
 - ▶ If *not* meeting goal – Quarterly
- ▶ **Also review Ambulatory Glucose Profile**



6. Glycemic Targets

Individualize Targets – ADA

- ▶ Pre-Prandial BG 80- 130
- ▶ 1-2 hr post prandial < than 180

*for nonpregnant adults

- ▶ Time in Range: 70%
- ▶ BG of 70-180 mg/dL



A1c and Estimated Avg Glucose (eAG)

<u>A1c (%)</u>	<u>eAG</u>
5	97 (76-120)
6	126 (100-152)
7	154 (123-185)
8	183 (147-217)
9	212 (170 -249)
10	240 (193-282)
11	269 (217-314)
12	298 (240-347)

***eAG = 28.7 x A1c-46.7 ~ 29 pts per 1%
Translating the A1c Assay Into eAG – ADAG Study***



6. Glycemic Targets: *Standards of Medical Care in Diabetes—2020*

American Diabetes Association
Diabetes Care 2020 Jan; 43(Supplement 1): S66-S76.
<https://doi.org/10.2337/dc20-S006>


Ambulatory Glucose Profile

- ▶ Standardized report with visual cues for those on CGM devices
- ▶ For most with type 1 or type 2 diabetes
 - > 70% of readings within BG range of 70-180mg/dL
 - < 4% of readings < 70 mg/dL
 - < 1% of readings < 54 mg/dL
 - < 25% of readings > 180 mg/dL
 - < 5% of readings > 250 mg/dL



For those with frailty or at high risk of hypoglycemia recommend:

- Target of 50% time in range
- Less than 1% time below range



Favor more stringent goal	Favor less stringent goal
Short diabetes duration	Long diabetes duration
Low hypoglycemia risk	High hypoglycemia risk
Low treatment risks and burdens	High treatment risks and burdens
Pharmacotherapy with cardiovascular, kidney, weight, or other benefits	Pharmacotherapy without nonglycemic benefits
No cardiovascular complications	Established cardiovascular complications
Few or minor comorbidities	Severe, life-limiting comorbidities

Table 6.2

ADA 2025 Goals Summary

A1c less than 7%
(individualize)

- Pre-meal BG 80-130
- Post meal BG <180
- Time in Range (70-180) 70% of time

Blood Pressure
<130/80



Cholesterol

- Statin therapy based on age & risk status
- If 40+ with ASCVD Risk, decrease LDL by 50%, LDL <70
- If 40+ with ASCVD, decrease LDL by 50%, LDL <55

Diabetes Bingo

“DiaBingo” Shout out Right Answer



DiaBingo- G

- G ADA goal for A1c is less than ____%**
- G People with DM need to see their provider at least every month**
- G Blood pressure goal is less than**
- G People with DM should see eye doctor (ophthalmologist) at least**
- G The goal for triglyceride level is less than**
- G Goal for LDL cholesterol for people 40+ with diabetes is _____**
- G The goal for blood sugars 1-2 hours after a meal is less than:**
- G People with DM should get this shot every year**
- G People with DM need to get urine tested yearly for _____**
- G Periodontal disease indicates increased risk for heart disease**
- G The goal for blood sugar levels before meals is:**
- G The activity goal is to do ___ minutes on most days**

Diabetes Care Guidelines- ADA

<u>Test / Exam</u>	<u>Frequency</u>
▶ A1c	At least twice a year
▶ B/P	Each visit
▶ Cholesterol (LDL, HDL, Tri)	Yearly or if med change
▶ Vaccinations	Flu yearly, pneumonia, hep
▶ Weight / BMI	Yearly
▶ UACR/GFR/Creat	Yearly
● Eye exam	Every 1-2 years
● Dental Care	At least twice a year
● Comprehensive Foot Exam	Yearly (more if high risk)
● Physical Activity Plan	As needed to meet goals
● Preconception counseling	As needed

Chronic Kidney Disease— 2025 Update

- ▶ Optimize glucose and BP to protect kidneys.
- ▶ Use SGLT-2 with demonstrated benefit to reduce CKD and CVD*
- ▶ To reduce CV risk and CKD, use a GLP-1* with demonstrated benefit.
- ▶ In people with CKD and albuminuria, a nonsteroidal MRA effective if GFR 25+
- ▶ Aim to reduce urinary albumin by $\geq 30\%$ in people with CKD

- ▶ *SGLT-2i's

- Empagliflozin (Jardiance), canagliflozin (Invokana), dapagliflozin (Farxiga)

- ▶ *GLP-1 RA's

- Semaglutide (Ozempic), liraglutide (Victoza), dulaglutide (Trulicity)

Albuminuria Categories	Urinary Albumin Creatinine Ratio (UACR)
Normal to mildly increased – A1	< 30 mg/g
Moderately increased – A2	30 – 299 mg/g
Severely increased – A3	300 mg/g +

Kidney Disease Stage	GFR
Stage 1 – Normal	90+
Stage 2 – Mild loss	89 - 60
Stage 3a – Mild to Mod	59 - 45
Stage 3b – Mod to Severe	44 - 30
Stage 4 – Severe loss	29 - 15
Stage 5 – Kidney failure	14 - 0

Mr. Jones - What are Your Recommendations?

MJ Profile

64 yr old with type 2 for 11 yrs. Hx of CVD.

Labs:

- ▶ A1c 9.3%
- ▶ LDL 137 mg/dl
- ▶ Triglyceride 260mg/dl
- ▶ UACR 32mg/g GFR 54
- ▶ B/P 132/94

Self-Care Skills

- ▶ Walks dog around block 3 x's a week
- ▶ Bowls every Friday
- ▶ 3 beers daily
- ▶ *What meds?*
- ▶ *What referrals?*
- ▶ *My foot hurts*


Lower Extremities

- ▶ Lift the Sheets and Look at the Feet






No Bathroom Surgery



5.07 monofilament = 10gms linear pressure



Three Most Important Foot Care Tips

- ▶ Inspect and apply lotion to your feet every night before you go to bed.
- ▶ Do NOT go barefoot, even in your house. Always wear shoes!
- ▶ Every time you see your provider, take off your shoes and show your feet.

Diabetes Visit – What do you think?

A small adjustment can make a BIG Difference

- ▶ John arrives at the clinic with a Time in Range slightly above 60%.
- ▶ HCP says “You do know that the goal for TIR is 70% ”
(Door shut)


How Does John Feel?

- ▶ Defeated
- ▶ Embarrassed
- ▶ Ashamed
- ▶ Angry
- ▶ Hurt



How does the HCP feel?

At least 42 factors affect glucose!






Missed Appointments due to Stigma and Shame

A recent survey of over 2,600 people with diabetes across eight countries revealed that nearly 40% of missed doctor's appointments are due to stigma or shame.

Diabetes Distress Stories

Common events you will hear about:

- Scary or embarrassing lows
- Surprising highs
- Difficulty managing BG
- Eating challenges
- Managing all of the tech
- Situations with friends, family, colleagues
- Managing health care (feeling judged and misunderstood), insurance, etc.



Diabetes Distress – Assess Annually

Type 1 Diabetes Distress Scale (T1-DDS)

Instructions: Living with type 1 diabetes can be tough. Listed below are a variety of distressing things that many people with type 1 diabetes experience. Thinking back over the past month, please indicate the degree to which each of the following may have been a problem for you by circling the appropriate number. For example, if you feel that a particular item was not a problem for you over the past month, you would circle 1. If it was very tough for you over the past month, you might circle 6.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20

		Not a problem	Slight problem	Moderate problem	Somewhat serious problem	Serious problem	Very serious problem
1	Feeling that I am not as skilled at managing diabetes as I should be.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
2	Feeling that I don't eat as carefully as I probably should.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
3	Feeling that I don't notice the warning signs of hypoglycemia as well as I used to.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
4	Feeling that people treat me differently when they find out I have diabetes.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
5	Feeling discouraged when I see high blood glucose numbers that I can't explain.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
6	Feeling that my family and friends make a bigger deal out of diabetes than they should.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
7	Feeling that I can't tell my diabetes doctor what is really on my mind.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
8	Feeling that I am not taking as much insulin as I should.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
9	Feeling that there is too much diabetes equipment and stuff I must always have with me.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
10	Feeling like I have to hide my diabetes from other people.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
11	Feeling that my friends and family worry more about hypoglycemia than I want them to.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
12	Feeling that I don't check my blood glucose level as often as I probably should.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
13	Feeling worried that I will develop serious long-term complications, no matter how hard I try.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
14	Feeling that I don't get help I really need from my diabetes doctor about managing diabetes.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
15	Feeling frightened that I could have a serious hypoglycemic event when I'm asleep.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
16	Feeling that thoughts about food and eating control my life.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
17	Feeling that my friends or family treat me as if I were more fragile or sick than I really am.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
18	Feeling that my diabetes doctor doesn't really understand what it's like to have diabetes.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
19	Feeling concerned that diabetes may make me less attractive to employers.	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6
20	Feeling that my friends or family act like "diabetes police" (bother me too much).	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4	<input type="checkbox"/> 5	<input type="checkbox"/> 6

ing
_isted
each
mber.
NOT
I would

Very
serious
problem

6

6

6

6

6

6

6

6

6


6

6

6

6

6



5. Facilitating Positive Health Behaviors and Well-being to Improve Health Outcomes.
Standards of Care in Diabetes—2024
American Diabetes Association Professional Practice Committee

Abstracts View article PDF

Topics: carbohydrates, diabetes mellitus, type 1, diabetes mellitus, type 2, eating, health personnel


Diabetes Care December 2023, Vol. 47, 577-5110 doi:https://doi.org/10.2337/6c24-5095

www.behavioraldiabetes.org

https://professional.diabetes.org/sites/default/files/media/ada_mental_health_toolkit_questionnaires.pdf

Releasing the Brake


- ▶ This strategy recognizes that diabetes distress acts as a brake on the application of existing diabetes knowledge and skills.
- ▶ By releasing the diabetes distress brake through emotion-focused intervention, the negative cycle can be efficiently ended.



Embark Trial – Emotions as Priority

I have finally given myself permission to make addressing the emotional aspects of diabetes a priority.

~Coach Beverly



Diabetes Visit – Let's Go *through*

A small adjustment can make a BIG Difference


- ▶ John arrives at the clinic with a Time in Range slightly above 60%.
- ▶ HCP says “You do know that the goal for TIR is 70% ” (Door closed)
- ▶ HCP smiles and says, “Wow John, I can see you are making an effort to improve your time in range.”
- ▶ Door Open – Connection made

How Does John Feel?


- ▶ Reassured
- ▶ Heard & Seen
- ▶ Recognized
- ▶ Confident
- ▶ Connected



How does the HCP feel?




Relationship Building | Three Tools To Make It Happen




Used with permission from ReVive 5 Program; Larry Fisher, PhD & Susan Guzman, PhD

Commit to Listening at least Half of the Time






- ◆ Ask about their life (SDOH)
- ◆ Assess current self-management behaviors
- ◆ Assess your feelings
- ◆ Accept without judgement
- ◆ Acknowledge one thing they are doing right
- ◆ Advocate for needed resources



- ◆ Beliefs about health and diabetes
- ◆ Barriers can be confused with non-compliance
- ◆ Burnout lookout. On extended diabetes vacation due to diabetes distress?
- ◆ Bouncing back – leaning into resilience



- ◆ Having the Conversation
- ◆ Coaching that highlights *their* knowledge and resilience.
- ◆ Carrots – problem solve together and dig for solutions that are meaningful in everyday life.
- ◆ Compassion for the people in our care and ourselves.
- ◆ Connection through – opening the door.

Thank You



- ▶ Questions?
- ▶ Email: info@diabetesed.net
- ▶ Web: www.diabetesed.net
- ▶ Phone 530-893-8635

