



**Welcome to
Diabetes in the 21st Century**

Beverly Dyck Thomassian, RN, MPH, BC-ADM, CDE
President, Diabetes Education Services
www.DiabetesEd.net

© Copyright 1999-2014, Diabetes Education Services, All Rights Reserved. 

Diabetes in the 21st Century:
A Clinical and Educational Update

1. Describe impact of diabetes
2. Discuss prevention, management strategies
3. Discuss different types of diabetes
4. Describe insulin therapy
5. Review glucose patterns and determine how to adjust therapy to improve glucose.
6. Discuss medical nutrition therapy
7. Gain understanding of Type 2 Meds.
8. Demonstrate successful teaching strategies

CDC Announces



**35% of
Americans will
have Diabetes
by 2050**

Boyle, Thompson, Barker, Williamson
2010, Oct 22-8(1)29
www.pophealthmetrics.com

Diabetes in America 2014

- ▶ 29 million or > 9.3%
- ▶ 27% don't know they have it
- ▶ 37% of US adults have pre diabetes

Diabetes



CDC's Division of Diabetes Translation. National Diabetes Surveillance System available at <http://www.cdc.gov/diabetes/statistics>



Diabetes Education SERVICES

Type 2 in Kids



- ▶ 7 fold increase 1990
- ▶ 1 in 6 overweight kids (age 12- 19) have prediabetes.
- ▶ ~2,500 to 3,700 new cases in U.S. annually.
- ▶ Highest risk: very obese, minority, female, low socioeconomic status, limited education
- ▶ In age range 12-19, less than 1% have Type 2 – NHANES
- ▶ Environmental changes to urgently needed



Diabetes Education SERVICES

Global Epidemic

- ▶ Every 10 seconds
 - ▶ 1 person dies with diabetes
 - ▶ 2 people develop diabetes
- ▶ Every year
 - ▶ 3 million deaths
 - ▶ 6 million new cases
- ▶ World Diabetes Day is November 14
- ▶ March is ADA Sound the Alert Day "find people w/ undetected diabetes"



Diabetes Education SERVICES

World Diabetes Day
November 14

The right education
for all

Diabetes:
protect our future

www.worlddiabetesday.org

The right environment
for all

Diabetes:
protect our future

www.worlddiabetesday.org

Diabetes Education SERVICES

Age-adjusted Diabetes Prevalence
20 yrs or older, by race/ethnicity— U.S. 20014

Age-adjusted* percentage of people aged 20 years or older with diagnosed diabetes, by race/ethnicity, United States, 2010–2012

Non-Hispanic whites	7.6
Asian Americans	9.0
Hispanics	12.8
Non-Hispanic blacks	13.2
American Indians/Alaska Natives	15.9

*Based on the 2000 U.S. standard population.
Source: 2010–2012 National Health Interview Survey and 2012 Indian Health Service's National Patient Information Reporting System.

- Among Hispanic adults, the age-adjusted rate of diagnosed diabetes was 8.5% for Central and South Americans, 9.3% for Cubans, 13.9% for Mexican Americans, and 14.8% for Puerto Ricans.
- Among Asian American adults, the age-adjusted rate of diagnosed diabetes was 4.4% for Chinese, 11.3% for Filipinos, 13.0% for Asian Indians, and 8.8% for other Asians.
- Among American Indian and Alaska Native adults, the age-adjusted rate of diagnosed diabetes varied by region from 6.0% among Alaska Natives to 24.1% among American Indians in southern Arizona.

Diabetes Education SERVICES

Why Should Zip Code Determine Life Expectancy?

STOCKTON
95202
Life Expectancy
73

IRVINE
92606
Life Expectancy
88

California Endowment – look up your zip code at www.measureofamerica.org

Diabetes Education SERVICES

Engaging and supporting Kids to help slow the epidemic

▶ Phases of Life

▶ During Childhood



▶ Environment

- ▶ Access to safe places to exercise
- ▶ Access to healthy foods
- ▶ Access to learning rich environments
- ▶ Access to health care

▶ LifeStyle

- ▶ Limit screen time to 2 hours a day
- ▶ 1 hour a day of activity
- ▶ Healthy Snacks
- ▶ Limit junk food, sugary beverages
- ▶ Fruits and Veggies

Thoughts on Diabetes, Weight, Social Change



▶ “The only way on a societal basis to reduce the prevalence of obesity is through community action” – Dr. Frieden, CDC

- ▶ Obesity (BMI 30+) prevalence 22% to 40%
- ▶ Poverty, Obesity, Diabetes inter-related



Diabetes Education SERVICES

Weight and Gut Bacteria New and Early Research

▶ Lower levels among obese and DM patients compared with healthy controls of:

- ▶ *Firmicutes*: 4% lower in obese patients, 13% lower in DM
- ▶ *Bifidobacteria*: 14% lower in obese patients, 28% lower DM
- ▶ *Clostridium Leptum*: 14% lower in obese patients, 11% lower DM

▶ Based on prospective study involving:

- ▶ 27 morbidly obese pts with mean BMI of 40
- ▶ 26 pts with new type 2 – BMI 29
- ▶ 28 healthy controls (mean BMI 23 kg/m²).



▶ “The human gut microbiome consists of some 100 trillion bacteria, or some 100 trillion friends you didn't know you had.”

- ▶ Yalcin Basaran, MD, presented at [International Endocrine Meeting](#)



Diabetes Education SERVICES

Free Live Webinars and Live Seminars at DiabetesEd.net

- ▶ Free Webinars
 - ▶ Preparing to take CDE
 - ▶ New Frontiers
 - ▶ New Medications
 - ▶ BC-ADM



Diabetes Education SERVICES

Role of the Pancreas Endocrine Functions

Beta Cells - Insulin

- Anabolic hormone - helps store glucose as glycogen in muscle, liver
- ▶ secreted in response to elevated glucose
- ▶ halts breakdown of glycogen in liver
- ▶ increases protein synthesis, fat storage
- ▶ powerful hypoglycemic


Beta Cells - Amylin

- ▶ secreted in 1:1 ratio with insulin
- ▶ Causes satiety
- ▶ Lowers post-prandial glucagon response
- ▶ Slows gastric emptying
- ▶ Type 1 make none
- ▶ Type 2 make less than normal amounts



Diabetes Education SERVICES

Role of the Pancreas Endocrine Functions



Alpha cells - Glucagon

Opposes action of insulin at the liver

- stimulated in response to low glucose levels
- stimulates liver to convert glycogen to glucose
- inhibits liver from glucose uptake
- causes hyperglycemia



Diabetes Education SERVICES


Hormones Effect on Glucose

Hormone	Effect
▶ Glucagon (pancreas)	⬆
▶ Stress hormones (kidney)	⬆
▶ Epinephrine (kidney)	⬆
▶ Insulin (pancreas)	⬇
▶ Amylin (pancreas)	⬇
▶ Gut hormones - incretins (GLP-1) released by L cells of intestinal mucosa, beta cell has receptors)	⬇



Diabetes Education SERVICES

GLP-1 Effects in Humans Understanding the Natural Role of Incretins



Diabetes Education SERVICES


Bariatric Surgery

- ▶ Consider on diabetes pts w/ BMI >35, esp with comorbidities
- ▶ Remission (BG normalized)
 - ▶ rates range from 40 – 95%
 - ▶ Better results with newer diabetes (more beta cell mass)
 - ▶ Due to increase incretins (gut hormones)
- ▶ Still researching long term benefits, cost effectiveness and risk



Diabetes Education SERVICES

Natural History of Diabetes



Signs of Diabetes



- ▶ Polyuria
- ▶ Polydipsia
- ▶ Polyphasia
- ▶ Weight loss
- ▶ Fatigue
- ▶ Skin and other infections
- ▶ Blurry vision
- ▶ Glycosuria, H₂O losses
- ▶ Dehydration
- ▶ Fuel Depletion
- ▶ Loss of body tissue, H₂O
- ▶ Poor energy utilization
- ▶ Hyperglycemia increases incidence of infection
- ▶ Osmotic changes



Diabetes Education SERVICES

Diabetes Classifications

- ▶ Type 1
- ▶ Type 2
- ▶ Gestational
- ▶ Secondary



Diabetes Education SERVICES

Case Study

1. Pt profile: 5'8", 192 lb male

Diabetes 12 years, on insulin 3 yrs

What type of DM and how do you know?



2. Pt profile: 5'6", 108 lb female

On insulin 3u Novolog before meals,
10u Lantus at bedtime

What type of DM and how do you know?



Diabetes Education
SERVICES

Type 1 Rates Increasing Globally

▶ 23% rise in type 1 diabetes incidence from 2001-2009

▶ Why?

- ▶ Autoimmune disease rates increasing over all
- ▶ Changes in environmental exposure and gut bacteria?
- ▶ Hygiene hypothesis
- ▶ Obesity?



Diabetes Education
SERVICES

Type 1 Diabetes Facts

- As many as **3 million Americans** may have type 1 diabetes.
- Each year, approximately **80 people per day** are diagnosed with type 1 diabetes in the U.S.
- Approximately **85 percent** of people living with type 1 diabetes are adults, and **15 percent** are children.
- The rate of type 1 diabetes incidence among children under age 14 is estimated to **increase by 3 percent annually** worldwide.
- Type 1 diabetes accounts for **\$14.9 billion** in healthcare costs in the U.S. each year.

Source: JDRF



Diabetes Education
SERVICES

Type 1 – 10% of all Diabetes Genetics and Risk Factors

- Auto-immune pancreatic beta cells destruction
- Most commonly expressed at age 10-14
- Insulin sensitive (require 0.5 - 1.0 units/kg/day)

- Combo of genes and environment:
 - Autoimmunity tends to run in families
 - Higher rates in non breastfed infants
 - Viral triggers: congenital rubella, coxsackie virus B, cytomegalovirus, adenovirus and mumps.



Diabetes Education SERVICES

Incidence of Type 1 in Youth



- ▶ **General Pop 0.3%**
- ▶ **Sibling 4%**
- ▶ **Mother 2-3%**
- ▶ **Father 6-8%**
- ▶ Rate doubling every 20 yrs
- ▶ Many trials underway to detect and prevent (Trial Net)



Diabetes Education SERVICES

Autoantibodies Assoc w/ Type 1

Panel of autoantibodies –

- ▶ GAD65 - Glutamic acid decarboxylase –
- ▶ ICA - Islet Cell Cytoplasmic Autoantibodies
- ▶ IAA - Insulin Autoantibodies



Diabetes Education SERVICES

Type 1 Diabetes Associated with other immune conditions

- ▶ Celiac disease (gluten intolerance)
- ▶ Thyroid disease
- ▶ Addison's Disease
- ▶ Rheumatoid arthritis
- ▶ Other



Diabetes Education SERVICES

What Does Type 1 Look Like?



Mary Tyler Moore



Adam Morrison



Nick Jonas



Justice Sonia Sotomayor

Bret Michaels



From Debbie Nagata's slide collection



Diabetes Education SERVICES

Medalist Study – Harvard Joslin Diabetes Center

- ▶ After 50 years with diabetes
 - ▶ Many still produced some insulin
 - ▶ Many had no eye disease



Diabetes Education SERVICES

Type 1 Summary

- ▶ Autoimmune
- ▶ Complete pancreatic destruction
- ▶ Need insulin shots
- ▶ Often first present in DKA



Diabetes Education SERVICES

Type 1 in Hospital


- ▶ 43 yr old admitted to evaluate angina.
- ▶ Morning blood sugar is 92.
- ▶ Based on Regular insulin sliding scale, no insulin required.
- ▶ Breakfast tray shows up and patient says, I need my insulin shot before I eat.



What do you say?




Diabetes Education SERVICES



Diabetes Education SERVICES

Natural Progression of Type 2 Diabetes



Adapted from Bergenstal et al. 2000; International Diabetes Center.



Diabetes Education SERVICES

Cardio Metabolic Risk - 5 Hypers -

- ▶ Hyperinsulinemia (resistance)
- ▶ Hyperglycemia
- ▶ Hyperlipidemia
- ▶ Hypertension
- ▶ Hyper"waistline"emia (35" women, 40" men)



Manifestations of Insulin Resistance



Diabetes Education SERVICES

Diabetes 2 - Who is at Risk?

(ADA Clinical Practice Guidelines)

1. Testing should be considered in all adults who are overweight (BMI \geq 25) and have additional **risk factors**:
 - ▶ First-degree relative w/ diabetes
 - ▶ Member of a high-risk ethnic population
 - ▶ Habitual physical inactivity
 - ▶ PreDiabetes
 - ▶ History of heart disease



Diabetes Education SERVICES

Diabetes 2 - Who is at Risk?

(ADA Clinical Practice Guidelines)



Risk factors cont'd

- ▶ HTN - BP > 140/90
- ▶ HDL < 35 or triglycerides > 250
- ▶ baby >9 lb or history of Gestational Diabetes Mellitus (GDM)
- ▶ Polycystic ovary syndrome (PCOS)
- ▶ Other conditions assoc w/ insulin resistance:
 - ▶ Severe obesity, acanthosis nigricans (AN)



Diabetes Education SERVICES

Acanthosis Nigricans (AN)

- ▶ Signals high insulin levels in bloodstream
- ▶ Patches of darkened skin over parts of body that bend or rub against each other
 - ▶ Neck, underarm, waistline, groin, knuckles, elbows, toes
 - ▶ Skin tags on neck and darkened areas around eyes, nose and cheeks.
- ▶ No cure, lesions regress with treatment of insulin resistance



Diabetes Education SERVICES


Diabetes Detectives Needed



- ▶ On average – takes 6.5 years to diagnose diabetes
- ▶ 1/4 of all people with diabetes don't know they have it



Diabetes Education SERVICES



Comparison of Type 1 and Type 2

	<u>Type 1</u>	<u>Type 2</u>
Obesity	x	xxx
Insulin dependence	xxx	30%
Respond to oral agents	0	xxx
Ketosis	xxx	x
Antibodies present	xxx	0
Typical Age of onset	teens	adult
Insulin Resistance	0	xxx

Diabetes Education SERVICES

Diabetes is also associated with:

- ▶ Fatty liver disease
- ▶ Obstructive sleep apnea
- ▶ Cancer; pancreas, liver, breast
- ▶ Alzheimer's
- ▶ Depression

Diabetes Education SERVICES

Gestational DM ~ 7% of all Pregnancies

- ▶ GDM prevalence increased by
 - ▶ ~10–100% during the past 20 yrs
- ▶ Native Americans, Asians, Hispanics, African-American women at highest risk
- ▶ Immediately after pregnancy, 5% to 10% of GDM diagnosed with type 2 diabetes
- ▶ Within 5 years, 50% chance of developing DM in next 5 years.



Diabetes Education SERVICES

Diabetes in pregnant mothers associated with ...

- ▶ Offspring
 - ▶ Fetal Complications
 - ▶ Obesity and diabetes later in life
- ▶ Mother
 - ▶ More complicated pregnancy and delivery
 - ▶ Diabetes later in life
- ▶ Intrauterine environment is important



Diabetes Education SERVICES

Screen Pregnant Women Before 13 weeks

- ▶ Screen for undiagnosed Type 2 at the first prenatal visit using standard risk factors.
- ▶ Women found to have diabetes at their initial prenatal visit treated as “Diabetes in Pregnancy”
- ▶ If normal, recheck at 24-28 weeks



Diabetes Education SERVICES

Increasing Prevalence – A public health perspective

- ▶ Body weight before and during pregnancy influences risk of GDM and future diabetes
- ▶ Children born to women with GDM at greater risk of diabetes
- ▶ Focus on prevention



Diabetes Education SERVICES

Postnatal Health: Maternal Behavior

- ▶ Encourage breastfeeding for one year
 - ▶ (25% of women achieving this goal)
- ▶ Screening 6-12 weeks post partum using non-pregnant OGTT criteria (50%)
- ▶ Repeat at 3 yr intervals or signs of DM
- ▶ Encourage weight control and exercise
- ▶ Make sure connected with health care
- ▶ Preconception counseling



Diabetes Education SERVICES

Start Metformin therapy

- ▶ For women with PreDiabetes and History of GDM



Diabetes Education SERVICES

Other Causes of Hyperglycemia

- ▶ Steroids
- ▶ Agent Orange
- ▶ Tube feedings / TPN
- ▶ Transplant medications
- ▶ Cystic Fibrosis

Regardless of cause, requires treatment

- ▶ Insulin always works
- ▶ Sign of pancreatic malfunction



Diabetes Education SERVICES

Flash Mob – World Diabetes Day to Beat It –

- ▶ March R/C/R
- ▶ Fred Astaire
- ▶ Point R/L
- ▶ Arms up, down
- ▶ Shoulder Walk
- ▶ Punch down/up
- ▶ Scoot Rt/Left
- ▶ Reach up R/L
- ▶ Shoulder Walk
- Open door
- Ride Horse
- Scoot Rt/Left
- Turn R & Clap, then L
- Shoulder Walk
- Punch down/up
- To print out steps, go to DiabetesEd.net, bottom of home page – Flash Mob steps



Diabetes Education SERVICES

Life Study – Mrs. Jones

Mrs. Jones is 62 years old, overweight and complaining of feeling tired and urinating several times a night. She is admitted with a urinary tract Infection. Her WBC is 12.3, glucose 237. She is hypertensive with a history of gestational diabetes. No ketones in urine.

- ▶ What are her risk factors, signs of diabetes
- ▶ What type of diabetes does she have?
- ▶ Does she have insulin resistance?



What Do You Say? Mrs. Jones asks you

- ▶ What is type 2 diabetes?
- ▶ Will this go away?
- ▶ Will I get complications?
- ▶ Will I need to take diabetes medication for the rest of my life?
- ▶ How come I got diabetes?
- ▶ Do I have to check my blood sugars?



Diabetes Education SERVICES

Running into Roadblocks?

- ▶ *HUG Patients*
- ▶ Help with
- ▶ Unconditional
- ▶ Guidance and Support

Anne Peters, MD, CDE
ADA Post Grad

▶ **Unconditional Positive Regard –**
involves showing complete support and acceptance of a person no matter what that person says or does.
Carl Rogers



Diabetes Education SERVICES

No one is Unmotivated

... to lead a long and healthy life

- ▶ **These are the 3 usual Critical Barriers**
- ▶ Perceived worthlessness
- ▶ Too many personal obstacles
- ▶ Absence of support and resources



Bill Polonsky, PhD, CDE



Diabetes Education SERVICES

Overcoming barriers

- ▶ Confront the key misbelief. Ask the question, does dm cause complications?
 - ▶ Offer pts evidence based hope message –
 - ▶ Frequent contact
 - ▶ Paired glucose testing
 - ▶ Ask pt, “Tell me 1 thing that is driving you crazy about your diabetes”
 - ▶ Discuss medication beliefs
 - ▶ To improve outcomes, see pts more often
- Bill Polonsky, PhD, CDE



Diabetes Education SERVICES

How Often Should I Check?

- Be realistic!!
- Type 1 – as often as needed
- Type 2 – as needed
- Consider:
 - ▶ Types and timing of meds
 - ▶ Goals
 - ▶ Ability (physical and emotional)
 - ▶ Finances



Diabetes Education SERVICES


How will it help me?

- ▶ See if your treatment plan is working
- ▶ Make decisions regarding food and/or med adjustment when exercising
- ▶ Find out how that pizza affected your BG
- ▶ Avoid unwanted weight gain
- ▶ Enhanced athletic performance
- ▶ Find patterns
- ▶ Manage illness



Diabetes Education SERVICES

New Meters – a little goes a long way



- 0.3 microliters of blood
- minimal pain



Customer Service (toll-free): Look for 800 number



Diabetes Education SERVICES

DiaBingo

- B Frequent skin and yeast infections.....
- B A BMI of ____ or greater is considered overweight
- B To reduce complications, control **A1c**, **B**lood pressure, **C**holesterol
- B PreDiabetes – fasting glucose level of ____ to ____
- B Erectile dysfunction indicates greater risk for ____
- B Diabetes – fasting glucose level ____ or greater
- B Type 1 diabetes is best described as an _____ disease
- B People with diabetes are _____ times more likely to die of heart dx
- B Elevated triglycerides, < HDL, smaller dense LDL
- B Each percentage point of A1c = _____ mg/dl glucose
- B At dx of type 2, about ____% of the beta cell function is lost
- B Diabetes – random glucose ____ or greater



Diabetes Education SERVICES

Complications - Why?



- ▶ Degree of hyperglycemia “glucose toxicity”
- ▶ Duration of hyperglycemia
- ▶ Genes
- ▶ Multiple risk factors: smoking, vascular disease, dyslipidemia, hypertension, other



Diabetes Education SERVICES

Diabetes Complications

- ▶ Heart disease leading cause of death.
- ▶ CAD death rates are about 2 -4x's as high as adults without diabetes (it's not getting better)
- ▶ Risk of stroke is 2 - 4 times higher
- ▶ 60% - 65% of people with DM have HTN.
- ▶ DM accounts for 40% of new cases of ESRD
- ▶ 60 - 70% have mild - severe forms of neuropathy
- ▶ Diabetes is the leading cause of blindness
- ▶ Accounts for 50% of lower limb amputations



Diabetes Education SERVICES

Control Matters

- ▶ **Trials**
- ▶ **Practice Recommendations**



Diabetes Education SERVICES

Financial Advisor

- ▶ Mid 30s, friendly, he smiles to greet you and you notice his gums are inflamed. You'd guess a BMI of 26 or so, with most of the extra weight in the waist area.
- ▶ If you could give him some health related suggestions, what would they be?



Diabetes Education SERVICES

Can Type 2 be Prevented in Older Adults?



- Physical activity (30 mins a day)
- Dietary score (higher fiber intake, low saturated fat and *trans*-fat, lower mean glycemic index)
- Not Smoking
- Alcohol use (up to 2 drinks a day);
- BMI <25 and waist circumference

Dariusz Mozaffarian, MD,
Arch Intern Med. 2009;169(8):798-807.

Overall, 9 of 10 new cases of diabetes attributable to these 5 lifestyle factors.

89% risk reduction when all at goal.

35% rel risk reduction for each additional



Diabetes Education SERVICES

We Made It – 3129 miles- Now Walking Back

Track miles with fitbit
Fitbit.com
Lifespan Treadmill
Desk - Amazon

Can we stop pre diabetes from progressing?

3, 234 people w/ Pre-Diabetes randomized:

- ▶ Placebo
- ▶ Diet/Exercise or
- ▶ Metformin

over a three year period

Diabetes Prevention Program (DPP) 2001



Diabetes Education SERVICES

Diabetes Prevention Program

- ▶ Standard Group - 29% developed DM
- ▶ Lifestyle Results - 14% developed DM
 - ▶ 58% (71% for 60yrs +) Risk reduction
 - ▶ 30 mins daily activity
 - ▶ 5-7% of body wt loss
- ▶ Metformin 850 BID - 22% developed DM
- ▶ 31% risk reduction (less effective with elderly and thinner pt's)




Diabetes Education SERVICES

Weight loss and Prevention

- ▶ For every 2.2 pounds of weight loss, risk of type 2 diabetes was reduced by 13%.



Diabetes Education SERVICES



Diabetes Education SERVICES

ABCs of Diabetes

A1C

Blood Pressure

Cholesterol

Standards of Medical Care – American Diabetes Association



Diabetes Education SERVICES

Glucose and BP Control Matter

- ▶ 1% decrease in A_{1c} reduces microvascular complications by 35%
- ▶ 1% decrease in A_{1c} reduces diabetes related deaths by 25%
- ▶ B/P control (144/82) reduced risk of:
 - ▶ Heart failure (56%)
 - ▶ Stroke (44%)
 - ▶ Death from diabetes (32%)

Lancet 352: 837-865, 1998



Diabetes Education SERVICES

A1c and Estimated Avg Glucose (eAG) 2008

A1c (%)	eAG
5	97
6	126
7	154
8	183
9	212
10	240
11	269
12	298

Order teaching tool kit free at diabetes.org



$eAG = 28.7 \times A1c - 46.7 \sim 29$ pts per 1%

*Translating the A1c Assay Into Estimated Average Glucose Values – ADAG Study
Diabetes Care: 31, #8, August 2008*



Diabetes Education SERVICES

ABCs of Diabetes –

- ▶ **A1c less than 7% (avg 3 month BG)**
 - ▶ Pre-meal BG 70-130
 - ▶ Post meal BG <180
- ▶ **Blood Pressure < 140/80**
- ▶ **Cholesterol**
 - ▶ HDL >40
 - ▶ LDL <100 (if CHD, <70)
 - ▶ Triglyceride < 150



Diabetes Education SERVICES


“Legacy Effect”

- ▶ **For participants of DCCT and UKPDS**
 - ▶ long lasting benefit of early intensive BG control prevents
 - ▶ microvascular complications
 - ▶ Macrovascular complications (15-55% decrease)
 - ▶ Even though their BG levels increased over time
 - ▶ Message – Catch early and Treat aggressively



Diabetes Education SERVICES

How are we doing? Reaching goal



Diabetes Care, 2/13



Diabetes Education SERVICES

Vaccinations- Immunizations

- ▶ Flu vaccine
 - ▶ every year starting 6 months
- ▶ Pneumococcal starting at 2 years.
 - ▶ One time Revaccination for those over 64 and had first vaccine >5 years prior
- ▶ Hepatitis B Vaccine (ADA Stds 2013, pg s28)
 - ▶ For diabetes pts age 19 – 59 (not previously vaccinated)
 - ▶ Double risk of Hep B due to lancing devices/ glucose meter exposure



Diabetes Education SERVICES

DiaBingo- G

- G ADA goal for A1c is less than ____%
- G People with DM need to see their provider at least every month
- G Blood pressure goal is less than
- G People with DM should see eye doctor (ophthalmologist) at least
- G The goal for triglyceride level is less than
- G Goal for my HDL cholesterol is more than
- G The goal for blood sugars 1-2 hours after a meal is less than:
- G People with DM should get this shot every year
- G People with DM need to get urine tested yearly for _____
- G Periodontal disease indicates increased risk for heart disease
- G The goal for blood sugar levels before meals is:
- G The activity goal is to do ___ minutes on most days



Diabetes Education SERVICES

Mr. Jones - What are Your Recommendations?

Patient Profile

64 yr old with type 2 for 11 yrs. Hx of CVD.

Labs:

- ▶ A1c 9.3%
- ▶ HDL 37 mg/dl
- ▶ LDL 114 mg/dl
- ▶ Triglyceride 260mg/dl
- ▶ Proteinuria - neg
- ▶ B/P 142/92

Self-Care Skills

- ▶ Walks dog around block 3 x's a week
- ▶ Bowls every Friday
- ▶ 3 beers daily
- ▶ Widowed, so usually eats out
- ▶ 15 lbs overweight
- ▶ My foot hurts



Diabetes Education SERVICES

Diabetes Care Guidelines- ADA

Test / Exam	Frequency
› A1c	At least twice a year
› B/P	Each diabetes visit
› Cholesterol (LDL, HDL, Tri)	Yearly (less if normal)
› Weight	each diabetes visit
› Microalbumin/GFR/Creat	Yearly
• Eye exam	Yearly
• Dental Care	At least twice a year
• Comprehensive Foot Exam	Yearly (more if high risk)
• Physical Activity Plan	As needed to meet goals
• Preconception counseling	As needed



Diabetes Education SERVICES


Foot Care

Lift the sheets
and look at the
Feets!



Diabetes Education SERVICES

Foot Wounds



Blisters
Calluses

Ulcers

Bone infection



Diabetes Education SERVICES

No Bathroom Surgery

Diabetes Education SERVICES

5.07 monofilament = 10gms linear pressure
If pt can't feel pressure = neuropathy

Diabetes Education SERVICES

3 Most Important Foot Care Tips

- ▶ Inspect and apply lotion to your feet every night before you go to bed.
- ▶ Do NOT go barefoot, even in your house. Always wear shoes!
- ▶ Every time you see your doctor, take off your shoes and show your feet.

Diabetes Education SERVICES

Glucose Management and Hospitalized Patients



▶ In hospitalized patients with critical illness, hyperglycemia is a signal that warrants our attention.



Diabetes Education SERVICES

Hospitals and Hyperglycemia – What’s the Big Deal?


- ▶ Hyperglycemia is associated with increased morbidity and mortality in hospital settings.
 - ▶ Acute Myocardial Infarction
 - ▶ Stroke
 - ▶ Cardiac Surgery
 - ▶ Infection
 - ▶ Longer lengths of stay



Diabetes Education SERVICES

Hyperglycemia*: A Common Comorbidity in Medical-Surgical Patients in a Community Hospital

Umplieriez G et al, J Clin Endocrinol Metabol 87:976, 2002




- Normoglycemia
 - Known Diabetes
 - New Hyperglycemia
- n = 2,020
 * Hyperglycemia: Fasting BG \geq 126 mg/dl or Random BG \geq 200 mg/dl X 2



Diabetes Education SERVICES

Effect of Hyperglycemia on Hospital Mortality



BG Above Normal = Trouble

- ▶ Pre Diabetes
 - ▶ Fasting Glucose = 100-125mg/dl
 - ▶ A1c 5.7 – 6.4%
- ▶ Diabetes
 - ▶ Fasting Glucose = 126 mg/dl +
 - ▶ Random Glucose = 200 mg/dl +
 - ▶ A1c 6.5% +



▶ Any blood glucose above 140 requires treatment
Umpierrez et al

WHAT SHOULD WE AIM FOR?

- Critically Ill pts
- BG > 180- Start insulin
 - BG goal 140-180



- Non Critically Ill patients BG Goals
- Premeal <140
 - Post meal <180

• Insulin therapy preferred treatment

Consensus: Inpt Hyperglycemia, Endocr Pract. 2009;15 (No.4)

Management of Hyperglycemia and Diabetes

- ▶ Stop oral agents (ie) metformin & sulfonylurea on admission
- ▶ “The sole use of Sliding Scale insulin is discouraged” – ADA 2014
- ▶ For discharge, oral meds can be resumed

Start Basal/bolus therapy

- ▶ NPH and Regular insulin
- ▶ Long-acting and rapid-acting insulin
- ▶ Premixed insulin




Diabetes Education SERVICES

In Patient Strategies – Start Early, Focus on Survival Skills



Diabetes Education SERVICES

Discharge insulin Algorithm



Clinical Guidelines for the Management of Hyperglycemia in Hospitalized Patients in a Non-Critical Care Setting



Diabetes Education SERVICES

Now What?

▶ Nurse had an emergency and pt already ate lunch?



▶ Nurse administered insulin and pt only ate a few bites of turkey and drank non sugar tea?

▶ You just gave 3 units of Aspart and patient needs to go to OR NOW!

5 Survival Skills

1. Basics of Diabetes
 2. Can patient perform self blood glucose monitoring? Do they need meter?
 3. Can pt safely take meds / insulin? Teach side effects.
 4. Meal Planning?
 5. Self Care including hypo prevent/treat
- ▶ Follow-Up plan - Does pt know who to contact when need help?
- ▶ Diabetes Ed, PCP, Home Health



Diabetes Education SERVICES

Bottom Line

- ▶ 30-40% of hospitalized patients have diabetes
 - ▶ 10% aren't officially diagnosed
- ▶ Cardiovascular disease is the leading cause of hospitalization for people with diabetes
- ▶ Look for patients with hyperglycemia and cardiometabolic risk factors: smokers, HTN, central obesity, abnormal lipids, Acanthosis.
- ▶ Provide education and promote self-advocacy



Diabetes Education SERVICES


Diabetes Self-Management

- ▶ Self Monitor Blood Glucose
- ▶ Meal Plan
- ▶ Exercise / Activity
- ▶ Medications




Diabetes Education SERVICES

Obesity in America



- ▶ 68% overweight or obese
- ▶ 34% BMI 30+, 34% BMI 25-29
- ▶ 1/3 of all overwt people don't get diabetes
- ▶ We burn 100 cals less a day at work
- ▶ Overall, food costs ~ 10-15% of income
- ▶ Calorie Intake is on the rise



Average Daily Per Capita Calories Adjusted for Weight. Source: Economic Research Service of the United States Department of Agriculture (Per Capita) Data System.



Diabetes Education SERVICES


Average American Consumes 25 teaspoons of sugar a day (400 cals)

- ▶ Warning label on sodas proposed
- ▶ One soda has 12 teaspoons of sugar
- ▶ On avg, 1 person consumes 40 gallons of soda each year
- ▶ ADA guidelines "limit sodas and beverages with sugar, High Fructose Corn Syrup, (HFCS)



Diabetes Education SERVICES

BMI – Visual Image



Diabetes Education SERVICES

Medical Nutrition Therapy – ADA 2014 Updates



- No ideal percentage of calories from protein, carbohydrate and fat for people with diabetes.
- Macronutrient distribution should be based on an *individualized assessment* of eating patterns, preferences and metabolic goals.



Diabetes Education SERVICES

Medical Nutrition Therapy – ADA 2014

- ▶ Focus on the Individual
- ▶ Maintain pleasure of eating
- ▶ Provide positive messages about food
- ▶ Limit food choices only when backed by science
- ▶ Provide practical tools
- ▶ Refer to a RD and Diabetes Education – Lowers A1c by 1-2%



Diabetes Education SERVICES

Sodium, Fat and Fiber

- ▶ Sodium – Try and keep less than 2,300 mg a day
- ▶ Vitamin and mineral supplements not recommended -lack of evidence.
- ▶ Fat - same as recommended for general population
 - ▶ Less than 10% saturated fat,
 - ▶ Limit trans fats
 - ▶ Less than 300 mg cholesterol daily
 - ▶ Mediterranean Diet looks like good option
- ▶ Fiber 25 -38 gms a day



Diabetes Education SERVICES

Approach Depends on Patient

- New Type 2
 - Portion Control
 - Plate Method
 - Record Keeping
 - Education
- On Insulin?
 - Carb counting
 - Post prandial checks



Diabetes Education SERVICES

Losing 2-8kg Early in diagnosis Type 2 Helpful

ADA 2014

- ▶ Weight Loss –
 - ▶ *The optimal macronutrient intake to lose weight not known*
 - ▶ *The literature does not support one particular nutrition therapy to reduce weight, but rather a spectrum of eating patterns that result in reduced energy intake.*
- ▶ To lose one pound – avoid 3,500 cal
 - ▶ Decrease intake 250-500 cal daily + exercise



Diabetes Education SERVICES

Successful weight loss strategies include

- ▶ Weekly self-weighing
- ▶ Eat breakfast
- ▶ Reduce fast food intake.
- ▶ Decrease portion size
- ▶ Increase physical activity
- ▶ Use meal replacements
- ▶ Eat healthy foods



Diabetes Education SERVICES

Diabetes Prevention Program Focus on fat = wt loss success

To help you lose weight and improve your health, stay as close as possible to your fat and calorie goals.
Find your starting weight below. Your fat and calorie goals are in the same row. Circle your fat and calorie goals.

Weight (lb)	Fat Goal (grams)	Calorie Goal
120-174	33	1,200
175-219	42	1,500
220-249	50	1,800
>250	55	2,000

<http://www.cdc.gov/diabetes/prevention/recognition/curriculum.htm>



Diabetes Education SERVICES


Public Health Issue?

- ▶ 66% of our people are obese/overweight
- ▶ Rates of gestational diabetes on rise
- ▶ 30% of kids are obese/overweight



Diabetes Education SERVICES


How nutrients affect blood sugar



Diabetes Education SERVICES

Teaching About Eating Healthy

- Major food groups
- "Handy Diet"
- Plate Method
- Exchange Lists
- Food Diaries / Glucose Records
- Carbohydrate Counting



Assess what is best for the situation.



Diabetes Education SERVICES

Move toward the Tomato



Diabetes Education SERVICES

ADA recommendation Eat Less Junk Food & Sugary Drinks –

- ▶ Less Processed Foods
- ▶ Less Sugary Beverages
 - ▶ increase visceral adiposity
 - ▶ With sugar or
 - ▶ High fructose corn syrup

- ▶ Soda Tax?
- ▶ Junk Food Tax?



Diabetes Education SERVICES

10 Superfoods

- ▶ Beans
- ▶ Dark Green Leafy Veggies
- ▶ Citrus Fruit
- ▶ Sweet Potatoes
- ▶ Berries
- ▶ Tomatoes
- ▶ Fish High in Omega-3 Fatty Acids
- ▶ Whole Grains
- ▶ Nuts
- ▶ Fat-Free Milk and Yogurt



Diabetes Education SERVICES

USDA Food Pyramid www.myplate.gov

Balancing Calories

- ▶ Enjoy your food, but eat less.
- ▶ Avoid oversized portions.

Foods to Increase

- ▶ Make half your plate fruits and vegetables.
- ▶ Make at least half your grains whole grains.
- ▶ Switch to fat-free or low-fat (1%) milk.

Foods to Reduce

- ▶ Compare sodium in foods like soup, bread, and frozen meals — and choose the foods with lower numbers.
- Drink water instead of sugary drinks.



Diabetes Education SERVICES

Carbs affect Post meal Blood Glucose

- o Starch
- o Fruit
- o Milk
- o Desserts

Starchy foods



*ADAM



Diabetes Education SERVICES

Carbohydrate Needs for Most Adults

	<u>Grams</u>	<u>Servings</u>
Each Meal	45-60 gm	3 - 4
Snacks	15-30 gm	1- 2



Carbs affect Post Meal Blood Glucose



Diabetes Education SERVICES

Choose Healthy Carbs

- o Carbs have fiber, vitamins, minerals and phytonutrients
- o 25 gms of fiber a day
- o Power Carbs include:
 - o Beans
 - o Veggies
 - o Fruits
 - o Whole grain foods



Diabetes Education SERVICES

Handy Meal Plan

▶ Per Meal Serving


- ▶ Each finger = 15 gms carb (can have 3-4 servings/meal)
- ▶ Palm of hand = 3 oz's protein
- ▶ Thumbnail = 1 tsp fat serving



Diabetes Education SERVICES

Carb Counting - Starch


Each Food has:
80 Calories
15 grams carb



Diabetes Education SERVICES

Carb counting- fruit

Each Food has:
60 Calories
15 grams carb



Diabetes Education SERVICES

Carb Counting - Milk

Each Food has:
90-150 calories
12-15 grams carb

1 slice bread

- 8 oz buttermilk
- 1 packet diet hot cocoa
- 8 oz milk
- 8 oz soy milk
- 6 oz plain yogurt
- 6 oz light fruit yogurt

Diabetes Education SERVICES

Carb Counting - Sweets

Each Food has:
Calories vary
15 grams carb

1 slice bread

- 2 inch square cake or brownie, unfrosted
- 1/2 cup diet pudding
- 1/2 cup regular jello
- 2 tsp light syrup
- 2 small cookies
- 1/2 cup ice cream or frozen yogurt
- 1/2 cup sherbet
- 1/4 cup sorbet
- 1 tbsp syrup, jam, jelly, table sugar, honey

Diabetes Education SERVICES

Go Lean with Protein

- Choose lean protein
 - Poultry, fish, egg, lean beef
 - Plant sources- beans, lentils, nuts
 - Low fat cheese- cottage cheese, mozzarella cheese
- Limit high fat protein
 - Bacon & sausage
 - High fat cuts of beef
 - Whole milk cheese
- Serving size
 - 1 oz = 1/4 cup
 - 3 oz = deck of cards

Diabetes Education SERVICES

Fats- Aim for heart health

• Saturated fats (LIMIT)

- Solid
- Animal
- Tropical (palm, coconut)
- Trans fats (deep fried)

○ Monounsaturated

- Olive & canola oils
- Nuts
- Avocado



○ Polyunsaturated

- veg oils: canola, corn, walnut, safflower, soybean

Serving sizes

- 1 tsp butter, margarine, oil, mayonnaise
- 1 Tbsp salad dressing, cream cheese, seeds
- 2 Tbsp avocado, cream, sour cream
- 1 slice bacon



Diabetes Education SERVICES

Using Alcohol Safely

▶ Women- 1 or fewer alcoholic drinks a day

▶ Men 2 or fewer alcoholic drinks a day

▶ 1 alcoholic drink equals

- ▶ 12 oz beer, 5 oz glass of wine, or 1.5 oz distilled spirits (gin etc)

▶ If drink, limit amount and drink w/ food.

▶ Ask HCP if safe for you to drink. Tell them your usual quantity and frequency.

▶ Can cause hypo and worsen neuropathy



Diabetes Education SERVICES

Ms. Gonzales' Daily Meal plan

Break	Lunch	Dinner	Night
5 corn tortillas, 1/2 c. beans, salsa, peppers, egg beaters	Sandwich, low fat potato chips, 1c. juice, 2-4 lowfat cookies	Lg bowl low salt soup, 1c. rice, BBQ meat, salad & cooked vegs 1 glass wine	1 bowl of cereal
Avg BG 120's	Avg BG 200's	Avg BG 200's	Avg BG 180's



Diabetes Education SERVICES

Resources

- ▶ www.eatright.org American Dietetic Association website for nutrition information, resources, and access to Registered Dietitians
- ▶ www.diabetes.org American Diabetes Association website, advocates to prevent, cure and improve the lives of all people affected diabetes
- ▶ www.americanheart.org American Heart Association website; resources, recipes and tips; learn about efforts to reduce death caused by cardiovascular disease
- ▶ www.dce.org/publications/education-handouts/



Diabetes Education SERVICES

Resources

- ▶ www.nhlbi.nih.gov contains information for professionals and the general public about heart and vascular diseases, lung diseases, blood diseases.
- ▶ www.niddk.nih.gov National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) information and resources clearinghouse.



Diabetes Education SERVICES

DiaBingo - N

- N Injected hormone called an incretin mimetic
- N DPP demonstrated that exercise and diet reduced risk of DM by ___ %
- N An _____ a day can help prevent heart attack and stroke
- N Rebound hyperglycemia
- N Scare tactics are effective at motivating patients to change behavior
- N Losing ___ % of body weight, can improve blood glucose, BP, lipids
- N Drugs that can cause hyperglycemia
- N 2/3 cups of rice equals _____ serving carbohydrate
- N A1c of 7% equals glucose of _____
- N One % drop in A1c reduces risk of complications by ___ %
- N 1 gm of fat equal _____kilo/calories
- N Metabolic syndrome = hyperglycemia, hyperlipidemia, hypertension
- N 1% A1c = _____ of Blood Glucose




Diabetes Education SERVICES

Insulin – the Ultimate Hormone Replacement Therapy

Objectives:

- Discuss the actions of different insulins
- Describe using pattern management as an insulin adjustment tool.



Diabetes Education SERVICES

Psychological Insulin Resistance (PIR)




- ▶ 50% of providers in study threatened pts “with the needle”.
- ▶ Less than 50% of providers realized insulins’ positive effect on type 2 dm
- ▶ Most pts don’t believe that insulin would “better help them manage their diabetes”.
- ▶ Solutions: Find the root of PIR and address it, use more insulin pens

Diabetes Attitudes, Wishes, Needs Study - Rubin



Diabetes Education SERVICES

Needle Size often a Barrier Size *Does* Matter




BD Nano 4mm and BD Mini 5mm only available in pen needles

- ▶ Use more short needles – 4 mm
- ▶ Effective for pts with BMI of 24- 49
- ▶ Keeps it subq
- ▶ If pt thin, inject at angle
- ▶ To avoid leakage, count to 10 before withdrawing needle
- ▶ ½ the patients who could benefit from insulin are not using it due to needle phobias
- ▶ Consider inhaled insulin



Diabetes Education SERVICES

Physiologic Insulin Secretion: 24-Hour Profile



Insulin Action Teams

- ▶ **Bolus: lowers after meal glucose levels**
 - ▶ Rapid Acting
 - ▶ Aspart, Lispro, Glulisine
 - ▶ Short Acting
 - ▶ Regular
 - ▶ Afrezza - Inhaled
- ▶ **Basal: controls glucose between meals, hs**
 - ▶ Intermediate
 - ▶ NPH
 - ▶ Long Acting
 - ▶ Detemir (Levemir)
 - ▶ Gargine (Lantus)




Cost Per Vial in Northern CA

Per vial cost	Walmart	Walgreens	Costco
Regular Insulin	\$25*	\$92	\$99
NPH	\$25*	\$92	\$99
70/30	\$25*	\$92	\$101
Humalog	\$200	\$220	\$178
Novolog	\$197	\$217	\$178
Apidra	\$180	\$246	\$178
Levemir	\$300	\$300	\$300
Lantus	\$226	\$221	\$206




Afrezza – Inhaled Insulin – Approved 2014



For adults over 18
Not indicated for pregnancy, while breastfeeding

Figure 3. Baseline-Corrected Glucose Infusion Rate (A) and Baseline-Corrected Serum Insulin Concentrations (B) after Administration of AFREZZA or Subcutaneous Insulin Lispro in Type 1 Diabetes Patients*



* Despite the faster absorption of insulin (FI) from Afrezza, the onset of activity (OT) was comparable to insulin lispro.



Diabetes Education SERVICES

Bolus Insulins (½ of total daily dose ÷ meals)

Name	Onset	Peak Action
▶ Lispro (Humalog)	15-30 min	1-1.5 hrs
▶ Aspart (NovoLog)		
▶ Glulisine (Apidra)		
▶ Afrezza (Inhaled)		
▶ Regular	30 mins	2-4 hrs



Diabetes Education SERVICES

Afrezza Dosing and Considerations


- ▶ Bolus regular insulin – inhaled before meals
- ▶ Dosing: 4 and 8 unit cartridges
 - ▶ Convert with 1:1 ratio to existing insulin dose
- ▶ Lung function test before start (incentive spiro)
 - ▶ Not for pts w/ chronic lung issues
 - ▶ Asthma, COPD, history of lung cancer, smokers
 - ▶ Can cause acute bronchospasm
- ▶ Side effects:
 - ▶ Hypoglycemia, sore throat, cough



Diabetes Education SERVICES

Afrezza Inhaler

Know your AFREZZA® inhaler:




Replace inhaler every 15 days



Diabetes Education SERVICES

Afrezza – Foil Packages Contain 30 cartridges – Use w/in 10 days

There are two strengths of AFREZZA® cartridges:



Let cartridges and inhaler sit at room temp for 10 minutes before using




Diabetes Education SERVICES

Afrezza– Storage

Storage:

- ▶ Refrigerated - Not in use and sealed –till expires
- ▶ Foil package at room temp – use within 10 days
- ▶ Once strips opened, good for 3 days



1 foil package

2 blister cards per package (5 strips per blister card)

3 cartridges per strip



Diabetes Education SERVICES

If you cannot find your injected dose in the chart below, you must talk to a healthcare provider before using AFREZZA®.

To switch from injected mealtime insulin to AFREZZA®...

Afrezza – Combos to get right dose

Find your injected insulin dose in the chart.

Injected Mealtime Insulin Dose	AFREZZA® Dose	# of 4 unit (blue) cartridges needed	# of 8 unit (green) cartridges needed
up to 4 units	4 units	1	0
5-8 units	8 units	0	1
9-12 units	12 units	1	1
13-16 units	16 units	2	0
17-20 units	20 units	2	1
21-24 units	24 units	3	0

Example: If you need to take 12 units of AFREZZA® you can use... 1 blue (4 unit) cartridge + 1 green (8 unit) cartridge

Find the correct dose of AFREZZA®.

Afrezza – Loading Cartridge into device

- ▶ Hold inhaler level

- ▶ Open inhaler by lifting white mouthpiece

- ▶ Hold insulin cartridge with cup facing down.

- ▶ Place cartridge inside and close lid. Keep level.

- ▶ Make sure cartridge has been at room temp for 10 minutes

Afrezza – Proper Inhale Technique

- ▶ Exhale

- ▶ Position inhaler in mouth (take off cover)

- ▶ Tilt inhaler down toward chin, keep head level
- ▶ Inhale deeply and hold breath for as long as comfortable

- ▶ Remove cartridge
- ▶ Replace cover

Bolus Insulin Summary

- ▶ Regular, Novolog, Humalog, Apidra, Afrezza
- ▶ Starts working fast (15-30 mins)
- ▶ Gets out fast (3-6 hours)
- ▶ Post meal BG reflects effectiveness
- ▶ Should comprise about ½ total daily dose
- ▶ Covers food or hyperglycemia.
- ▶ 1 unit
 - ▶ Covers ≈ 10 -15 gms of carb
 - ▶ Lowers BG ≈ 30 – 50 points



Diabetes Education SERVICES

Bolus Insulin Timing

- ▶ How is the effectiveness of bolus insulin determined?
 - ▶ 2 hour post meal (if you can get it)
 - ▶ Before next meal blood glucose
- ▶ Glucose goals (ADA) – may be modified by provider/pt
 - ▶ 1-2 hours post meal <180
 - ▶ Before next meal – 70 - 130



Diabetes Education SERVICES

Bolus – Insulin Sliding Scale

Starts at 150, 2 units for every 50 mg/dl >150

	Break	Lunch	Dinner	HS
Day 1	94 no insulin	212 4 uR	148 no insulin	254 6 uR
Day 2	243 4uR	254 6 uR	201 4uR	199 no insulin
Day 3	189 2uR	243 4uR	162 2uR	244 4uR
Day 4	66 No insulin	287 6uR	144 none	272 6uR



Diabetes Education SERVICES

Basal Insulins
(½ of total daily dose)

Intermediate Acting Peak Action Duration

▶ NPH 4-12 hrs 12-24

Long Acting Peak Action Duration

▶ Detemir (Levemir) peakless 20 hrs

▶ Glargine (Lantus) No peak 24 hrs

Fasting BG reflects efficacy of basal



Diabetes Education SERVICES

Basal Insulin Summary

- ▶ NPH, Levemir, Lantus
- ▶ Covers in between meals, through night
- ▶ Starts working slow (4 hours)
- ▶ Stays in long (12-24 hours)
 - ▶ NPH/ Lente 12 hrs
 - ▶ Levemir, Lantus 20-24 hrs
- ▶ Fasting blood glucose reflects effectiveness



Diabetes Education SERVICES

Basal Only
Type 2, 60kg – A1c 8.7%

	Break	Lunch	Dinner	HS
Mo 1	170s			298 10uLan
Mo 2	160s			233 20uLan
Mo 3	140s	283	265	206 30uLan



Diabetes Education SERVICES

Pattern Management



Diabetes Education SERVICES

Pattern Management

- ▶ Safety 1st!! - Evaluate 3 day patterns
- ▶ **Hypo:** eval 1st and fix:
 - ▶ If possible, decrease medication dose
 - ▶ Timing of meals, exercise, medications
- ▶ **Hyperglycemia:** evaluate 2nd
 - ▶ Identify patterns
 - ▶ Before increase insulin, make sure not missing something (carbs, exercise, omission)



Diabetes Education SERVICES

Type 2 – New diagnosis – No meds Patterns? Questions

	Break	Lunch	Dinner	HS
Day 1	164			181
Day 2		124	106	195
Day 3	149		102	242
Day 4	151	81		211



Diabetes Education SERVICES

Type 2 – Amaryl 4mg AM, 10u Lantus pm

	Break	Lunch	Dinner	HS
Day 1	164	94	66	162
Day 2	169		59	195
Day 3		84	81	242
Day 4	159		43	211



Diabetes Education SERVICES

Basal Bolus – What Adjustments?
Pt weighs 80kg

	Break	Lunch	Dinner	HS
Day 1	69 7H	79 5H	245 8H	190 22u Det
Day 2	81 7H	87 5H	170 8H	133 22u Det
Day 3	73 7H	94 5H	194 8H	110 22u Det
Day 4	62 7H	83 5H	211 8H	127 22u Det



Diabetes Education SERVICES

Intensive Diabetes Therapy
Insulin Dosing Strategy

50/50 Rule

▶ 0.5-1.0 units/kg day

▶ Basal = 50% of total

- Glargine QD
- NPH or Detemir BID

● Bolus = 50% of total

- usually divided into 3 meals

Example

▶ Wt 50kg x 0.5 = 25 units of insulin/day

▶ Basal dose: 13 units

- Glargine 13 units QD
- NPH/Detemir 6u BID

▶ Bolus dose: 12 units

- ▶ 4 units NovoLog, Apidra Humalog, Regular each meal



Diabetes Education SERVICES

Intensive Diabetes Therapy
Insulin Dosing Strategy

50/50 Rule

- ▶ 0.5-1.0 units/kg day
- ▶ Basal = 50% of total
 - Glargine QD
 - NPH or Detemir BID
- ★ Bolus = 50% of total
 - usually divided into 3 meals

Example – You Try

- ▶ Wt 60 kg x 0.5 = ____ units of insulin/day
- ▶ Basal dose: ____ units
 - Glargine ____ QD
 - NPH/Detemir __ BID
- ▶ Bolus dose: ____ units
 - ____units NovoLog, Apidra
 - Humalog, Reg each meal



Diabetes Education SERVICES

Intensive Diabetes Therapy
Insulin Dosing Strategy

50/50 Rule

- ▶ 0.5-1.0 units/kg day
- ▶ Basal = 50% of total
 - Glargine QD
 - NPH or Detemir BID
- ★ Bolus = 50% of total
 - usually divided into 3 meals

Example – You Try

- ▶ Wt 60kg x 0.5 = 30 units of insulin/day
- ▶ Basal dose: 15 units
 - Glargine 15 QD or
 - NPH/Detemir 7u BID
- ▶ Bolus dose: 15 units
 - ▶ 5 NovoLog, Apidra, Humalog, Reg each meal



Diabetes Education SERVICES

Basal Bolus – Using 50/50 Rule - Pt weighs 80kg

	Break	Lunch	Dinner	HS
Day 1	84 6H	89 7H	145 7H	190 20 u Det
Day 2	81 6H	97 7H	107 7H	133 20u Det
Day 3	79 6H	104 7H	124 7H	110 20u Det
Day 4	69 6H	103 7H	208 7H	193 20u Det



Diabetes Education SERVICES

Insulin Teaching Keys

- ▶ Bolus insulin with meals
- ▶ Basal 1-2xs daily
- ▶ Abdomen preferred injection site
- ▶ Stay 1" away from previous site
- ▶ Don't re-use ultra fine syringes
- ▶ Keep unopened insulin in refrigerator
- ▶ Toss opened insulin vial after 28 days
- ▶ Proper disposal
- ▶ Review patients ability to withdraw and inject.
- ▶ Side effects include hypoglycemia/wt gain
- ▶ Insulin pens –
 - ▶ Prime needle to assure accurate insulin dose given
 - ▶ Hold needle in for 5 seconds after injection
 - ▶ Roll 70/30 pens



Diabetes Education SERVICES

Sharps Disposal: Product and Info

- ▶ Look in the Government section white pages for a household hazardous waste listing for your city or county.
- ▶ Call 1-800-CLEANUP (1-800-253-2687)
- ▶ Search for collection centers on the California Integrated Waste Management Board (CIWMB) Web site:
<http://www.ciwmb.ca.gov/HHW/HealthCare/Collection/>



Diabetes Education SERVICES

DiaBingo - I

- I Injected hormone that is an analog of amylin
- I Glargine, Detemir, NPH are types of
- I Breakdown of glycogen into glucose
- I Anabolic hormone
- I Insulin is released when glucose levels are low
- I Once opened, insulin vials are good for one _____
- I Elevated post-prandial glucose indicate need for pre-meal
- I Epinephrine increases insulin resistance
- I Creation of glucose from amino acids and lactate
- I Decreasing renal function for people on insulin can cause
- I Bolus insulins
- I A hormone that increases blood glucose levels



Diabetes Education SERVICES

Diabetes Meds for Type 2: Objectives



1. Describe the main action of the 5 different categories of type 2 diabetes medications.
2. Discuss strategies to determine the right medication for the right patient.
3. List the side effects and clinical considerations of each category of medication.



Diabetes Education
SERVICES

Resources for Medications

- ▶ Partnership for Prescription Assistance
 - ▶ www.pparx.org
- ▶ NeedyMeds.org
- ▶ www.rxassist.org



Diabetes Education
SERVICES

Diabetes Agents Considerations

- ▶ Diabetes medications can be used as monotherapy, in combo or with insulin
- ▶ Combining agents from different classes has additive effect
- ▶ Most reduce A1c 0.5 – 2.0%
- ▶ Not to be used during preconception, pregnancy or when breastfeeding



Diabetes Education
SERVICES

ADA-EASD Position Statement:
Management of Hyperglycemia in T2DM

Patient-Centered Approach

“...providing care that is respectful of and responsive to individual patient preferences, needs, and values - ensuring that patient values guide all clinical decisions.”

- Gauge patient’s preferred level of involvement.
- Explore, where possible, therapeutic choices.
- Utilize decision aids.
- Shared decision making – final decisions re: lifestyle choices ultimately lie with the patient.



Diabetes Care 2012;35:1364–1379
Diabetologia 2012;55:1577–1596



Diabetes Education SERVICES

Approach to management of hyperglycemia:




Figure 1

Diabetes Care 2012;35:1364–1379
Diabetologia 2012;55:1577–1596
(Adapted with permission from: Ismail-Beigi et al. *Ann Intern Med* 2011;154:554)

Ideal Diabetes Med -

- ▶ No hypoglycemia
- ▶ No weight gain
- ▶ Affordable
- ▶ Lowers CV risk
- ▶ Most people can tolerate /use?



Diabetes Education SERVICES

Action/Classes of Type 2 Meds

- | | |
|------------------------|--|
| 1. Suppressor | Biguanide – Metformin |
| 2. Squirter | Sulfonylureas
Meglitinides |
| 3. Satiators | AmylinoMimetics
Incretin Mimetics
DPP-4 Inhibitors |
| 4. Sensitizer | Thiazolidinediones (TZD) |
| 5. Glucoretics | SGLT2 Inhibitors |
| 6. Circadian Switchers | Dopamine Receptor
Agonists |
| 7. Slower | Alpha-glucosidase inhibitors |



Diabetes Education SERVICES

Biguanides – Suppressor Metformin (Glucophage®)

- ▶ Action: suppresses release of glycogen from the liver
- ▶ Who?
 - ▶ Fasting hyperglycemia
 - ▶ Dysmetabolic Syndrome
 - ▶ For pediatrics starting age 10
 - ▶ (XR age 17)



Glycogen
Stopper



Diabetes Education SERVICES

Biguanides - Metformin

- ▶ **Action:** decrease hepatic glucose (glycogen)
- ▶ **Names:**
 - ▶ Metformin (Glucophage)
 - ▶ Starting dose: 500 BID, max 2500mg daily
 - ▶ Metformin XR - extended release – less GI upset
 - ▶ Starting dose 500mg at dinner, max dose 2000 to 2500 mg daily
- ▶ **Efficacy:**
 - ▶ Decrease fasting plasma glucose 60-70 mg/dl
 - ▶ Reduce A1C 1.0-2.0%



© Copyright 1999-2013, Diabetes Educational Services, All Rights Reserved

Diabetes Education SERVICES

Biguanides - Metformin

▶ Benefits

- ▶ Decrease LDL cholesterol and triglycerides
- ▶ No weight gain, possible modest weight loss
- ▶ Cancer protective?

▶ Concerns

- ▶ Diarrhea and abdominal discomfort – Use XR
- ▶ Lactic acidosis if improperly prescribed
- ▶ Watch for B12 deficiency
- ▶ Hold prior to IV contrast dye studies and use caution during acute illness. Resume when kidney function adequate



Diabetes Education SERVICES

Considerations

Biguanide - Metformin (Glucophage®)

▶ Contraindications due to lactic acidosis:

- ▶ creatinine >1.4 females, >1.5 males
- ▶ liver disease
- ▶ alcohol abuse
- ▶ over 80 years old
- ▶ risk of acidosis
- ▶ during IV dye study
- ▶ CHF requiring meds



© Copyright 1999-2013, Diabetes Educational Services, All Rights Reserved.

Diabetes Education SERVICES

Metformin – How does it rate?

<u>Question</u>	<u>Answer</u>
▶ Cause hypoglycemia?	No
▶ Cause weight gain?	No
▶ Affordable?	Yes
▶ Lowers CV risk?	Yes
▶ Can most tolerate /use?	Yes/No




Diabetes Education SERVICES

What is next step?

69 year old male, BMI 25, on Metformin 1000mg BID.
AM glucose 120s, A1c 8.1%.
Creat 1.3



Diabetes Education SERVICES



Sulfonylureas –

- ▶ Action: tells pancreas to squirt insulin all day
- ▶ Who?
 - ▶ Lean type 2



Diabetes Education SERVICES

Sulfonylureas - Squirts

- ▶ Action: Increase endogenous insulin secretion
- ▶ Efficacy:
 - ▶ Decrease FPG 60-70 mg/dl
 - ▶ Reduce A1C by 1.0-2.0%
- ▶ Secondary failures: 5-10% shortly after initial response, many more later
 - ▶ Usually after 5 or more years of therapy due to natural history of DM 2



Diabetes Education SERVICES

Sulfonylureas: 2nd Generation

Generic	Trade	Duration
▶ Glyburide	Diabeta, Micronase, Glynase Prestabs	12-24 hrs
▶ Glipizide*	Glucotrol, Glucotrol XI	12-24 hrs
▶ Glimepiride	Amaryl	16-24 hrs



Diabetes Education SERVICES

Sulfonylureas

- ▶ Other Effects
 - ▶ Hypoglycemia
 - ▶ Weight gain
 - ▶ Cleared by kidney, use caution for pts with kidney problems
 - ▶ Generally the least expensive class of medication
 - ▶ Amaryl safest for those with CV Disease



Diabetes Education SERVICES

Squirters – How does they rate?

Question	Answer
▶ Cause hypoglycemia?	Yes
▶ Cause weight gain?	Yes
▶ Affordable?	Yes
▶ Lowers CV risk?	No
▶ Can most tolerate /use?	Yes/No



Diabetes Education
SERVICES

What Medications Cause Hypoglycemia?

- ▶ Insulin
- ▶ Sulfonylureas
- ▶ Meglitinides
- ▶ Or any combo medication that includes these



Diabetes Education
SERVICES

Hypoglycemia = “Limiting Factor”

- ▶ Defined as glucose of 70mg/dl or below
- ▶ 50% of episodes occur during the night
- ▶ Higher mortality rate with severe hypoglycemia secondary to sulfonylureas
 - ▶ Especially (glyburide) Micronase®, Diabeta®
- ▶ Blood glucose levels don’t describe severity, response is individual



Diabetes Education
SERVICES

Hypoglycemic Symptoms

- ▶ Autonomic
 - ▶ Anxiety
 - ▶ Palpitations
 - ▶ Sweating
 - ▶ Tingling
 - ▶ Trembling
 - ▶ Hypoglycemic Unawareness
- Neuroglycopenia
 - ↳ Irritability
 - ↳ Drowsiness
 - ↳ Dizziness
 - ↳ Blurred Vision
 - ↳ Difficulty with speech
 - ↳ Confusion
 - ↳ Feeling faint



Diabetes Education SERVICES

Treatment of Hypoglycemia

- ▶ If blood glucose **70mg/dl** or below:
 - 10-15 gms of carb to raise BG 30 - 45mg/dl
- Ⓞ Retest in 15 minutes, if still low, treat again, even without symptoms
- Ⓞ Follow with usual meal or snack
- Ⓞ If BG less than 40, allow recovery time



Diabetes Education SERVICES

15 - 20 Gms Carb Sources

- Ⓞ 3 - 4 Glucose Tablets
- Ⓞ 8 - 10 Lifesavers candy
- Ⓞ 8 - 10 Hard candies
- Ⓞ 2 Tablespoons Raisins
- Ⓞ 4 - 6 oz's Nondiet soda
- Ⓞ 4 - 6 oz's Fruit Juice
- Ⓞ 8 oz Milk (non fat)



Diabetes Education SERVICES

What questions?

- ▶ 72 yr old, thin, lives alone, A1c 7.3%. History of MI, stroke. DM for 12 yrs, “diet controlled”. Limited income. Creat 1.4.



Diabetes Education SERVICES

DPP-4 Inhibitors – “Incretin Enhancers”

Januvia (sitagliptin) – Tradjenta (linagliptin)
Onglyza (saxagliptin) Nesina (alogliptin)

▶ Action:

- ▶ Increase insulin release w/ meals
- ▶ Suppress glucagon

▶ Dosing:

Januvia – 100mg a day
Onglyza – up to 5mg a day
Tradjenta – 5mg a day
Nesina – up to 25 mg a day

▶ Efficacy: Decreases A1c by 0.6 -0.8%

▶ Indication: For type 2s



Diabetes Education SERVICES

DPP-4 Inhibitors – “Incretin Enhancers”

Januvia (sitagliptin) – Tradjenta (linagliptin)
Onglyza (saxagliptin) Nesina (alogliptin)

- ▶ Januvia, Onglyza eliminated via kidney, lower dose needed
- ▶ Do not cause wt gain or hypoglycemia
- ▶ Side effects – headache, runny nose, sore throat - watch for pancreatitis
- ▶ Cost \$100 - \$150 mo



Diabetes Education SERVICES

DPP-IV Inhibitors – How do they rate?


Question	Answer
▶ Cause hypoglycemia?	No
▶ Cause weight gain?	No
▶ Affordable?	No
▶ Lowers CV risk?	No
▶ Can most tolerate /use?	Yes



Diabetes Education SERVICES


Incretin Mimetics – “Gut Hormone Imitators” GLP-1 Agonists

▶ How do they work?



Diabetes Education SERVICES

GLP-1 Effects in Humans Understanding the Natural Role of Incretins



Adapted from Flint A, et al. J Clin Invest. 1998;101:513-520.
Adapted from Larsson HK, et al. Acta Physiol Scand. 1997;150:413-422.
Adapted from Nauck MA, et al. Diabetologia. 1995;38:1546-1553.
Adapted from Drucker DJ. Diabetes. 1998;47:189-193.

GLP-1 degraded by DPP-4 w/in minutes



Diabetes Education SERVICES

Incretin Mimetics

Exenatide (Byetta), Exenatide XR (Bydureon)

▶ **Action:**

- ▶ Insulin release in response to meal
- ▶ Slows gastric emptying
- ▶ Causes Satiety
- ▶ Preserves Beta Cells

▶ **Exenatide Dosing:**

- ▶ 5-10 mcg before break, dinner
- ▶ Long acting version - 1x week (available in pens in 2015)

▶ **Efficacy:** Decreases A1c by 0.7%, wt by 3lbs

▶ **Indication:** For type 2s only - mono or in combo



Diabetes Education SERVICES

Incretin Mimetics —

Exenatide XR - Bydureon

▶ **Once a Week Dosing:** 2mg

▶ **Efficacy:** Decreases A1c by 1.6%, wt by ~6lbs

▶ **Indication:** For type 2s only

▶ **Other:** Pt will need to mix powdered form and inject – Pen in future

▶ **Caution:** not indicated for those with history of medullary thyroid tumor - pancreatitis warning



Diabetes Education SERVICES


\$323.44 for four doses, or about \$4,200 a year.



Diabetes Education SERVICES

Incretin Mimetics – Albiglutide - Tanzeum

- ▶ **Once a Week Dosing:** 30 – 50mg
- ▶ **Efficacy:**
Decreases A1c by ~ 1%, wt by ~2lbs
- ▶ **Indication:** For type 2s only
- ▶ **Other:** Pen injector
- ▶ **Caution:** not indicated for those with history of medullary thyroid tumor - pancreatitis warning



Incretin Mimetics - GLP-1 Analog Liraglutide (Victoza)

- Liraglutide Dosing:** 1x daily, time not critical
- 0.6 x 1 week – if tolerated (nausea), go to >
 - 1.2 x 1 week – if tolerated go to >
 - 1.8 mg daily
 - ▶ **Efficacy:** lowers; A1c by 1%, body wt by ~ 2.5kg
 - ▶ **Indication:** Monotherapy or in combo . Type 2 only
 - ▶ **Other:** In pen, with preset dosing
 - ▶ **Black box**–thyroid tumor warning (avoid if family hx, notify MD of hoarseness, lump).



For all the Previous GLP-1 Agonists

- **Pancreatitis Warning**
 - Please tell all patients to report signs right away and discontinue meds
 - Signs include:
 - Sudden abdominal pain, nausea and vomiting



Incretin Mimetics – How do they rate?

Question	Answer
▶ Cause hypoglycemia?	No
▶ Cause weight gain?	No
▶ Affordable?	No
▶ Lowers CV risk?	No
▶ Can most tolerate /use?	Yes/No (GI)



Diabetes Education SERVICES

Next Step?

- ▶ 69 year old male, BMI 25, on Metformin 1000mg BID and Exenatide 10mcg before breakfast and dinner.
- ▶ Pt overweight - A1c 8.1%. Creat 1.2



Diabetes Education SERVICES

SGLT2 Inhibitors- “Glucoretics”

▶ **Action:** “Glucoretic” decreases renal reabsorption in the proximal tubule of the kidneys (reset renal threshold and increase glucosuria)

▶ **Names:**

- ▶ Canagliflozin (Invokana)
- ▶ Dosing: 100 – 300 mg once daily ac first meal
 - ▶ If eGFR 45-60: do not exceed 100mg a day
 - ▶ If eGFR <45, do not use



Decreases Glucose Reabsorption

- ▶ Dapagliflozin (Farxiga)
- ▶ Dosing: 5 – 10 mg once daily ac first meal
 - ▶ If eGFR <60, do not use
 - ▶ Don't use if pt has bladder cancer and report blood in urine

▶ **Efficacy:**

- ▶ Weight loss of 1-3 lbs
- ▶ Reduce A1C ~0.7-1.5%



Diabetes Education SERVICES

Considerations



- Monitor B/P, K+ & renal function.
- Side effects: hypotension, UTI, increased urination, genital yeast infections.
- Improves beta cell function?
 - Reverses glucoses toxicity by increasing GLUT4 transport in muscle
 - Increase liver sensitivity to insulin and decreases gluconeogenesis.



Diabetes Education SERVICES

SGLT2 Inhibitors- How do they rate?

Question	Answer
▶ Cause hypoglycemia?	No
▶ Cause weight gain?	No
▶ Affordable?	No
▶ Lowers CV risk?	No
▶ Can most tolerate /use?	Yes?



Diabetes Education SERVICES

Indications for Insulin Sensitizers

Rosiglitazone (Avandia), Pioglitazone (Actos)

- ▶ **Action:** decrease insulin resistance by making muscle and adipose cells more sensitive to insulin. Decrease free fatty acids
- ▶ **Names:**
 - ▶ pioglitazone (Actos) – bladder cancer warning
 - ▶ Dosing: 15-45 mg daily
 - ▶ rosiglitazone (Avandia) – restriction relaxed
 - ▶ Dosing: 4-8 mg daily
- ▶ **Efficacy/ Considerations**
 - ▶ Reduce A1C ~0.5-1.0%
 - ▶ 6 weeks for maximum effect
 - ▶ \$100 a month
 - ▶ Can cause fluid retention, not indicated w/ CHF



Diabetes Education SERVICES

TZDs – How do they rate?

Question	Answer
▶ Cause hypoglycemia?	No
▶ Cause weight gain?	Yes
▶ Affordable?	Yes/No
▶ Lowers CV risk?	??
▶ Can most tolerate /use?	Yes/No



Diabetes Education SERVICES

Keeping Healthy in Long Run

- ▶ HUG Patients
 - ▶ Help with
 - ▶ Unconditional
 - ▶ Guidance and Support
- Anne Peters, MD, CDE
ADA Post Grad

▶ **Unconditional Positive Regard –**
involves showing complete support and acceptance of a person no matter what that person says or does.
Carl Rogers



Diabetes Education SERVICES

Thank You



- ▶ Questions?
- ▶ Email
bev@diabetesed.net
- ▶ Web
www.DiabetesEd.net



Diabetes Education SERVICES
