



Thank you for downloading our PocketCards! We found these resources helpful so we want to share them with you!

- **FREE Live Webcasts** - check here to see our upcoming schedule
- **FREE Resource Pages** -
 - **Preparing for the CDE Resource Page and important articles**
 - **Preparing for the BC-ADM Resource Page and important articles**
- **Comprehensive Diabetes Articles Library**
- **Send for your free laminated copies of the PocketCards**
- **Monthly Newsletters** - Review our archived editions
- **Instructional videos** - on how to use our Teaching Tools

Thanks again for your efforts to improve diabetes care. To see and print your PocketCards, scroll to the next page.

www.DiabetesEd.net



Advancing Your Career
in Diabetes Education

Connect with us online!

Main Website | Online University | Online Store | Facebook



Oral Diabetes Medications

Class/Main Action	Name(s)	Daily Dose Range	Considerations
Biguanides <ul style="list-style-type: none"> Decrease hepatic glucose output First line med at diagnosis of type 2 	metformin (Glucophage)	500 – 2500 mg (usually BID w/meal)	Side effects: nausea, bloating, diarrhea. Use XR to minimize. Lactic acidosis precaution: avoid in pts with creat >1.4 women, 1.5 men, during illness or surgery. Benefits: decreased cholesterol, no wt gain or hypoglycemia. Lowers A1c 1.0% – 2.0%.
	Extended Release-XR (Glucophage XR) (Glumetza) (Fortamet)	(1x daily w/dinner) 500 – 2000 mg 500 – 2000 mg 500 – 2500 mg	
Sulfonylureas <ul style="list-style-type: none"> Stimulates sustained insulin release 	glyburide: (Micronase, Diabeta) (Glynase)	1.25 – 20 mg 0.75 – 12 mg	Can take once or twice daily before meals. Side effects include hypoglycemia and weight gain. Eliminated via kidney. Caution: Glyburide most likely to cause hypoglycemia. Lowers A1c 1.0% – 2.0%.
	glipizide: (Glucotrol) (Glucotrol XL)	2.5 – 40 mg 2.5 – 20 mg	
	glimepiride (Amaryl)	1.0 – 8 mg	
DPP – 4 Inhibitors <ul style="list-style-type: none"> "Incretin Enhancers" Prolongs action of gut hormones Increases insulin secretion Delays gastric emptying 	sitagliptin (Januvia)	100 mg daily (eliminated via kidney*)	*If creatinine elevated, see pkg insert for dosing info. No wt gain or hypoglycemia. Side effects include nasopharyngitis, headache and upper-respiratory tract infection. Report signs of pancreatitis (abdominal pain, nausea, vomiting). Lowers A1c 0.6% – 0.8%.
	saxagliptin (Onglyza)	Up to 5 mg daily (eliminated via kidney*, feces)	
	linagliptin (Tradjenta)	5 mg daily (eliminated via feces)	
	alogliptin (Nesina)	25 mg once daily (eliminated via kidney)	

More medications on back. Note: These meds are for people with Type 2 diabetes and should not be used during pregnancy. Content is for educational purposes only; please consult prescribing information for details.

Class/Main Action	Name(s)	Daily Dose Range	Considerations
SGLT2 Inhibitors <ul style="list-style-type: none"> Decrease glucose reabsorption in kidneys "Glucretic" 	Canagliflozin (Invokana) Dapagliflozin (Farxiga) Empagliflozin (Jardiance)	100 – 300 mg 1x daily 5 – 10 mg 1x daily 10 – 25 mg 1x daily	For all, monitor B/P, K+ and renal function. If GFR<45, stop Invokana. If GFR<60, stop Farxiga. Do not start pts w/ GFR<45 on Jardiance. Side effects: hypotension, UTIs, increased urination, genital infections. Avoid Farxiga in pts w/ bladder cancer. Lowers A1c 0.7% – 1.5%, lowers wt 1 – 3 lbs.
Thiazolidinediones "TZDs" <ul style="list-style-type: none"> Increase insulin sensitivity 	pioglitazone (Actos) rosiglitazone (Avandia)	15 – 45 mg daily 4 – 8 mg daily	Black Box Warning: TZDs may cause or worsen CHF. Monitor for edema and weight gain. Increased peripheral fracture risk. Actos may increase risk of bladder cancer. Lowers A1c 0.5% – 1.0%
Glucosidase Inhibitors <ul style="list-style-type: none"> Delay carb absorption 	acarbose (Precose) miglitol (Glyset)	25 – 100 mg w/meals; 300 mg max daily dose	Start low dose, increase at 4-8 wk intervals to decrease GI effects. Caution with liver or kidney problems. In case of hypo, treat w/ glucose tabs. Lowers A1c 0.5– 1.0%.
Dopamine Receptor Agonists <ul style="list-style-type: none"> Resets circadian rhythm 	bromocriptine mesylate— Quick Release "QR" (Cycloset)	1.6 to 4.8 mg a day (each tab 0.8 mg)	Take within 2 hrs of waking. Side effects: nausea, headache, fatigue, hypotension, syncope, somnolence. Lowers A1c 0.6% – 0.9%.
Meglitinides <ul style="list-style-type: none"> Stimulates rapid insulin burst 	repaglinide (Prandin) nateglinide (Starlix)	0.5 – 4 mg w/meals (metabolized in liver) 60 – 120 mg w/meals (eliminated via kidney)	Take before meals. Side effects may include hypoglycemia and weight gain. Lowers A1c 1.0% – 2.0%.